Sky and TELESCOPE



Total eclipse and shadow

In This Issue:

Eclipse over New England

Reflections of a Spectroscopist

The Great Aurora of Early September

American Astronomers Report

Philipp Fauth and the Moon

Radio Astronomy Receivers — I

Northern and Southern Star Charts

★

Vol. XIX, No. 1

NOVEMBER, 1959

50 cents

★



j. w. fecker inc.

A subsidiary of American Optical Company

6592 HAMILTON AVENUE, PITTSBURGH 6, PA.

Sky and TELESCOPE

CHARLES A. FEDERER, JR., *Editor*
JOSEPH ASHBROOK, *Technical Editor*

Copyright, 1959, by SKY PUBLISHING CORPORATION, all rights reserved.

Vol. XIX, No. 1 NOVEMBER, 1959

CONTENTS

COVER: The October 2nd total eclipse of the sun, as seen above the eastern horizon from an American Airlines plane at 21,000 feet, 16 miles off the Massachusetts coast. Surrounding the sun is the silvery corona, showing the relatively round form typical in years when sunspots are most numerous. The shadow of the moon on the atmosphere is conspicuous in this picture, the bright clouds at the right and left being on the edges of the narrow cylinder of totality. Photograph by Edward F. Carr, courtesy of the Boston "Globe." (See page 4.)

SOME NEW AMERICAN RADIO TELESCOPES 3

ECLIPSE OVER NEW ENGLAND 4

REFLECTIONS OF A SPECTROSCOPIST
— Otto Struve 7

THE GREAT AURORA OF EARLY SEPTEMBER 14

AMERICAN ASTRONOMERS REPORT 16

PHILIPP FAUTH AND THE MOON
— Hermann Fauth 21

RADIO ASTRONOMY RECEIVERS – I
— Frank D. Drake 26

AMATEUR ASTRONOMERS 29
A Young Amateur's Observatory in Texas

BOOKS AND THE SKY 44
Handbuch der Physik, Vol. 53; Astrophysics IV: Stellar Systems
Films and the Sky — The Story of the Universe, Unit I: The Earth and Its Moons

CELESTIAL CALENDAR 60

GETTING ACQUAINTED WITH ASTRONOMY 25
Techniques of Lunar and Planetary Observing — II

GLEANINGS FOR ATM's 51
A Low-Power Microscope System for High Magnifications
Some Advantages of a Transfer Lens

NEWS NOTES 19

OBSERVER'S PAGE 32
Observing the Moon — Parry and Bonpland
Astronomical Seeing
Deep-Sky Wonders

OBSERVING THE SATELLITES 11

PLANETARIUM NOTES 30

QUESTIONS 24

SOUTHERN STARS 64

STARS FOR NOVEMBER 65

SKY AND TELESCOPE is published monthly by Sky Publishing Corporation, Harvard College Observatory, Cambridge 38, Mass. Second-class postage paid at Boston, Mass.

Subscription: $5.00 per year in the United States and possessions; $9.00 for two years; $13.00 for three years. To Canada, Mexico, and all countries of the Pan American Postal Union, $6.00 for one year; $11.00 for two years; $16.00 for three years. To all other foreign countries, $7.00 for one year; $13.00 for two years; $19.00 for three years. Canadian and foreign remittances should be made in United States currency. Single copy 50 cents. Back numbers, as available, 50 cents each.

All notices of change of address must be sent one month in advance and accompanied by old and new addresses. When sending your renewal order, or writing in regard to your subscription, your current mailing address must be given. Return our bill form with renewal payment. Circulation manager, Nancy R. Bolton. Office staff: Janet G. Condon, Virginia K. Cox, J. Caroline Nason, Mary Silva, John Simmons, Helen B. Sniegiecki, Stacia Strong.

Editorial and advertising offices: 49-50-51 Bay State Rd., Cambridge 38, Mass. Advertising rates are listed in STANDARD RATE AND DATA SERVICE, or sent upon request. Advertising manager, Henry M. Corrado. Editorial staff: Francis B. Russell, William E. Shawcross, Donald B. Silverman.

Unsolicited articles and pictures are welcome, bearing adequate return postage, but we cannot guarantee prompt editorial attention, nor are we responsible for the return of unsolicited material. The principal articles are indexed in THE READERS' GUIDE TO PERIODICAL LITERATURE.

Although the contents of this magazine, its text, pictures, diagrams, and charts, are copyrighted with all rights reserved, we are pleased to give permission, upon request, for the use of SKY AND TELESCOPE material for educational and scientific purposes, with the exception of the monthly star and planet charts, which may not be copied or reproduced in any manner for any purpose. Permission to use matter from signed articles, and photographs and diagrams for which credit is given, must also be obtained from the person, observatory, institution, or agency concerned.

Some New American Radio Telescopes

TO many astronomers, the 1950's will be remembered as the decade of an explosively rapid expansion in radio astronomy, when nearly every branch of the science was invigorated by the findings with giant instruments of varied designs. An even faster growth in the near future is promised by additional large radio telescopes just completed or under construction in this country.

On October 7th, the University of Michigan dedicated the new 85-foot paraboloidal radio telescope at its Peach Mountain station, 16 miles northwest of Ann Arbor. Primarily intended for solar research, the 155-ton instrument joins a 28-foot antenna that has been at work there since August, 1957. Being fully steerable, the equatorially mounted 85-foot dish can track the sun from rising to setting.

This instrument will also be used for 21-centimeter and other studies of Milky Way and extragalactic sources. Many refinements are incorporated, including a newly developed broad-band traveling-wave-tube radiometer, representing a major advance over other similar receivers. It will permit observations to be made at wave lengths as short as three centimeters. Another feature is a ruby maser, a virtually noise-free amplifying device. The project has been supported by the Office of Naval Research.

Another 85-foot radio telescope is planned to be in operation by June, 1960, at the University of California's Hat Creek station in the Lassen National Forest, in northern California. The 60-acre tract, 14 miles southeast of the town of Burney, was selected after a five-month survey of 30 possible localities. A second paraboloidal antenna, 33 feet in diameter, is expected to be in use at the same site next month. Harold F. Weaver is director of the new radio observatory, which will cost about $500,000. ONR is providing 70 per cent of the amount, and will share the operating expenses with the University of California.

Much larger than these instruments is the 140-foot dish under construction at the National Radio Astronomy Observatory at Green Bank, West Virginia. On page 26 of this issue, NRAO astronomer Frank D. Drake reviews the functions of radio receivers and explains the importance of radiometer design in the operation of radio telescopes.

Thirty airline miles away, at Sugar Grove, West Virginia, the Naval Radio Research Station is constructing the largest steerable radio antenna of all, a 600-foot paraboloid. Although it is designed primarily for military communications research, it will also be used for basic astronomical work.

ECLIPSE OVER NEW ENGLAND

This series of eclipse pictures was taken from an airplane by David H. Freeman, Massachusetts Institute of Technology, aided by John S. Robie, Lexington, Massachusetts, and John A. Griner, Harvard Business School. The plane was flying some 120 miles an hour at 6,500 to 6,800 feet when these pictures were made.

Dr. Freeman used two Leica cameras, with 135-mm. lenses and ultraviolet filters. One camera, containing Kodachrome film, was preset for the partial phases, and secured the topmost picture, of the thin rising crescent before totality. The other camera, with high-speed Ektachrome (ASA 160), took the totality shots, the diamond-ring effect, and the early partial phase (bottom).

AT MOST a few hundred observers, borne aloft in airliners and small private planes, saw the total phase of October 2nd's dawn eclipse in the United States. Meanwhile rain, fog, and cloudy skies disappointed many thousands of ground observers who flocked to shore sites north of Boston and inland vantage points in Massachusetts and southern New Hampshire.

By 9 o'clock that morning the sun was shining brightly at Salem, where 500 observers had gathered for the SKY AND TELESCOPE eclipse party at the Coast Guard Air Rescue Station and the Plummer Home for Boys. But at 6:50 a.m., Eastern daylight time, when the moon's shadow briefly touched New England, these travelers from 25 states and five foreign countries were standing in a dismal rain with visibility less than a mile. The remnant of Hurricane Gracie, passing northward across New York State and western New England, was too slow in moving out of the eclipse area. No report has been received that any ground observer in North America saw the total phase.

Photography was possible only from the air. Probably the highest observer was J. Allen Hynek, Smithsonian Astrophysical Observatory, who was in a jet trainer



Observing near Lynn, Massachusetts, Armand N. Spitz, of Yorklyn, Delaware, was able to secure this record of the changes in zenith brightness on eclipse morning. He used a Weston photoelectric photometer, making observations every five minutes until near totality. The sudden darkening experienced by ground observers is shown by the curve, as well as the rapid increase in brightness thereafter. When the partial eclipse was over, the heavy overcast gave a reading of about 45 foot-candles.

A somewhat globular corona is shown by this picture, taken from an American Airlines plane at 21,000 feet by David Wurzel, New England picture manager of United Press International Photos. He used a 4-by-5 Speed Graphic with a 15-inch lens, exposing 1/125 second on Royal Pan film with no filter. Note the reflection of coronal light from the layer of clouds.

at 35,000 feet, where the eclipse was "perfect, except for a few little streaky clouds." Several airlines sent up large planes, carrying scientists and newspaper photographers, some of their pictures being shown here and on the front cover.

One amateur astronomer, David H. Freeman, a physical chemist at Massachusetts Institute of Technology, had chartered a plane a month ahead of time, only to find it on eclipse morning carefully stowed away in a hangar at Bedford airport, behind a large bomber! After some hasty searching, he found another plane for his small party. Piloted by Frank D. Comerford, an instrument flying instructor, the twin-engine Apache was among the few that succeeded in breaking through the storm into clearing skies a mile or so high. As totality came, they saw the moon's shadow racing across the cloud layer below them.

The city of Boston, within the totality zone, was plagued by early morning traffic jams as tens of thousands of would-be observers sought vantage points. Large numbers crowded Lynn Beach, where the

About 25 times as many observers as in this picture attended the eclipse party at Salem, Massachusetts. Here they gaze forlornly across Salem Harbor past the lighthouse at old Ft. Pickering, while the rain comes down. Marblehead is in the distance. Much photographic and telescopic equipment was set up here, including an elaborate array by the Grumman Astronomical Society of Bethpage, New York. United Press International photographs.

Above: The total eclipse seen from a Northeast Airlines plane at 16,000 feet by Frank Curtin, an Associated Press photographer. He used a Hasselblad camera with a 180-mm. lens. Wide World photograph.

Right: A multiple exposure of four phases of the partial eclipse, taken atop the R. C. A. building in New York City. Exposures were made at five-minute intervals from 7:25 a.m. (bottom to top), 1/1,000 second at f/22 with an orange filter. Wide World photograph.

Amateur Telescope Makers of Boston had their official station, and thronged to Winthrop, Revere, and Cape Ann. On Nahant, the American Association of Variable Star Observers set up their instruments within the grounds of the U. S. Army's Nike rocket base.

Inland, at Mt. Wachusett, observing teams from the Detroit Astronomical Society, the Smithsonian Astrophysical Observatory, the Boston Museum of Science, and the Air Force Cambridge Research Center, were stationed. With hundreds of other watchers, their pre-dawn hopes were raised by breaks in the cloudy mantle, but during the crucial moments of totality the eastern horizon was completely obscured.

The party led by Peter A. Leavens, of Freeport, New York, arrived at Salem early, then decided to search for better weather. They found a vantage point near Nashua, New Hampshire. During totality, which was hidden by clouds, they saw the bright sky to the north outside of the shadow path.

Arthur Stockellburg, W1SS, of Lincoln, Massachusetts, was one of several amateurs who co-operated to make successful radio observations of the effects of the eclipse. He transmitted an unmodulated, constant-intensity 3,820-kilocycle signal, which was monitored at Westfield, Massachusetts; Ellsworth, Maine; Springfield, Vermont; and other points. The Westfield record showed a short-lived sharp rise in signal strength during totality; other stations reported a broad peak.

First reports from American observers outside the totality zone indicate that clouds also interfered with seeing the partial phases. In New York City, the best view came about 7:30 a.m., when some one-fourth of the sun's diameter was covered by the moon.

From early information, even parts of the highly favored Canary Islands had clouds, possibly affecting some expeditions adversely. Perhaps the most dramatic view of the eclipse was obtained by the crew of a jet F-101B Voodoo, which raced the moon's shadow for seven minutes from Fuerteventura Island to the African desert at a speed of 1,080 miles per hour. During the flight, a magnetic tape recorded measurements of polarization in the sun's corona, planned as a test, particularly in the infrared, of the presence of synchrotron radiation in the outermost solar atmosphere.

Reflections of a Spectroscopist

OTTO STRUVE

*National Radio Astronomy Observatory**

The dome of the 100-inch Hooker reflector at Mount Wilson Observatory as it looks today. In use since 1918, for three decades this was the world's largest telescope. Its enormous light grasp particularly suits it for spectroscopic work. The room of the high-dispersion coude spectrograph is located under the slanting roof, at the southern end of the polar axis.

SINCE my recent appointment to the staff of the National Radio Astronomy Observatory, I have been thinking a great deal about some of the questions raised at a symposium in Paris last July. There, under the same title as this article, P. Swings, of the University of Liège, made an eloquent appeal to young French scientists to develop an interest in stellar and nebular spectroscopy, in order to take advantage of the fine instrumental facilities now available to them at St. Michel Observatory.

As Professor Swings pointed out, "The study of spectra of the celestial objects seems to have lost some of its appeal during the past few years, due to the emergence of accurate photometric methods, of radio astronomy investigations, of semitheoretical applications of nuclear physics, and of such recent developments as plasma physics, magnetohydrodynamics, and shock waves."

Yet, he continued, a large number of purely spectroscopic problems remain unsolved or only partly solved. "Progress in our knowledge of the Milky Way and other galaxies will, perhaps forever, depend to a large extent upon special spectroscopic investigations. We would, in fact, not have been able to go very far in our theories of stellar evolution if it had not been for the patient and admirable spectroscopic researches carried out at Mount Wilson, Palomar, McDonald, Lick, Victoria, Haute Provence, and elsewhere."

Although in turning to radio astronomy I am giving up, or at any rate greatly reducing, my work in optical stellar spectroscopy, my decision is in no sense caused by loss of interest in this field. It is one of the most rewarding branches of scientific endeavor, and I would like to urge young astrophysicists to choose it for their careers.

My first exposure to the marvels of stellar spectra occurred just 40 years ago. As a student at the University of Kharkov in Russia, I was taking an astrophysics course given by B. P. Gerasimovich, who had just returned after participating for several years in A. A. Belopolsky's spectrographic investigations at Poulkovo Observatory. Dr. Gerasimovich had brought back with him a series of spectrograms of different stars. He used these to teach his students the technique of measuring radial velocities.

My studies were interrupted by the civil war in Russia, and I was only able to resume them after arriving at Yerkes Observatory in October, 1921, as a graduate student and assistant in stellar spectroscopy to director E. B. Frost. I remember vividly how he told me that the measurement of stellar radial velocities was the most exciting, gratifying scientific occupation he had ever known!

Forty years ago the application of Doppler's principle to astronomy was still fairly new. Many astronomers then living had personally known Huggins, Vogel, and other pioneers in the measurement of the line-of-sight motions of stars, and could recapture the almost mystical delight brought by this great scientific advance. Frost, Belopolsky, W. S. Adams, J. S. Plaskett, and W. W. Campbell belonged to that period. But to me, of a later generation, the excitement that these men felt was foreign: I recognized the great importance of determining radial

*Operated by the Associated Universities, Inc., under contract with the National Science Foundation.



The first observatory especially devoted to spectroscopic studies of heavenly bodies. From 1860 to 1869, William Huggins used this 8-inch Clark refractor at Tulse Hill, England, for visual observations of the spectra of stars and nebulae. Huggins in 1868 was the first to attempt to measure the line-of-sight velocities of stars. From H. C. King, "The History of the Telescope."

velocities of stars, but the application of Doppler's principle to astronomy seemed an old story.

Perhaps today many young astronomers regard the more recent developments of stellar spectroscopy in the same light. Yet, I have experienced the same kind of exhilaration as did Frost in his radial velocity work, for I have witnessed enormous advances, such as A. S. Eddington's proof that sharp absorption lines of ionized calcium and neutral sodium can be produced in interstellar space; M. N. Saha's theory of excitation and ionization, and its extension by H. N. Russell, E. A. Milne, and others; Cecilia Payne Gaposchkin's application of this theory to stellar spectra; A. Unsöld's derivation of abundances of chemical elements in the atmospheres of stars, and many more.

But this very feeling comes in other ways. Ever since obtaining my first stellar spectrogram with the 82-inch telescope at McDonald Observatory, I have not spent a night observing there, or during the past nine years at Mount Wilson, without experiencing excitement and anticipation. Never have I returned from an observing visit at either place without some new and usually unexpected features on some of my spectrograms. The predicted characteristics were not always there; the unexpected ones made every spectrogram a potential gold mine!

To achieve important results it is necessary, as Swings said, "to keep abreast of all technical advances, and all new theoretical developments. . . . The spectroscopist must first seek to increase the range of wave lengths he can cover, and to shorten the time required to record a spectrum by increasing the sensitivity of his receivers." Although many important spectroscopic problems can be attacked with relatively small reflectors and simple, medium- or small-dispersion spectrographs, the scope of the work may be so limited that frustration might destroy the investigator's initiative.

I experienced this frustration at Yerkes. The combination of the 40-inch refractor and a very old spectrograph, employing three 60-degree prisms of optically dense glass, allowed me to reach stars no fainter than 4th or 5th magnitude with a linear dispersion of 10 angstroms per millimeter. Today, a 60-inch parabolic reflector and a modern grating spectrograph, with a range of dispersions between 10 and perhaps 75 angstroms per millimeter, should normally be regarded as the minimum instrumental requirement, except for such spectral classification programs as those undertaken by W. W. Morgan and P. C. Keenan, or by D. Chalonge in France.

Edwin B. Frost (1866-1935), second director of Yerkes Observatory, devoted his entire career to stellar spectra. From his autobiography, "An Astronomer's Life," Houghton Mifflin Co., 1933.

Photographic procedures are still good for many problems, but the spectroscopist should try to develop other methods. B. Strömgren has been very successful in measuring the intensities of absorption lines using a photoelectric photometer with narrow-transmission-band filters. At Mount Wilson, a promising method of scanning spectra has been developed by L. Spitzer, Jr., and his collaborators, and a somewhat different technique by O. C. Wilson and J. B. Oke.

Spectroscopists can profitably exploit and improve the image-tube methods of A. Lallemand, W. A. Baum and J. S. Hall, or W. A. Hiltner. Because the image tube promises to reduce the time required for recording the spectrum of even a faint star to minutes or seconds, this device should be well suited for observing certain very rapid changes in stellar spectra. I have described some of these short-lived events in my article on Nova Herculis in the July issue, and the one on flare stars in September.

A problem of this nature that has intrigued me for many years is in connection with the famous eclipsing binaries U Cephei and U Sagittae, when only a narrow crescent of the early-type component is unhidden just before or after totality. My McDonald spectrograms show during these brief intervals extraordinary changes in the absorption lines of the star undergoing eclipse. But the exposure times were too long to record the changes without serious contamination by the light of the late-type subgiant producing the eclipse. The use of image-tube techniques should be especially profitable in this case.



The 82-inch reflector of McDonald Observatory on Mt. Locke, Texas, was completed in 1939. This fine instrument has been very effective for astrophysical observations. Note the counterweight close to the northern pier, as well as the observing platform in the background.

Professor Swings also recommends efforts to observe with reasonably high dispersion the infrared spectral region between one and 15 microns. In addition, he would like to see someone try M. Czerny's evaporograph technique, in which long-wave radiation is detected by its heating effect upon some highly volatile material, such as a very thin film of camphor. Others should attempt to bridge the gap between the optical infrared, which extends to about 15 microns, and the present short-wave limit of ground-based radio telescopes at about two or three centimeters. This will, of course, require a radio instrument located in a satellite above the ionosphere, a region which is opaque in the range of wave lengths mentioned.

Satellite instruments should soon be informing us of the appearance of stellar spectra at wave lengths shorter than 3000 angstroms. Preliminary calculations by L. Aller indicate, however, that in the far ultraviolet the absorption lines of the Lyman series of hydrogen and the Lyman continuum, both produced by interstellar atoms, may be so strong that even the nearest stars will be invisible at wave lengths below 1300 angstroms. But there should be much to learn from spectra between the first line in the Lyman series — Lyman alpha — and the present ultraviolet limit imposed by our atmosphere at about 3000 angstroms.

It was a revelation to most spectroscopists when recent solar spectrograms that were obtained by the Naval Research Laboratory (Sky and Telescope, July, 1959, page 496) turned out to be full of emission lines in the far-ultraviolet region. After this discovery had been made, it was easy to explain the phenomenon, but to my knowledge no one had clearly predicted a similarity between the solar spectrum from 1500 to 2500 angstroms and the emission-line spectra of such stars as Z Andromedae at the usual photographic wave lengths.



This two-prism spectrograph is used at the Cassegrainian focus of the 82-inch McDonald reflector pictured on the facing page.



Spectral differences between giant and dwarf stars are shown in these spectrograms by W. W. Morgan, Yerkes Observatory. The upper spectrum in each pair is that of the giant. In the two B4 stars, Chi Aurigae has much sharper hydrogen lines than the dwarf Eta Aurigae. There is a similar contrast between the spectra of Alpha Cygni (Deneb) and Lambda Ursae Majoris. Ionized strontium lines are stronger in Zeta Capricorni than in the sun.

Perhaps the most exciting problem in stellar spectroscopy today is the determination of abundances of chemical elements in the atmospheres of stars. We now realize that these abundances are not the same in all stars, and many of the more striking differences have been attributed to evolutionary processes. But I believe that we are beginning to wonder whether chemical abundances are *exactly* the same in *any* two stars.

It is true that the spectrum of the sun can be quite closely matched by spectra of a large number of other stars of similar surface temperature and luminosity. This resemblance is partly due to the multitude of absorption lines produced by atoms of neutral iron, which are prominent features in the spectra of stars like the sun. But on examining the lines of other, less common elements, we almost always find small differences, often too delicate for visual discrimination on the spectrograms, and detectable only by refined spectrophotometric techniques. When photography has been replaced by photoelectric scanning, a large field of study will be opened, and every star, even the brighter ones, will probably reveal remarkable differences in composition.

To illustrate what has already been achieved with spectra of relatively small dispersion, let me describe some work on differences in chemical composition among stars of spectral classes *F* to *K*. This material has been adapted from a chapter by Margherita Hack of Merate Observatory, Italy, in a forthcoming joint textbook on stellar spectroscopy.

In 1950 and 1952, Nancy G. Roman published a study of several hundred stars of spectral classes *F*4 to *K*5. These were divided into two nearly equal groups according to whether the absorption lines of metals appeared to be weak or strong. This was followed by P. Wellmann and Miss Hack's finding that in the stars with weak lines the molecular band due to CH is more intense than in strong-line objects.

The weak-line and strong-line stars show other differences. Because metallic absorption lines tend to be more numerous in the violet region than in the green and yellow, we should expect weak-line stars to show less "blanketing" of the violet continuous spectrum, and they should appear bluer than strong-line stars of similar spectral types. Miss Hack finds that this is apparently the case.

Miss Roman had also noticed that the weak-line stars usually have much larger



Rapidly rotating stars can be recognized by their widened spectral lines, as illustrated by these spectrograms of two double stars. In each pair, the brighter component (labeled A) is rotating faster than the fainter star (B), which consequently has sharper lines. Otherwise, the spectra are similar. In ADS 8257, the primary has an axial rotation of perhaps 100 kilometers per second. Mount Wilson spectrograms by the author.

space motions than those with strong lines. She did not use space motions as a criterion for attributing the stars to one or the other group. Martin and Barbara Schwarzschild, on the other hand, investigated spectra of stars selected by their kinematical properties. Similar studies have been made by J. L. Greenstein and Keenan, and by the latter and G. Keller. All of their results, while differing in details, were quite consistent with the previous work.

Apparently, in this range of spectral types there are three groups of stars. In one, the metal abundances are somewhat larger (about two times) than in the "normal" group; in another they are slightly smaller. In the latter group, absorption bands of CH and perhaps CN are relatively strong. These weak-line stars are often identified with Population II, while "normal" and strong-line stars belong to Population I.

Is there a sharp distinction between these groups? Apparently not. Miss Hack says that although "there is a strong correlation between spectroscopic features and kinematical properties, there are some high-velocity stars which do not show spectroscopic peculiarities, and there are also low-velocity stars that do." She therefore believes that there is a continuous range of chemical compositions from conspicuously low abundances of elements heavier than hydrogen to noticeably high abundances of the same elements. This, in turn, might mean that among the stars of spectral types *F*5 to *K*5 all ages are represented, from the oldest, Population II with extreme weak-line character, to the youngest, Population I with very strong lines.

Such a diversity in the ages of main-sequence stars is not unexpected. The oldest were formed when the interstellar medium contained mostly hydrogen and few metallic atoms. The youngest were created after the interstellar gas had been enriched with heavy elements by the escape of matter from supergiants, novae, and supernovae. However, a marked difference in the ages of giants is not so easily explained. There is every reason to believe that giants in a galactic cluster like the Hyades are younger than those in a globular cluster or a very old galactic cluster, such as M67. But it is not yet certain that we can directly correlate spectroscopic features of giants in the general field of the Milky Way with their ages.

Some of the most recent work on chemical abundances, based upon the best existing spectrograms from Mount Wilson and Palomar Observatories, has been published in the March and May, 1959, issues of the *Astrophysical Journal*. G. Wallerstein and H. L. Helfer find that two *G*-type stars in the Hyades cluster have the same abundances (within 25 per cent) as the sun for sodium and nine heavier elements. Barium may be slightly overabundant, as a result of element building in the stellar interiors. But these results indicate that, in general, between the time that the sun was formed (five billion years ago) and the origin of the Hyades (one billion years ago), there has been little, if any, change in the composition of the interstellar medium.

On the other hand, extremely old Population-II stars show marked metal deficiencies compared to the sun and younger objects. Helfer, Wallerstein, and Greenstein have studied two globular-cluster giants and one high-velocity giant star not belonging to a cluster (HDE 232078), the metal deficiency factor being 20 in two cases and at least 100 in the third (a star in M92). A *K*-type giant in the relatively young galactic cluster M41 (about a hundred million years old) gave roughly the same abundances as the sun. In the old, high-velocity star 85 Pegasi, a visual double, the elements from sodium upward are less abundant than in the sun by factors of three to nine.

These are challenging studies. Persons not acquainted with the many observational and theoretical difficulties involved in these determinations may be surprised that the results are so uncertain. The reason is not imperfect observations and measurements, but the complex process of ascertaining the excitation and ionization of the stellar atmosphere, as well as its turbulence, before the atomic abundances themselves can be computed.



Spectra of four K-type giant stars showing differences in chemical composition. At the top, a star in the galactic cluster M41 has much stronger metallic lines than does the high-velocity star HDE 232078 or two stars in the globular clusters M13 and M92. The hydrogen-gamma line is similar in the four cases, but there is great weakening of the ionized scandium, iron, and ionized titanium lines in the lower spectra. These Palomar spectrograms by H. L. Helfer, G. Wallerstein, and J. L. Greenstein, are courtesy the "Astrophysical Journal."

OBSERVING THE SATELLITES

Touching the Moon

THE first man-made object to reach the surface of the moon was the Soviet space probe that landed on September 13th at 21:02:23 Universal time. The 860-pound spherical instrument package and the 3,331-pound final-stage rocket both struck the target, according to Soviet tracking data.

Despite the incompleteness of information about this historic operation, the great accuracy with which it was performed is evident. For a probe to strike the moon after about 35 hours of flight, the speed of the final stage at burnout must be within 200 feet per second of an intended value of approximately 35,700 feet per second. Furthermore, the trajectory after burnout may not differ by more than half a degree from an angle of 12 degrees with the horizontal. In addition, since the earth is a rotating platform, actual launching has to occur within a fraction of a minute of the predetermined moment, unless course corrections are possible during flight. The success of Lunik II shows that these stringent requirements were met.

Very little has been announced concerning the multistage rocket itself. From the weights of the last stage and instrument package, it is evident that at least half a million pounds of rocket-engine thrust had been employed.

Inside the probe were radio transmitters, operating at three frequencies of 19.993, 39.986, and 183.6 megacycles. When the probe was so close to the moon that it was being speeded up by lunar gravity, observers at the 250-foot Jodrell Bank radio telescope in England noted the expected Doppler change in signal frequency. The signals ceased abruptly at 21:02:24 UT. With an allowance of 1¼ seconds for the time of transmission from the moon, the impact time was only 83 seconds later than the predicted 21:01, announced many hours earlier by Soviet scientists. This close agreement is an indication of the great accuracy of the early tracking from which the forecast was made. American and Japanese radio observers also reported successful tracking operations, but the moon was below the horizon for practically all of the United States at the moment of impact.

Like the earlier Russian lunar probe of January 2, 1959, which missed the moon and entered into orbit around the sun, the September capsule ejected a cloud of luminous sodium vapor, at 18:39 UT on September 12th. Seen from the earth, at a distance of less than 100,000 miles, the short-lived "artificial comet" was located at right ascension $20^h\ 41^m$, declination $-7°.2$. At the time, this part of the sky was unobservable in America, but the luminous cloud was photographed at six or more Soviet observatories.

There appears to be no hope whatever of recognizing the impact crater on the moon caused by the fall of the probe. According to G. P. Kuiper, the craterlet would be only about 100 feet in diameter and with walls 10 feet high, thus lost among the multitudes of tiny uncharted pits that dot the lunar surface. The dust cloud thrown up by the impact might have been detectable. Press accounts of a cloud lasting 58 minutes that was seen by a Hungarian watcher are probably erroneous, as any dust would have fallen back much sooner, lacking the support of an appreciable lunar atmosphere. Details are unavailable concerning the photographs said to have been taken at Lvov Observatory in Russia, showing a cloud immediately following impact that had not been recorded just before it.

According to the Soviet news agency Tass, analysis of the radio tracking data indicated that the point of fall was in the vicinity of the craters Archimedes, Aristillus, and Autolycus, at selenographic latitude +30°, longitude 0°. (The extent of uncertainty was not indicated.) This conclusion was based in part on the readings of a "lunar altimeter," presumably a radar device whose findings were telemetered to earth on the 183.6-megacycle channel.

Some preliminary results have been announced from the data telemetered by the probe. No indication of a lunar magnetic field was found, within the limits of sensitivity of the magnetometer in the instrument capsule. Measurements of cosmic ray flux were made along the flight path, in which nuclei of helium, carbon, nitrogen, oxygen, and heavier elements were counted.



The Vanguard III package being fitted to satellite launching vehicle 7 at the Atlantic Missile Range, Cape Canaveral, Florida, by R. J. Andryshak (left) and D. R. Corbin, both of the National Aeronautics and Space Administration. The satellite was instrumented to measure the earth's magnetic field, solar X-rays, and other conditions in space, such as numbers of meteoritic particles. This was the last ascent in the Vanguard program.

Vanguard III

THE LAST of the Vanguard series of artificial satellites, begun as an American contribution to the International Geophysical Year, was successfully injected into a low-inclination orbit on September 18th at 5:29:45 Universal time, from Cape Canaveral, Florida. This launching was sponsored by the National Aeronautics and Space Administration, which had taken over the Vanguard program from the Office of Naval Research a year ago.

The hour of launching favored optical observations in the evening sky from northern middle latitudes, and many early Baker-Nunn photographs and Moonwatch sightings were made. To most visual observers, satellite $1959\eta_1$ appeared as bright as a 7th-magnitude star.

During its initial revolutions around the earth, the satellite had an anomalistic period (counted from one perigee passage to the next) of 130.19 minutes. It rose to 2,330 miles above the earth's surface at apogee, and descended to 318 miles at perigee, moving in an orbit inclined $33\frac{1}{3}$ degrees to the equator. The orbital characteristics indicate a lifetime of more than 40 years for $1959\eta_1$.

The 100-pound satellite consists of a 20-inch spherical instrument package, bearing a 26-inch-long conical snout, and the unseparated 50-pound final-stage rocket. The fiberglass case of the last of the three stages was intentionally left attached to the instrument compartment, in order to minimize tumbling. (When the cloud-cover satellite Vanguard II was

The components of Vanguard III. At the left are the battery packs and battery compartment lid; next, the lower three-quarters of the satellite's magnesium shell; battery pack tie rods; turn-on plugs controlling transmitter; battery gas relief valve; magnetometer electronics; magnetometer tube and upper quarter of satellite shell; X-ray and environmental electronics and tracking transmitter (sitting atop part of internal structure). In the foreground of the two electronics packages are internal equipment clamps. The pictures are from the National Aeronautics and Space Administration.

sent up on February 17, 1959, separation of the final stage gave the sphere a wobble, thereby making its photoelectric scannings difficult to interpret.) The power supply of 1959γ_1 consists of 62 silver-zinc batteries, weighing 22½ pounds in all, which are expected to function for about 90 days, this depending on temperature conditions.

One major Vanguard III experiment is the measurement of the earth's magnetic field with a new apparatus. The magnetometer is carried at the tip of the 26-inch cone, which is made of fiberglass bonded with phenolic resin, so that the sensor is separated from the metal parts and magnetic fields in the sphere.

In the operation of this device, command from a ground station causes a 6½-ampere magnetizing current to flow for two seconds through a copper coil that surrounds a container of liquid hexane. The strong magnetizing field lines up the spin axes of the protons in the hexane, and this orientation tends to persist for a few seconds after the current is turned off. During the decay of this alignment, terrestrial magnetism imposes a precession upon the protons, thereby inducing a weak current in the copper coil. After amplification, this proton-precession current is telemetered to ground. Eight Minitrack stations have similar magnetometers, for simultaneous observations at the earth's surface. Field strengths as low as 10 gammas (0.0001 gauss) can be measured with this equipment.

Another purpose of Vanguard III is to measure solar X-rays that accompany flares. A pair of ion chambers, for detecting 1-to-10-angstrom radiation, are connected to memory circuits which store the count during a full orbital revolution and then telemeter the data. The count restarts when a photocell observes the emergence of the satellite from eclipse inside the earth's shadow into sunlight. In addition, the X-ray detectors continuously transmit their information to allow detailed study of rapid changes.

Four different methods of detecting meteoritic particles are included in the satellite. The erosion of three chromium strips by bombardment is measured through changes in electrical resistance. The second method employs a cadmium-sulfide photocell behind an opaque screen of mylar and aluminum; when an impact punctures the screen, light passing through the hole is detected by the cell. In addition, there are four microphones listening for impacts. Lastly, two pressurized tanks girdle the sphere, covering about 20 per cent of its area. A differential pressure gauge indicates puncture of one or both tanks by larger interplanetary particles.

The entire array of scientific apparatus in the satellite, exclusive of batteries and supporting structures, weighs only 8½ pounds.

First Two Years of the Space Age

OCTOBER 4th marked the second anniversary of the first successful launching of an earth satellite. The table on the following page lists all known attempts to send scientific payloads into space. This listing is incomplete, for the Soviet Union has not revealed its unsuccessful launching attempts.

In four firings, a total of seven major objects launched by the Russians have entered orbits around the earth, and two long-range probes have been sent aloft. The United States has made 33 attempts, 12 of which have put 15 major objects in orbit, and of five deep-probe launchings,

Satellite and Probe Launchings During the First Two Years of the Space Age

	Launched		Nation	Name and Designation	Status	Notes
1957	Oct.	4	R	Sputnik I — 1957α2	Down about Jan. 4, 1958	Last radio signal Oct. 27, 1957. Rocket, 1957α1, down Dec. 1, 1957.
	Nov.	3	R	Sputnik II — 1957β	Down April 14, 1958	Last radio signal Nov. 10, 1957.
	Dec.	6	A	Vanguard	No orbit	First-stage malfunction.
1958	Feb.	1	A	Explorer I — 1958α	In orbit	Radio silent May 23, 1958.
	Feb.	5	A	Vanguard	No orbit	First-stage control malfunction.
	Mar.	5	A	Explorer II	No orbit	Fourth-stage ignition failed.
	Mar.	17	A	Vanguard I — 1958β2	In orbit	Solar-powered radio still operating. Rocket, 1958β1, in orbit.
	Mar.	26	A	Explorer III — 1958γ	Down late June, 1958	Radio silent June 16, 1958.
	Apr.	28	A	Vanguard	No orbit	Relays for third-stage ignition failed.
	May	15	R	Sputnik III — 1958δ2	In orbit	Solar-powered radio still operating. Rocket, 1958δ1, down Dec. 3, 1958.
	May	27	A	Vanguard	No orbit	Second-stage cutoff at wrong orientation.
	June	26	A	Vanguard	No orbit	Second-stage cutoff premature.
	July	26	A	Explorer IV — 1958ϵ	In orbit	Radios silent Sept. 9 and Oct. 6, 1958.
	Aug.	17	A	Lunar probe	Failed	First-stage malfunction.
	Aug.	24	A	Explorer V	No orbit	Midflight collision of separated stages.
	Sept.	26	A	Vanguard	No stable orbit	Second-stage thrust low. Probably completed one revolution.
	Oct.	11	A	Pioneer I	High trajectory	Reached 70,700 miles from Earth.
	Oct.	23	A	Beacon	No orbit	Payload separated from booster before burnout.
	Nov.	8	A	Pioneer II	Failed	Third stage failed to ignite.
	Dec.	6	A	Pioneer III	High trajectory	Reached 63,580 miles from Earth.
	Dec.	18	A	Atlas-Score — 1958ζ	Down Jan. 21, 1959	Radio silent Jan. 13, 1959.
1959	Jan.	2	R	Mechta — Artificial Planet 1	Solar orbit	443-day period. Radio contact to 373,125 miles. Rocket also orbiting.
	Feb.	17	A	Vanguard II — 1959α1	In orbit	Radios silent Mar. 15, 1959. Rocket, 1959α2, in orbit.
	Feb.	28	A	Discoverer I — 1959β	Down early March, 1959	Radio malfunction.
	Mar.	3	A	Pioneer IV — Artificial Planet 2	Solar orbit	407-day period. Radio contact to 407,000 miles. Rocket also orbiting.
	Apr.	13	A	Discoverer II — 1959γ	Down Apr. 26, 1959	Last radio signal Apr. 21, 1959.
	Apr.	13	A	Vanguard	No orbit	Second stage caused tumbling.
	June	3	A	Discoverer III	Probably no orbit	No telemetry after second-stage firing.
	June	22	A	Vanguard	No orbit	Pressure valve in second stage caused failure.
	June	25	A	Discoverer IV	No orbit	Insufficient second-stage velocity.
	July	16	A	Explorer	No orbit	Power supply for guidance failed. Destroyed by safety officer.
	Aug.	7	A	Explorer VI — 1959δ2	In orbit	Solar-powered radio still operating. Rocket, 1959δ1, in orbit.
	Aug.	13	A	Discoverer V — 1959ϵ	Down Sept. 28, 1959	Radio frequencies undisclosed.
	Aug.	15	A	Beacon	No orbit	Booster fueling failed; orientation wrong.
	Aug.	19	A	Discoverer VI — 1959ζ	In orbit	Radio frequencies undisclosed.
	Sept.	12	R	Lunik II	On moon	Radios ceased operation upon lunar impact Sept. 13, 1959. Final stage also on moon.
	Sept.	17	A	Transit I	No orbit	Third stage failed to fire.
	Sept.	18	A	Vanguard III — 1959η	In orbit	Radios operating.
	Oct.	4	R	Lunik III	In orbit	Radios operating. Rocket also in orbit.

This table is compiled largely from information supplied by the National Aeronautics and Space Administration. Under *Nation*, A designates American launchings; R, Russian. The U. S. S. R. has not released information on unsuccessful attempts. Dates are in Universal time.

three have returned useful information.

Our American space programs have recently been reorganized. The military work, in the hands of the Advanced Research Projects Agency since February 7, 1958, has now been largely assigned to the Air Force, giving it responsibility for developing, producing, and launching military space vehicles. Project Midas, for infrared detection of ballistic missiles, and Project Samos, a satellite reconnaissance system, are also under Air Force control. The Navy will supervise development of an orbiting navigation system under Project Transit, and the Army will handle Project Notus, a satellite communication system.

The NASA has been responsible for the civilian space program since October 1, 1958. Important in this agency's plans is a series of spaceflight vehicles: Delta, Scout, Vega, Centaur, and Nova, progressively scaled to larger and heavier space missions. When Scout is ready, perhaps in mid-1961, three British scientific satellites, now being developed by 10 teams in the United Kingdom, will be launched in a joint operation with the Americans.

The 1960 fiscal budget of the United States' space effort amounts to about 800 million dollars, of which more than 500 million is allocated to NASA.

Translunar Orbit

EARLY on October 4th, exactly two years after the first Sputnik launching, Soviet scientists sent a 614-pound satellite into an immensely elongated trajectory past the moon, in company with the 3,424-pound final stage of the carrier rocket. While few of the details had become available at the time of writing, it was claimed by a Russian announcement that the satellite is moving around the earth in an orbit for which the apogee distance is 292,000 miles and the perigee distance about 25,000 miles. The corresponding period of revolution is about 14 days.

Presumably the final-stage rocket, which also carries a payload of 345 pounds of instruments and batteries, is traveling in a similar orbit. An earlier Soviet report gave the closest approach to the moon as about 4,400 miles, at 14:16 Universal time on October 6th.

As seen from the moon, the satellite's path was a hyperbola, as the velocity was too great to allow capture by the moon. If, as seems probable, the probe crossed the lunar orbit before the moon arrived at the intersection point, then the probe's motion would be decelerated, shrinking its orbit and shortening the period. (The reverse effects would have occurred if the moon arrived first.)

If the orbit is actually an ellipse of this character, it will be subject to major changes, as the perturbations by the lunar and solar attractions can be very large. While predictions for such a probe offer many complications, its orbital motion will be of considerable interest to astronomers in the field of celestial mechanics. Because the probe carries solar batteries as well as chemical ones, its radio transmissions, at 39.986 and 183.6 megacycles, should permit tracking for a long time.

MARSHALL MELIN
Research Station for Satellite Observation
P. O. Box 4, Cambridge 38, Mass.

The Great Aurora of Early September

Auroral rays diverging from the observer's zenith form a corona during the display of September 3rd, photographed at Sandwich, Illinois, by Nicholas Liepins at 10:16 p.m. Central standard time. The exposure was one minute, at f/2 on Tri-X film. At bottom center the stars of Delphinus are bright, while Epsilon Cygni is in the upper right.

AMATEUR ASTRONOMERS from coast to coast witnessed a spectacular auroral storm on the night of September 3-4, 1959. Auroral activity on a much smaller scale was observed on the following nights, but these weaker displays ended by midnight.

Solar observers noted increased activity on the sun three days prior to the northern lights. Sunday, August 30th, Howard C. Colton, of Fairport, New York, obtained the accompanying photograph, which shows a large number of sunspots and faculae.



Pronounced activity on the sun, indicated by the number of sunspots and faculae, preceded the widespread auroral display. This picture was taken on August 30th by Howard C. Colton, using Kodak Autopositive film, at the prime focus of his 6-inch f/15 telescope.

The duration and intensity of the September aurora varied with the locality, the shortest displays being reported from central and western sections of the country. The most distant observer was Alan McClure, who had set up cameras on Mt. Pinos, about 75 miles northwest of Los Angeles, California, to photograph Comet Alcock 1959f. He saw auroral activity at 7:40 p.m. and 12:35 a.m. Pacific standard time.

A long and detailed log of the event, as it appeared from Eightyfour, Pennsylvania, was kept by 15-year-old David R. Kaiser, from 10:30 p.m. to 4:00 a.m. Eastern standard time. He first noted a large rayed arc, blue-green in color, stretching from the western to the eastern horizon. In shape it resembled a huge rainbow.

Changes came rapidly, according to Mr. Kaiser. At 10:40, a beautiful green

Compare this picture of the auroral corona with the one at the top of the page, taken 14 minutes earlier, both in Illinois. Elmhurst amateur James M. Hyer used an exposure of 25 seconds, at f/2.8 on Plus-X film. The stars of Delphinus can be identified near the bottom in this picture, too.

coronal aurora formed near the zenith, where pulsating rays from all four quadrants converged. Five minutes later, the corona dimmed and became whitish. In the west he saw a low glow, with a red ray rising from it 70 degrees up the sky. Northward, a rayed arc extended to the zenith, while in the south and west a short-lived ray bundle reached to 10 degrees of overhead.

The corona had vanished completely by 10:53, and all that remained was a rayed arc, covering the west, north, and east. This showed waves of light, traveling upward and from east to west. At exactly 11 o'clock this ceased, and a new rayed arc formed in the north, its western end doubled. Flaming aurora appeared in the east, varying from red to green, and pulsating in brightness.

As the Pennsylvania amateur continued his watch, he observed a sudden increase in auroral activity 12 minutes later, the rayed arc rising to an altitude of 75 degrees and changing to a brilliant red. At 11:20 a corona appeared nearly overhead, brightest on the north and with a gap on its southern side. This green corona disappeared five minutes later, leaving an arc in the north and west, from which rose a pulsating sheaf of rays. Still another brilliant green corona formed at 11:30 and grew larger. From it diverged red rays toward the four points of the compass. Ten minutes later this spectacular complex was nearly gone, the most conspicuous feature being a rayed arc in the northern sky.

Thereafter auroral activity lessened. From 2 to 3:45 a.m., Mr. Kaiser saw only a glow in the north that shrank progressively until by 4 o'clock it was completely gone. But an hour earlier, in Pittsburgh, Pennsylvania, W. A. Feibelman had resumed observing the aurora after a rest of three hours, to find the whole northern sky lit up with waves and rays of intense green color. He finally lost the declining aurora in the dawn light.

The following amateurs also described the September display, including the distinctive coronal activity: L. E. Reinagel, Kenmore, N. Y.; D. Czarick, Pottstown, Pa.; R. R. Zappala, Cleveland, Ohio; J. Stamm, Jr., Akron, Ohio; J. M. Hyer, A. Stewart, F. Mraz, and J. A. Marshall, Elmhurst, Ill.; N. Liepins, Sandwich, Ill.; R. Spaulding, Royal Oak, Mich.; G. R. Peters, Detroit, Mich.; D. G. and R. J. Airhart, Toronto, Canada.

Other reports were submitted by A. Moehrke, Poughkeepsie, N. Y.; D. Della Pietra, E. Rochester, N. Y.; L. Rick, Lorain, Ohio; G. Hunter, Ann Arbor, Mich.; J. P. Hyde, Elkader, Iowa; P. Connor, Fremont, Nebr.; J. R. Otoupalik, Greeley, Colo.; H. W. Lang, St. Louis, Mo.; and P. Nash, Townsend, Mont.

On September 5th an aurora was seen by L. Mattersdorf at Chesterfield, N. H.; H. A. Luft, Oakland Gardens, N. Y.; and E. D. Alpert, New Bedford, Mass.

The display was noted for the variety of its forms. At Pittsburgh, W. A. Feibelman took these pictures that illustrate some main auroral types. The first, taken at 9:50 p.m. EST, shows a broad, homogeneous arc in the north. By 10:42, the arc had formed a rayed structure, as shown in the second picture. At 3:40 a.m., when the third photograph was made, the entire northern sky was bright with intensely green draperies. This was by far the brightest aurora of the year, rivaling the great display of February 10-11, 1958, according to Mr. Feibelman's report.

AMERICAN ASTRONOMERS REPORT

Here are highlights of some papers presented at the 103rd meeting of the American Astronomical Society at Toronto, Canada, Aug. 30-Sept. 2, 1959. Complete abstracts will appear in the Astronomical Journal.

Distance of the Sun

An extremely precise radio method of finding the sun's distance has been proposed by A. E. Lilley, Harvard Observatory, and Dirk Brouwer, Yale Observatory. It uses the same principle — Doppler shifts caused by the earth's motion around the sun — employed by an older optical technique involving observations of stellar spectra. When the earth moves toward a star, there is a shift of stellar spectrum lines toward shorter wave lengths, while for a receding earth the displacement is to longer wave lengths. In practice, the optical procedure did not give the accuracy that its radio counterpart now offers.

If a radio source is located behind a cloud of interstellar hydrogen, the 21-centimeter line of neutral hydrogen can be observed as a very sharp absorption feature. It is planned to measure the frequency of such an absorption line (for a source near the ecliptic) at different times of the year for several years. Because of the earth's revolution, the observed frequency of the absorption line will vary back and forth during the year. Thus it will be possible to ascertain the orbital velocity of the earth, and hence the precise size of its orbit and the distance to the sun.

Dr. Brouwer's computations show that the method should be capable of fixing the sun's distance to one part in 300,000 — a precision hitherto promised only by radar observations of Venus. The main limitation to the accuracy attainable is the rotation of the earth, which may change the observed frequency of the 21-cm. line by one cycle in the course of 20 seconds. This thus limits the time that can be profitably spent on any single measurement.

Actual observations will start early in 1960, at the Naval Research Laboratory and at the Agassiz station of Harvard Observatory. For recording the hydrogen line, a new radiometer has been developed, with financial support from the National Aeronautics and Space Administration.



The University of Toronto, where the American Astronomical Society met, operates this 74-inch reflector, which for 25 years has been Canada's largest telescope. It is housed in a dome 61 feet in diameter.

Moon's Distance by Radar

Hitherto, astronomers have relied upon triangulation for finding the distance from the earth to the moon, either by direct measurements of the moon's position in the sky as seen from widely separated observatories, or from times of occultations of stars. With either method, the moon's distance can be derived in terms of the earth's size, which is known rather precisely.

Preliminary results from a radically different approach — radar ranging — were reported by R. H. Bruton, K. J. Craig, and B. S. Yaplee, of the Naval Research Laboratory. During October and November, 1957, they measured the two-way travel times of 60,000 pulses of 10-centimeter radio waves reflected from the moon. Although the beam-width was 30 minutes of arc, at this wave length the moon reflects like a polished sphere, all of the echo coming from a very small area in the middle of its disk.

The NRL scientists found in this way that the average center-to-center distance to the moon is 238,855.95 miles, with an uncertainty of ±0.75 mile. This is in close agreement with the results previously found by triangulation.

One unexpected difficulty is that the daily measurements showed a monthly variation of a large fraction of a mile, after all known effects had been allowed for. To investigate this phenomenon, a new seven-month series of observations is under way.

Pressure-Induced Nuclear Reactions in Stars

Astrophysicists have long been familiar with thermonuclear reactions in stellar interiors, processes whose rates depend mainly on high temperature rather than density. A. G. W. Cameron, of Atomic Energy of Canada, Ltd., now calls attention to a different behavior of nuclear reactions occurring in extremely condensed matter. Such processes are little affected by temperature, but their rates are very sensitive to density.

For them, Dr. Cameron has coined the term *pycnonuclear reactions*, from the

Greek *pyknos* (compact, dense). He has computed the rates for three important processes of this kind: the conversion of C^{12} to O^{16}, N^{14} to O^{18}, and O^{16} to Ne^{20}, a helium nucleus being added in each case.

One instance where processes of this sort may take place was described by Dr. Cameron. In a Population-II star of about the sun's mass, helium reactions can build up elements as massive as neon, but the latter needs a temperature of 1.2 billion degrees to enter into further thermonuclear phenomena. However, in such a star this temperature might not be reached, and after the helium burning stops the star's core would contract to extreme density, becoming degenerate.

At this stage pycnonuclear reactions would set in, the star expanding (possibly explosively as in a nova) because of the heat generated. The C^{12}-O^{16} reaction may be the principal ignition process initiating nova explosions. For stars at the tip of the red giant sequence in Population I, the N^{14}-O^{18} reaction is probably the cause of core expansion.

Color-Magnitude Diagram for Visual Binaries

In recent years, color-magnitude diagrams have been derived for numbers of galactic and globular star clusters. Two important uses of such diagrams are to indicate observationally the evolutionary development of stars, and to determine accurate absolute magnitudes for stars of high intrinsic luminosity. Because so few star clusters are suitable for this kind of study, Gustav A. Bakos, now of the Smithsonian Astrophysical Observatory, points out that visual binary stars can add valuable information.

According to current views of stellar evolution, if the two components of a binary star system differ in brightness by several magnitudes, in the great majority of cases the fainter component will be on the main sequence. Accordingly, the color of the fainter star tells its absolute magnitude. Then, if the magnitude difference between the components is measured, along with the color of the brighter star, its location in the color-magnitude diagram is also established.

Dr. Bakos proceeded to measure photoelectrically the apparent magnitudes and colors of both components of 75 binary systems, using for this purpose the 19-inch reflector of David Dunlap Observatory. He took spectra with the David Dunlap 74-inch reflector at a dispersion of 33 angstroms per millimeter, in order to determine the absolute magnitudes of the primary components. In this way he compiled the color-magnitude array shown here. The primary components, plotted as open circles, are either giants or main-sequence stars. With few exceptions, the fainter members are on the main sequence, as expected.

In this diagram, curves have been added to show the evolutionary tracks of stars in a number of galactic star clusters. The greater the age of a cluster, the lower is the branching-off point of its track upward from the main sequence. Also, the more massive a star, the faster it will evolve, and the farther it will have moved in the diagram from the zero-age main sequence.

By matching the positions in the color-magnitude array of the components of a binary system with the track of a cluster, the approximate age of the binary can be found. Thus, a binary for which the plotted points agree with the pattern for NGC 752 should have about the same age, on the order of one billion years, as that cluster.

Of special interest are the double stars that lie in the same region of the diagram as the very old cluster M67, whose age is generally considered to be about five billion years. Those visual binaries whose brighter components lie to the right of the cluster track must be even older, about 10 billion years, according to Dr. Bakos' estimate. He points out that some recently observed galactic clusters have color-magnitude arrays indicating they, too, are older than M67.

Polarization of Light of Moon and Planets

A difficult but important observational problem is the comparison of the properties of interplanetary particles with those in interstellar space. But the moon, Mars, and Venus are continually collecting interplanetary particles, which may be expected to produce polarization effects in reflected sunlight. At McDonald Observatory in Texas, the Indiana University astronomer Thomas Gehrels has begun a study of these solar system bodies, as well as of the nebula NGC 7023, which shines by starlight scattered by interstellar dust.

Dr. Gehrels' observations are made with a Wollaston photometer attached to the 82-inch reflector. In this device, light is separated by a Wollaston prism into two beams, plane polarized at right



Left: Binaries with color-magnitude properties like those of stars in the cluster M67, indicating that they have similar evolutionary ages. Most of the primary stars lie on the giant branch, the secondaries on the main sequence of the cluster.



Right: The colors and absolute visual magnitudes of visual binaries, determined by Gustav A. Bakos with the 19-inch and 74-inch telescopes of the David Dunlap Observatory. The thin lines connect primary and secondary stars in a few special cases. Superimposed on the chart are the observed sequences of important galactic clusters, their ages ranging from a few million years, for the Double Cluster in Perseus, to five billion years for M67 in Cancer.

angles to each other, whose intensities are measured with photomultiplier tubes. A complete observation, requiring 30 minutes, consists of such measurements for six orientations of the prism, in ultraviolet, green, and infrared light. Because the two beams are recorded simultaneously, even during poor seeing or variable sky transparency the amount of polarization can be determined with an accuracy of about 0.1 per cent.

Seven small lunar regions, each about four miles in diameter, have been observed in detail to determine how the polarization and brightness change with the moon's phase. The polarization is least in the infrared (10,330 angstroms) and greater in the green. In the ultraviolet, at 3250 angstroms, the polarization is strongest, and for lunar maria at quarter phase can amount to as much as 23 per cent. In this respect, the floor of the crater Plato behaves as the maria do. Dr. Gehrels' measurements in green light agree well with the earlier visual polarimetric observations by B. Lyot in France.

Mars and Venus show the same wavelength dependence as the moon, strongest polarization occurring in the ultraviolet. A peculiar result is that for Venus the ultraviolet plane of polarization differs by 90 degrees from that at longer wave lengths.

The marked ultraviolet effect indicates that light is being scattered by particles smaller than about 0.3 micron (1.2×10^{-5} inch) in diameter. In the case of Venus, two types of particles may be involved. Dr. Gehrels comments: "On Mars, and even on Venus, we may be observing accreted interplanetary particles suspended in the atmosphere." He suggests that they are of the same nature as micrometeorites in the terrestrial atmosphere.



In the Meinel ratio spectrometer, the narrow slit allows only a 50-angstrom band-width to pass to its photocell, but the slit jaws are polished to reflect the remaining spectrum to the second slit, which is wide enough for 250 angstroms to pass.



Ordinary spectrographic intensity tracings are dominated by the basic energy curves of the stars, as shown by the dotted lines in these charts of Altair, an A7-type star, and Arcturus, of type K2. The Meinel spectrometer compensates for the energy curve, however, giving the normalized solid-line tracings. Kitt Peak National Observatory charts.

As for the reflection nebula NGC 7023, parts of it were observed at wave lengths of 3700, 5700, and 8200 angstroms, where 13, 19, and 22 per cent polarization are typical values. These indicate interstellar particle sizes of a few microns. But in the case of eight individual stars, the infrared interstellar polarization was found to be appreciably less than at 6500 angstroms.

Ratio Spectrometer

A. B. Meinel, of Kitt Peak National Observatory, has invented a simple and effective device for charting automatically the light intensities of features in a stellar spectrum. The starlight collected by a telescope falls on a grating that forms a spectrum which is scanned simultaneously by two slits. One slit views a 50-angstrom width of the spectrum, the other a 250-angstrom band centered upon the first.

During scanning, the two slits remain fixed while light reaches them from a turning mirror. Behind each slit is a 1P21 photocell, and the *ratio* of the two photocurrents is traced by a pen recorder. Localized absorptions and emissions in the spectrum appear as dips and peaks, respectively, on the tracing.

Besides its speed and convenience, the ratio spectrometer has other important advantages. Because the spectrum is in effect being compared with itself, the intensity ratio is unaffected by fluctuations in sky transparency or seeing, or by guiding errors. Also, the general course of the tracing is more-or-less horizontal, making small features easier to recognize.

Dr. Meinel has built an experimental model of the device, using it with a 16-inch Cassegrainian reflector at Kitt Peak to observe 200 stars of spectral type G. It is possible to record stars as faint as the 5th magnitude, 10 minutes being required for a scan from about 3500 to 5500 angstroms.

The ratio spectrometer can be used for accurate spectral classification and for the determination of absolute magnitudes of stars. Dr. Meinel suggests that the device, when used with a very large telescope, may serve for measuring the red shifts of remote galaxies, by noting the wave-length change of the entire spectrum pattern with respect to the spectrometer's calibration scale.

DAYTIME AURORAS

For many years there has been controversy concerning the possibility that intense auroral displays may be visible in full daylight. Until recently, the only report that had gained general acceptance was an observation in New Zealand, in May, 1921. Cicely M. Botley has now collected 15 probable cases of daylight auroras between the years 1744 and 1958.

From her list, published in the August *Journal* of the British Astronomical Association, she concludes that the phenomenon is by no means as rare as has been thought. "It therefore seems that aurora, at any rate in the great class, is not confined to the night hemisphere but is global in the most literal sense, a conclusion also supported by radar."

NEWS NOTES

UNIVERSITY GIVES OBSERVATORY TO AMATEUR ASTRONOMER

When Miami University, at Oxford, Ohio, recently decided to dismantle its observatory, the 12-inch Clark refracting telescope was presented to Leslie C. Peltier, of Delphos, Ohio. One of the nation's most famous amateurs, Mr. Peltier has discovered a dozen comets since 1925, and has been a very active variable star observer. Most of his work has been done with a 6-inch refractor lent to him by Princeton Observatory.

The 12-inch f/15.7 objective was made by Alvan Clark and Sons in 1868. For many years the telescope was used at the observatory of Wesleyan University, Middletown, Connecticut, and was transferred to Miami in 1925. The gift to Mr. Peltier also included the observatory building with its 22-foot dome, a 3-inch Gaertner transit instrument, and a sidereal clock.

Already moved by truck to Delphos, the building and telescope are being re-erected exactly as they were at Miami University, on Mr. Peltier's property about 100 yards north of his residence. The location is at the west end of the town, with no detrimental lights or smoke. The 12-inch is expected to be in active use by the end of the year, for observations of faint variable stars.

VELOCITY OF LIGHT

There is a curious discordance among modern experimental determinations of the velocity of light, for measurements made with visible light give slightly smaller speeds than do those with radio microwaves. R. A. Miller and A. Lopez, Manila Observatory, suggest that this difference arises from the time taken by reflection from mirror surfaces used in the optical methods. The work done with standing microwaves in resonant cavities should be essentially free of such an effect.

In support of their conjecture, the Philippine scientists call attention to the optical determination by L. E. Bergstrand a decade ago. This used only a single mirror reflection and gave a velocity of 299,792 kilometers per second — very close to the results of microwave experiments.

On the other hand, in the optical determinations made by A. A. Michelson and others up to 1935, light beams were sent back and forth over long paths by means of multiple reflections from rapidly rotating mirrors. Thus F. G. Pease and F. Pearson used 13 reflections between points more than 10 miles apart, finding a velocity 18 kilometers per second less than Bergstrand's.

Drs. Miller and Lopez have made a first attempt to calculate the delay suffered by visible light during reflection from silver, finding roughly 1.7×10^{-10} second. This lag is of about the right size to account for the differences between the early optical and the recent microwave work, they report in the September *Journal* of the Optical Society of America.

CATALOGUE OF DWARF GALAXIES

Sidney van den Bergh, of David Dunlap Observatory, has used National Geographic-Palomar Sky Survey prints to compile a catalogue of 222 dwarf galaxies, characterized by very low intrinsic luminosities. Faint objects were included in his list if they met two criteria: low surface brightness, and little or no central condensation on the photographs taken in red light.

Four different types of such galaxies are distinguished by Dr. van den Bergh. *Dwarf irregulars* resemble NGC 6822 in the local group of galaxies. *Dwarf spirals* exist in two varieties, either as a short bright bar on a fainter background, or with resolved stars and nebulosity in elongated patches resembling segments of a spiral arm. *Dwarf spheroidal* systems, of very low surface brightness, are rather easy to identify at large distances, but more difficult to recognize if they are near enough to be completely resolved into stars. IC 3475 in the Virgo cluster is the brightest known representative. Lastly, there are *dwarf ellipticals*, but they are difficult to distinguish from their giant counterparts, and Dr. van den Bergh's list contains very few.

Of the 22 known probable members of the local group of galaxies, no fewer than 17 are dwarf systems. A dozen were included in Dr. van den Bergh's survey, which is published as Vol. 2, No. 5, of the *Publications* of David Dunlap Observatory.

GUSTAV LAND DIES

American astronomy lost a major link with the German tradition of celestial mechanics and refined positional measurements when Gustav Land of Yale Observatory died on September 26th.

Born at Danzig on April 15, 1880, as Gustav Deutschland, he studied at Berlin University, where his thesis was an investigation of the motion of a comet during a very close approach to Jupiter. Afterward he was at Leipzig Observatory, and became well known as an expert in stellar statistics.

In 1939, Dr. Land left his native country, and, after working for a short time with the British Nautical Almanac Office, came to the United States as a staff member of Sproul Observatory. He was appointed research assistant at Yale in 1941, a position he held until his death.

While in America, his chief study concerned the measurement of star positions on photographic plates, and methods for pushing the accuracy of this work as far as possible. Particularly noteworthy were his studies of errors caused by emulsion shifts, and of practical ways of overcoming them. Dr. Land published in the Yale Observatory *Transactions* a searching analysis of the accuracy of the Yale measurements of trigonometric parallaxes of stars. The conclusions he reached in this memoir are of basic importance to improving our knowledge of stellar distances.

IN THE CURRENT JOURNALS

OCCURRENCE OF LIFE IN THE UNIVERSE, by Su-Shu Huang, *American Scientist*, September, 1959. "Granted that all kinds of stars have an equal chance of possessing planets, we ask: Is there any way of knowing which kinds of stars favor the existence of life on their planets? This question can be reasonably answered, we find, with our present knowledge."

METEOR TRAILS USED IN RADIO COMMUNICATION

In recent years, the overcrowding of the high-frequency bands normally employed for long-range radio communications has stimulated varied attempts to utilize wave lengths so short that ordinarily they would serve only over a line-of-sight path. The National Bureau of Standards now reports successful development of a two-way message transmission system, in which 49-megacycle signals are reflected from the ionized trails left by meteors.

The message to be sent is first recorded on magnetic tape. With both transmitters on the air, the presence of a suitably located meteor trail is detected within a few thousandths of a second. Then the message is sent, most satisfactorily at 2,400 words per minute — 40 times the speed of present teletype transmission. When the signal strength falls too low, as the meteor trail dissipates, the transmission is halted temporarily.

Extensive tests made over an 800-mile path shows the method can compete effectively with other long-range systems, and is relatively free from ionospheric disturbances. However, the simultaneous occurrence of two meteors can cause garbled signals.

In a project known as Janet, Canadian scientists had previously used meteor ionization to aid short-wave radio communications (SKY AND TELESCOPE, October, 1956, page 541).

INTERNATIONAL ASTRONOMICAL UNION 1961 MEETING

The 11th general assembly of the International Astronomical Union will be at the University of California in Berkeley during August, 1961. More than 1,000 astronomers from all parts of the world are expected to attend. IAU meetings are held every three years, the latest having been at Moscow in 1958.





The southern part of the moon, as represented on one sheet of Philipp Fauth's great lunar map, left incomplete when he died in 1941 and now being prepared for publication by his son, Hermann Fauth. The original is on a scale of one to a million, corresponding to 11½ feet for the moon's diameter. This reproduction is on a scale only 29 per cent as great. The small key chart, left, identifies several of the more prominent craters, including Clavius, 144 miles in diameter; Tycho, 54; and Moretus, with its massive central peak, 73 miles. Fauth uses contour lines to indicate vertical relief, a change from the hachures in his earliest charts. These contours are careful estimates rather than the result of detailed measurements. Possibly never again will a lone selenographer prepare so elaborate a map of the whole moon entirely from his own visual observations. The multitude of smaller formations visible in very large telescopes is so great that future visual charting will probably be co-operative, insofar as it is not superseded by photography or actual exploration. To appreciate the amount of detail in Fauth's map, and to judge its accuracy, the reader should compare the view in his own telescope with the depiction of, say, the crater Clavius.

All illustrations with this article are courtesy Hermann Fauth.

PHILIPP FAUTH and the Moon

HERMANN FAUTH

Philipp Fauth was 63 when this picture was taken in the summer of 1930. At that time, the German amateur was resuming lunar observations with a 15½-inch refractor at his fourth observatory (pictured on page 23), at Grünwald, in Bavaria.

ANYONE who writes about the history of selenography should no more omit the name of Fauth than those of Mädler and Schmidt. Fauth was a leader in lunar studies and the author of maps and publications showing his unprecedented knowledge of the moon's surface. Although this amateur was world-famed among astronomers during his lifetime, recent books either fail to mention him or give him only a few inadequate words. In America he has remained almost unknown.

Philipp Johann Heinrich Fauth was born on March 19, 1867, at Bad Dürkheim in the German Rhineland, the oldest of three children in a long-established family of pottery makers. From his father he acquired outstanding artistic talents, that later showed themselves in unsurpassed lunar maps, and a lifelong love for music. Tending the pottery kiln at night, the father would fetch his youngster out of bed and carry him outdoors, wrapped in a blanket, to show him the beauty of the stars and the sun rising over the vineyards of the Rhine Valley. Coggia's comet in April, 1874, made a deep impression on the boy.

At secondary school in Kaiserslautern, Fauth began an enthusiastic study of astronomy. In 1885 he started observing the sun and moon with a small 18-power telescope, and two years later acquired his first really serviceable instrument, a 3-inch refractor. His two aims were to become acquainted with everything to be seen in the sky, and to draw what he saw better than the usual pictures in books. (At that time astronomical photography was a little-known novelty.) The boy's inspiration came from books, for he had not yet visited an observatory or met an astronomer.

In 1890, when Fauth became a school teacher, he built his first observatory in the southern part of Kaiserslautern, equipping it with a 6-inch Pauly refractor. At that time the west German observatories of Karlsruhe, Heidelberg, and Cologne had nothing larger. He devoted himself at once to the moon, Jupiter, and the nebulae. But two years later, the school board transferred him to the remote town of Oberarnbach in the western Rhineland, for it was taken amiss that a young teacher should have such a good city post.

Now his observing had to be done long-distance, each working night requiring a three-mile walk over a hilly road, then a 10-mile train trip, and another half mile on foot to the observatory — and back again in the morning. He kept this up for four years, in addition to his regular school duties. Fauth recollected: "During those years I won my spurs as an astronomer. In the summers I more than once saw the sun set as I began to observe, and rise as I closed up the observatory. In winter I needed all my enthusiasm. Deep snow and bitter cold were obstacles, but steady seeing and a transparent sky encouraged me to persist; the unforgettable impressions even during winter nights are among my fondest observing memories."



Fauth's first observatory, in use from 1889 to 1895, was located on the Lämmchesberg, near Kaiserslautern. Here, with a 6-inch refractor, he made the observations on which was based his atlas of 25 lunar regions.

The fruits of these strenuous years





The third observatory that Fauth built, in use from 1911 to 1923, was on the Kirchberg near Landstuhl, in western Germany. At the left, Professor Fauth stands before the building. The interior view shows the 15½-inch medial refractor, an instrument of exceptional optical quality, which gave practically complete color correction. With this telescope the German amateur did much of his finest work.

were Fauth's first two memoirs, published in 1893 and 1895: *Astronomical Observations and Results in the Years 1890 and 1891,* and a similar report for 1893 and 1894. The latter included a topographic atlas of 25 lunar regions. Articles by him appeared in journals such as the *Astronomische Nachrichten* and *Sirius,* and he was already corresponding actively with fellow observers, especially Hermann Klein in Cologne, J. N. Krieger in Gern, Viktor Nielsen in Copenhagen, and Max Wolf in Heidelberg.

Fauth's success gained him the support of the Prussian Academy of Sciences for the construction of a more serviceable observatory in a better location. In 1895 he obtained a transfer to the school at neighboring Landstuhl, and near there on the Kirchberg he erected his second observatory, a dome-topped stone tower 26 feet high. It looked out over the treetops 450 feet above the town and 1,300 feet above sea level. Here Fauth obtained a wealth of observational results, and by 1902 had discovered some 5,600 new lunar craterlets and clefts. He was now established as a lunar specialist. In 1906, Wilhelm Foerster of Berlin encouraged him to write his first book, which also appeared in English translation as *The Moon in Modern Astronomy,* London, 1907. Professor Foerster furthermore recommended him for the post of assistant at a new observatory in Mexico, which, however, he decided not to accept.

In 1911 Fauth acquired a 15½-inch f/10 refractor of the medial type. This design, invented by Ludwig Schupmann of Aachen, provided images unusually free of chromatic aberration, and gave superb performance. The third observatory, built for this telescope, stood 600 feet south of the stone tower. To the low sheet-zinc dome in a grove of young pine trees, Fauth welcomed thousands of visitors in the following years.

This photograph, taken in 1890, shows the young schoolteacher at the outset of his career as a lunar specialist.

His 28 years of activity at Landstuhl furnished the basis for many memoirs and books: in 1898, *Observations of the Planets Jupiter and Mars at Their Oppositions of 1896-97,* with 147 drawings and five maps; in 1912, a 790-page work, *Hörbiger's Glacial Cosmogony;* and in 1916, his *25 Years of Planetary Investigation,* with 245 illustrations. There were also innumerable lunar drawings, thousands of sunspot sketches, and an analysis of almost 3,000 fine drawings of Jupiter. Moreover, he wrote numerous astronomical articles and kept up an extensive scientific correspondence.

This intense activity was interrupted in 1923, when Fauth left the Rhineland because of the French occupation, and he became a teacher in Munich. He could not bring his medial to Bavaria until 1930, when he set it up near the town of Grünwald, nine miles south of Munich. There he continued his work, publishing at the age of 70 a large collection of drawings of formations very near the edge of the moon, observed at times of especially favorable libration. Three years later he was planning to move his observatory



One of the 16 large-scale charts of Fauth's 1932 regional atlas shows the Hell Plain, in the south-central portion of the moon. The ruined and incomplete remains of an enormous crater, this plain has never been officially named, but Fauth designated it Hörbiger. About two inches left of the crater Hell on this chart is Cassini's bright spot, one of the most brilliant lunar areas at full-moon phase. The low-walled crater labeled Schupmann is generally known as Hell B.

to Rauhe Alb in Swabia, when he died on January 4, 1941. Philipp Fauth's last contribution was a yet-unpublished work of advice and suggestions for future lunar observers.

Today the sites of Fauth's four observatories have been absorbed by the growth of cities, and his famous medial telescope disappeared at the end of World War II. There remain only his dwelling and his grave in Landstuhl, and streets named for him in Bad Dürkheim, Landstuhl, and Grünwald. His main legacies are his great selenographical works: the comprehensive 600-page treatise of 1936, *Our Moon;* his lunar atlases of 1895 and 1932; and his great lunar map, 11½ feet in diameter and on a scale of 1:1,000,000.

The 22 sheets of this map survived World War II, but Fauth had lived only long enough to make the finished drawings for five of them. As his son, I have undertaken to complete the careful pencil drafts of the remaining 17 sections. The long-awaited publication of this map will probably be within a year. From the sample section reproduced on page 20, the reader may gather some impression of the remarkable detail and realism of Fauth's representation of the moon.

This map has had a long history. Over the years Fauth spent thousands of hours at the telescope, making the drawings that in 1936 he began to combine into a general map. Already in 1932 he had prepared a lunar atlas of 16 regional charts, the labor of a period when convalescence from a severe illness interrupted his observing. These 16 charts were beautiful depictions on a large scale of areas such as Eratosthenes, Taruntius, Plinius, Ptole-

Fauth's fourth observatory, at Grünwald, which he used from 1930 until his death in 1941. He is here sitting on the edge of the dome slit, and the medial can be seen inside. The hut at the left contains a small telescope for sunspot counts.

This map of the large crater Copernicus was prepared by Fauth in July, 1932, from observations with his 15½-inch refractor. The original scale is 1:200,000, the contour lines indicating 200-meter intervals in elevation. At top center is a conspicuous double crater, the larger being Fauth, the smaller Fauth A. Nearest the bottom edge is the crater Gay Lussac. Copernicus itself is remarkable for its intricate interior terraces, the inner wall descending in a series of irregular steps.

maeus, and Posidonius, as well as the Hell Plain, and the double crater Fauth (officially named after him). The great formation Copernicus is shown as it would be seen from directly above, without foreshortening, on a scale of 1:200,000.

The work on the 11½-foot map was delayed by Fauth's continual desire to add new details from additional observations. In 1939 he wrote, "I am constantly finding peculiarities that lend fresh interest to the work. In a way, all lunar maps are premature, for the detail is truly inexhaustible."

This cautious and responsible attitude helps explain Fauth's deep skepticism of reports of lunar change. In an important article (*Astronomische Rundschau, 3,* 172-176, 1901), he marshaled strong arguments that no alteration had actually occurred in the famous case of the object Linné, in Mare Serenitatis, and he again presented his proofs in his 1936 book. Yet others ignored his demonstration that Schmidt was mistaken in believing that Linné had altered from a crater to a bright patch.

"Without this error," he wrote, "lunar literature would be free from many of its fantasies; as it is, proofs of 'changes' on the moon spring up like weeds. We witness over and over again how 'scientific' methods and time and effort are squandered on unprofitable problems." Although Fauth was widely recognized as the leading lunar expert of his day, his warning went unheeded.

The German selenographer was deeply disappointed that for decades he had been publishing fine drawings of the moon without finding fellow workers whom he regarded as of his own caliber. "Either they were content with the success of lunar photography, or they shied away from creative work at the eyepiece, which demands patience and skill. Instead, the enthusiasm of students of the moon has been for selenological problems, and they have advanced wild theories without coming to grips with hard facts. Lunar work has not achieved the high level that should have been stimulated by the publication of Schmidt's lunar atlas of 1878."

Philipp Fauth's extensive contributions to lunar studies were nevertheless only one aspect of a versatile career. At the same time, he was a widely known educator and a highly accomplished musician. The biography that I am now writing should furnish a clear and faithful image of him.

QUESTIONS... FROM THE S+T MAILBAG

Q. How many natural satellites in the solar system can be seen with a 3-inch telescope?

A. About eight. These are the earth's moon, the four Galilean satellites of Jupiter, and three of Saturn's moons — Rhea, Titan, and perhaps Iapetus.

Q. Have any features on the invisible side of the moon been tentatively mapped?

A. Yes. Certain bright rays reach around the moon's edge into the visible hemisphere. On the assumption that these rays, like those of Tycho and Copernicus, diverge from craters, the locations of several craters have been derived on the far side of the moon.

Q. What does the symbol Δ mean when used with a number to label objects on a star chart?

A. These are southern clusters and nebulae in the catalogue of James Dunlop, who observed in Australia about 1825. Dunlop numbers are seldom used, as his list was superseded within a few years by John Herschel's catalogue.

Q. Which astronomical constants have been most precisely determined from observation?

A. The orbital periods of the earth and the moon, in terms of the length of the day.

Q. Who invented the telescope?

A. This question has never been fully settled. Apparently it was independently invented by several people at the beginning of the 17th century, among them Hans Lippershey, to whom credit is usually given. W. E. S.

GETTING ACQUAINTED WITH ASTRONOMY

TECHNIQUES OF LUNAR AND PLANETARY OBSERVING — II

AFTER the beginner has gained some acquaintance with the moon and planets, his growing experience is likely to suggest some specific observing programs, such as were described in the September issue. Whether his purpose is private enjoyment or adding to astronomical knowledge, the following hints should be helpful.

It is desirable to get in touch with other amateurs having similar interests, preferably by joining one or more of the major observing societies. The director of the Association of Lunar and Planetary Observers is Walter H. Haas, Pan American College Observatory, Edinburg, Tex. For information about the British Astronomical Association, write to the Assistant Secretary, 303 Bath Rd., Hounslow West, Middlesex, England. A new organization devoted to observations of the moon is the International Lunar Society, whose permanent secretary is A. Paluzie-Borrell, Diputacion 377, Barcelona, Spain. All three of these groups publish journals in which amateurs report their work. In addition, your local amateur society may have active observing sections.

An excellent habit for the observer is keeping a notebook in which every observation is written down at the time it is made, regardless of whether it seems important or not. Trust nothing to memory; a valuable discovery can be lost because the details were not written down when their recollection was fresh. Records should never be changed afterward. One convenient system is to keep the original in pencil and to make later additions in ink.

The practiced observer habitually records the time of observation for each entry in his notebook. On the moon, the advance of the sunrise or sunset line is so rapid that the appearance of details near it can change within a few minutes. The planets Mars, Jupiter, and Saturn can change their aspects rapidly as they rotate, so the exact moment of an observation must often be known for its proper interpretation.

In addition, notes should be made of the sky conditions at the time. We observe from the bottom of a deep ocean of turbulent air, which bends light rays irregularly as they pass through it. Consequently, stars are seen by the unaided eye to twinkle, and telescopic images vibrate and lose their sharpness. At the same time, there may be dimming by clouds, smoke, or dust. A careful distinction is drawn between *seeing*, which indicates the steadiness of the atmosphere, and *transparency*, which indicates its ability to transmit light.

It is customary to record the seeing on a scale of 0 (worst) to 10 (best). Similarly, sky transparency is often described on a scale of 0 (nothing visible) to 5 (very clear). The subject of seeing is discussed further on page 37 of this issue.

One reference work that is invaluable to the serious lunar or planetary student is the *American Ephemeris*, in particular the section containing the ephemerides for physical observations of the moon and planets. It gives complete information about conditions of solar illumination and the orientation of their surfaces with respect to the earth. Simplified compilations of such data are given in the annual *Handbook* of the British Astronomical Association, and in the *Observer's Handbook* of the Royal Astronomical Society of Canada.

Anything that adds to the comfort and convenience of the observer will increase the amount of work he can do in an evening. If the telescope does not already have a clock drive, adding one should be seriously considered. With a driveless instrument, objects drift rapidly out of the field of view, especially when high powers are used, and the frequent resetting of the telescope is a nuisance. Moreover, brief intervals of fine seeing can be missed because the observer's scrutiny has been interrupted.

Many amateurs have built their own drives at low cost. The first two volumes of *Amateur Telescope Making* contain many descriptions and pictures of homemade drives. Edgar Everhart discussed gear ratios for sidereal and other drive rates in SKY AND TELESCOPE for August, 1957, page 494 (out of print). Of course, many commercial models are available. Setting circles are less necessary for the planetary observer, who would need them only for locating Uranus or Neptune, or for observing the brighter planets by day.



An electric drive for a 10-inch telescope, made by the late J. H. Pruett.

The eyepiece of a long-focus Newtonian reflector can sometimes be awkwardly high above the ground, unless there is a suitable support or short sturdy stepladder available. There should be a convenient place close at hand for the observer's notebook, sharpened pencils, and charts. One accessory that can be strongly recommended is a box with a top of heavy ground glass, perhaps one foot square, and containing an electric light. With such an arrangement, drawing paper can be illuminated from below, making sketching the moon or planets much easier.



Several years ago, Thomas R. Cave, Jr., Long Beach, California, designed this observing platform and stand for his clock-driven 12½-inch f/9.7 reflector.

Radio Astronomy Receivers—I

FRANK D. DRAKE, *National Radio Astronomy Observatory*

AT FIRST SIGHT, the task performed by a radio telescope is incredible. The total power falling on the entire surface of the earth from the brightest radio source other than the sun, Cassiopeia A, is about 100 watts — just enough to operate a single light bulb of that rating. From this supply, a giant radio telescope will collect only about 10^{-16} watt.

The total radio power striking our earth in the famous 21-centimeter line of neutral hydrogen is scarcely 20 watts. Here a large radio telescope, receiving only a small fraction of the narrow band of available frequencies, may gather 10^{-17} watt. The faintest detectable radio sources provide our planet with but 1/100 watt of total power. Yet we can still manage to detect the millionth of a millionth of a millionth of a watt that is one telescope's share of this supply.

At the present time spectacular radio telescopes are springing up all over the world, especially in the United States. In each telescope, buried almost invisibly in the vast structure of the antenna, is a special radio receiver, usually called a *radiometer,* which detects and measures the radio energy collected by the antenna.

It is true that the resolving power is determined completely by the size and quality of the antenna, and these depend mainly on the money available to build it. But the faintness of detectable signals and the precision of radio intensity measures are largely controlled by the quality of the radiometer, and this depends chiefly on the ingenuity, knowledge, experience, and skill of the builder. For this reason, the design and construction of radiometers have appealed to some of the most nimble minds, who have brought to the field great vitality and fascinating advances.

Not only are the celestial signals weak, but they have no distinguishing features. Except for the 21-cm. sources, the radiation is at all frequencies, and the intensity of a particular frequency does not change with time, except in the case of solar system bodies. The radiation has the characteristics of random noise, just like static in a radio set. There are no dots and dashes, or tones, to tell us when we have found a signal. Furthermore, presently available receivers create far more noise power within themselves than is collected by the antenna. The problem facing the designer is to separate accurately the very faint celestial noise from that of the receiver, which is identical in nature but vastly stronger.

As a rule, the heart of the radiometer is a conventional receiver with all the usual tubes, coils, resistors, and other parts. To this is added electronic circuitry to increase stability and to distinguish signal noise from receiver noise. The conventional receiver performs the basic functions of all radios: It selects a certain band of frequencies, amplifies the power in that band, detects the amplified power, and presents the results in a convenient form.

The receiver design most frequently employed is the superheterodyne circuit, used in all home radios and TV sets, the incoming signal being mixed in a tube or crystal with a signal of nearly equal frequency generated internally. The two signals "beat," as in an out-of-tune piano, and the beat frequency, which now carries the signal information, is extracted, amplified, and detected. The advantage of this circuit is that the extracted beat frequency is much lower than the original signal frequency, and is more easily and cheaply amplified.

Until recently, the receiver output was universally recorded in graphical form by means of a moving-chart pen recorder. Lately, equipment has become commercially available that allows the output to be presented in digital form — as printed numbers or punched on tape in the proper code for direct insertion into an electronic computer. This latter method is rapidly gaining favor, as it eliminates the tedious work of measuring the pen recordings, avoids human error in data reduction, and enables the process to be carried out automatically by high-speed computers.



Fig. 1. The operation of an ideal radiometer, in which there would be no internal receiver noise, is compared with the situation in actual practice, where the output is a combination of external signal and internal noise.

Among the various complete radiometer systems, the most straightforward and obvious arrangement to extract signal from receiver noise is the *direct* or *total-power* radiometer. If a receiver is made extremely stable, the output power due to its noise will be nearly constant. When a radio source enters the antenna beam, there is a slight increase in output power, and this increase is a measure of the source intensity. This situation is shown in Fig. 1b.

In the direct radiometer a battery is used to provide a voltage equal to that of the receiver noise. These two voltages are applied to a circuit whose output is their difference. This differencing circuit thus acts to subtract out the receiver noise power, leaving only the increases in receiver output due to the presence of celestial radiation, as shown in Fig. 2a.

The proper performance of the direct radiometer depends entirely on high receiver stability. Because the receiver noise is being amplified, changes in the amplification, or gain, will cause fluctuations simulating changes in the input signal power. Since the receiver noise may be 10,000 times greater than the signal noise, it is necessary to have the gain vary by no more than one part in 10,000. Then the output fluctuations due to receiver gain variations will be less than changes due to the coming and going of celestial signals.

This is a very difficult standard to meet, and is a far cry from the one part in two or three that is acceptable in a radio or TV set. It is necessary to employ extremely stable electrical power supplies in radio astronomy receivers, particularly in the units that feed the tube filaments. Since slight variations in temperature affect gain, all receiver components in a direct radiometer are kept in rooms or containers that are held at very nearly constant temperatures, usually by means of heat pumps, air conditioners, or heaters controlled by sensitive thermostats. All this is costly, but provides the advantage of simplicity in the radio circuitry.

A variation of the direct radiometer, the *d.c.-comparison* type, is often used in 21-cm. work. Two separate bands of

frequencies are amplified by the receiver, separated, and detected individually. They are then put through the differencing circuit of a direct radiometer, one band replacing the battery. Since receiver noise and any wide-band cosmic noise are present equally in both channels, these are eliminated by the differencing circuit. However, if the receiver is tuned so that one band covers the 21-cm. line, the 21-cm. power will be in that band but not in the other. The output will then represent only the 21-cm. radiation being collected by the antenna.

A second, more sophisticated technique for overcoming receiver noise is the one used in the *switched* or *Dicke* radiometer (Fig. 2b), named after R. H. Dicke, who invented the method in 1946. The essence of this technique is to place a switch in front of the receiver, for alternate connection with the antenna and with a resistor. The signal power then comes to the receiver in pulses whose frequency and duration are controlled by the switch-frequency generator. Since the receiver is located after the switch in the circuit, the receiver noise does not have a pulsed character. The pulsing has, in effect, put a "tag" on the signal to distinguish it from the receiver noise.



Fig. 2. Principles of direct and Dicke radiometers, explained in the text.

Hence, if we put a device at the receiver output which recognizes only power with the proper tag on it, we may eliminate the receiver noise. Such a device is called a *synchronous detector*. It is connected to the switch-frequency generator, and detects only signals that are synchronized in time and duration with the switch. Its output is a direct-current voltage proportional to the amplitude of the pulsed signal applied to it. Such a radiometer is shown in Fig. 3.



Fig. 3. This Dicke radiometer, made by Ewen-Knight Corp., employs traveling-wave tubes (in left-hand unit), and operates at 8,000 megacycles with a bandwidth of 1,000 megacycles. The upper-center unit is the switch. All parts except the control rack (right) must be mounted on the telescope antenna.

Although the circuitry is more complicated in a Dicke radiometer, the gain-stability requirements are much less than in a direct radiometer. And in cases where the gain stability desired for the latter cannot be obtained with existing technology, the only answer is to use a Dicke radiometer. Disadvantages of this method, however, are that switches which do not waste signal power are often difficult or impossible to build, and that half the power collected by the antenna is thrown away because the receiver is connected to it only half the time.

A variation of the Dicke radiometer is also used in 21-cm. work. The receiver is connected to the antenna at all times, but the incoming energy is switched between the 21-cm. frequency and a nearby frequency. The receiver noise and wide-band cosmic noise are contained in both channels, and consequently not pulsed by the switching process, but the 21-cm. radiation is in only one band and is therefore pulsed, becoming the only power to appear in the final radiometer output. Such 21-cm. radiometers are necessarily about the most complex and expensive to be found in radio astronomy. An example of these receivers is the one used by Leiden Observatory astronomers to map the spiral structure of the Milky Way.

Would not perfect gain stability in the direct radiometer or the Dicke type allow us to detect infinitesimally faint signals? With existing electronic components, the answer is no. This inescapable situation exists because receiver noise *is* noise, a random sequence of voltages, and therefore not constant in intensity. Thus, receiver noise power in the output varies



Fig. 4. This installation at the National Radio Astronomy Observatory serves the 85-foot radio telescope (pictured on page 428, June issue). The control unit of Fig. 3 is at the left, and next to it a data digitizer. At right center, the large black assembly is a direct radiometer for 20 to 27 centimeters, and a d.c.-comparison radiometer for 21 centimeters. At the far right is a 68-centimeter direct radiometer. National Radio Astronomy Observatory photograph.

with time even if the gain is absolutely constant; hence the output of a direct radiometer will vary. Similarly, fluctuations in receiver noise in the Dicke radiometer will often, just by chance, create a series of noise pulses that are synchronized with the switch-frequency generator. Here again, the radiometer output varies.

The formula that follows is probably the most famous in radio astronomy, and gives the apparent input power fluctuation, ΔP, actually caused by receiver noise:

$$\Delta P = P_R/(Bt)^{\frac{1}{2}}.$$

This value is equivalent to the fluctuation in radiometer input power that would cause the same changes in output power as are produced by receiver noise. P_R is the receiver noise power in the output divided by the gain. B is band-width, the band of frequencies accepted by the receiver, and t is the time over which the output is averaged.

In a way, measuring the average output power is like trying to determine the mean level of the ocean when it is wavy. We might do this by inserting a number of sticks in the water and reading the water level at each stick. As the ocean gets rougher, which is equivalent to an increase of the receiver noise in the formula above, our results become less accurate. But if we use more sticks in the water, equivalent to enlarging the band-width, we get a more precise answer. Similarly, by making measurements repeatedly over longer periods of time — equivalent to increasing the averaging time — our answer becomes more accurate.

The value of the receiver fluctuation given by the formula sets the ultimate limit on the sensitivity of any radiometer, since if we are to distinguish a signal it must obviously be stronger than the random fluctuations. The continuous effort to improve sensitivity is one of the most challenging and busy fields of radio astronomy. It is only because we have succeeded in reducing the fluctuations below the level of 10^{-14} watt mentioned earlier that we have been able to make radio observations at all, and our success in pushing sensitivities below 10^{-17} watt has made radio astronomy a booming science.

What can be done to improve sensitivity? The averaging time is limited by the interval an object to be observed is above the horizon. Thus, it cannot exceed something like 12 hours for many objects. However, such long averaging times are rarely used, because over long periods instabilities in electronic components produce fluctuations greater than those given by the formula. The band-width B is, of course, as large as possible. In superheterodyne receivers it cannot be made greater than about 10^7 cycles per second — values above this figure cause increases in receiver noise that lead to a net deterioration in sensitivity, despite the larger band-width. Typical superheterodyne receivers today may detect signal powers which are about 1/10,000 as strong as the receiver noise.

The effort to improve sensitivity through increasing band-width scored a major advance when a special amplifier known as the traveling-wave tube was invented. Quite different from ordinary vacuum ones, these tubes have the special characteristic of amplifying a very wide band of frequencies, for instance 10^9 cycles per second. Dicke radiometers containing very simple receivers using these tubes are becoming quite popular in radio astronomy. The receiver in Fig. 3 can detect the presence of a noise signal which increases the noise power output by only one part in 350,000!

Sometimes, however, as in 21-cm. work, the signal covers only a narrow range of frequencies. This sets a limit on the useful band-width, and the only way left to improve sensitivity is to reduce receiver noise. Much work has gone into this, mainly in an effort to improve the electronic components, such as tubes and crystals, that govern receiver noise.

Looking back over the short history of radio astronomy, it appears that sensitivity has been improved, on the average, by about 50 per cent every year. This means that each year we have been able to see about 20 per cent farther into space with a given antenna, or to study almost twice as much space as the year before. This rapid advancement may now go even faster. Larger antennas are being built more quickly. And in just the last year or so, solid-state physics has given radio astronomers an assortment of new "wonder devices" that promise startling improvements in receiver sensitivity. A description of some of these devices, which may bring another revolution in astronomy, will be given in the succeeding article.

(To be continued)

Amateur Astronomers

A Young Amateur's Observatory in Texas

NOBLE OBSERVATORY is the product of 3½ years of work which began when I was 14 years old, without much experience in telescope making. But the three volumes of *Amateur Telescope Making* provided me with enough information to assemble and mount a refractor large enough for detailed observations of the moon and planets.

First, a 6-inch achromatic lens was bought from a very good amateur optical worker, and an equatorial mounting procured. As presently set up, the refractor is equipped with a large finder, a 3-inch guide telescope, a 3-inch astrograph, a 35-mm. camera, an astro-camera, setting circles, and a clock drive. The last is unsatisfactory, as its output torque is insufficient to turn the telescope, so I am attempting to make another clock drive.

All of the accessories listed above, except the two cameras, were made by me from locally available materials. I provided the mounting with slow motions and clamps, and plan for the future a console unit at which the instrument can be set for any position on the sky.

I bought the building, finished as shown in the photograph, for $300. It is 10 feet in diameter, with a full-sized entrance door. One of the top sections opens to permit observing from the zenith down to about 30° altitude. Another section is for from 30° to the top of the door, and anything lower is observed through the open door. Mounted on 31 wheels, the entire observatory rotates on an angle-iron track.



The 10-foot-diameter building of Tommy May's observatory turns on 31 wheels. Inside is a 6-inch refractor.

If this project were repeated, the total cost of $900 would be much less, as I have learned a great deal. The observatory is named after Miss Charlie M. Noble, an astronomy teacher of Ft. Worth, Texas, who has pioneered in encouraging junior astronomers in this area. The observatory is open to all visitors, and correspondence concerning it is welcome.

TOMMY MAY
2422 Gibbins Dr.
Arlington, Tex.

WASHINGTON, D. C.

In the summer of 1958, several members of the National Capital Junior Astronomers began a systematic program of drawing the planet Mars. They used an 8-inch reflector and a 5-inch refractor for most of their work, which was completed last February.

From 47 drawings, an over-all map of Mars was compiled. A similar program is planned for next year. Junior groups who may be interested in conducting a project of this type are invited to write Roy R. Troxel, 3017 Cleveland Ave., Washington 8, D. C.

HUMBOLDT COUNTY, CALIFORNIA

Two amateur societies are in operation in Humboldt County, in northern California. In Eureka, the Astronomers of Humboldt recently purchased 11 acres of land for a possible observatory site. The secretary of the club is William N. Abbay, Jr., 1745 Margaret Lane, Arcata, Calif.

Founded two years ago, the Astronomers of Southern Humboldt in Fortuna has nine active members, who meet on the second and fourth Sundays at private homes. Its secretary is William Shreeve, P. O. Box 862, Fortuna, Calif.

STAMFORD, CONNECTICUT

The primary mirror of the 20-inch Cassegrainian-Maksutov telescope, described on page 622 of the September issue with the project of the Fairfield County Astronomical Society, will have a diameter of 24 inches. The 4-inch apochromatic spotting scope, another instrument of the observatory at the Stamford Museum and Nature Center, was designed by E. L. McCarthy, Perkin-Elmer Corp.

ALBUQUERQUE, NEW MEXICO

There are 12 members in the Albuquerque Astronomers. Interested persons should communicate with Dan Judd, 402 Central Ave., S.W., Albuquerque, N. M.

BURLINGTON, IOWA

Ten adults and two juniors comprise the Burlington Astronomy Club. The president is Jack R. Polson, 2214 Barrett St., Burlington, Iowa.

RIVERSIDE, CALIFORNIA

Nineteen amateurs have formed the Riverside Astronomical Society. More information is available from H. E. Kaiser, 4868 Hedrick Ave., Arlington, Calif.

THIS MONTH'S PROGRAMS

Cleveland, Ohio: Cleveland Astronomical Society, 8 p.m., Warner and Swasey Observatory. November 6, Dr. G. de Vaucouleurs, Harvard Observatory, "The Atmospheres of Mars and Venus."

Dallas, Tex.: Texas Astronomical Society, 8 p.m., Dallas Health and Science Museum. November 23, Michael Gardner, "External Galaxies."

New York, N. Y.: Amateur Astronomers Association, 8 p.m., American Museum of Natural History. November 4, Dr. George A. Morton, R. C. A. Laboratories, "New Eyes for Our Telescopes."

New York, N. Y.: Junior Astronomy Club, 8 p.m., Waverly building, New York University. November 20, Dr. Francis J. Heyden, S. J., Georgetown Observatory, "Observing Total Eclipses for Geodetic Measurements."

Washington, D. C.: National Capital Astronomers, 8:15 p.m., Commerce Department auditorium. November 7, Dr. Jack Green, North American Aviation Corp., "Geochemical Aspects of Lunar Exploration."

BARTLESVILLE, OKLAHOMA

A new member of the Mid-States Region of the Astronomical League is the Bartlesville Astronomical Society, which is comprised of 19 amateurs. The president is E. L. Clark, 2054 Johnstone, Bartlesville, Okla.

A MICHIGAN AMATEUR'S BASEMENT PLANETARIUM

SINCE August, 1957, I have developed a small, inexpensive planetarium, with many auxiliary pieces of equipment. Although the seating capacity is only four to six, over 100 persons view the demonstrations each year, many of them returning every two months when I change the lecture subject.

So that the stars as seen from either the Northern Hemisphere or the Southern may be shown, the Spitz, Jr., projector is equipped with interchangeable spheres. These have been pierced to project 3rd- and 4th-magnitude stars not included by the manufacturer, and small holes punched together give a rather realistic impression of the Milky Way.

Half of a Spitz, Jr., sphere forms the

David DeBruyn's basement planetarium. The dome illuminator is at left.

dome illuminator, using a flashlight bulb with a blue filter. The dome is an 8½-foot beach umbrella, painted white on the inside and with linen added around the edge to complete the hemisphere. A silhouette of downtown Muskegon has been cut out, and compass points are marked.

For the sun, moon, and five bright planets, an equatorially mounted auxiliary projector with interchangeable slides was constructed. Fourteen slides depict the moon's phases, and there are slides of comets, meteors, and artificial satellites. A separate projector is used for auroras and the zodiacal light.

Photographs and other exhibits are set up outside the planetarium chamber. A phonograph with two remote speakers can be operated from the planetarium switchboard for special sound and musical effects.

DAVID DEBRUYN
2221 Oak Ave.
N. Muskegon, Mich.

Planetarium Notes

(Most planetariums give group and special showings by appointment.)

BALTIMORE: *Davis Planetarium.* Maryland Academy of Sciences, Enoch Pratt Library Building, 400 Cathedral St., Baltimore 1, Md., Mulberry 5-2370.

SCHEDULE: (Sept.-June), Thursday, 7:15, 7:45, 9 p.m.; Saturday, 2 and 3 p.m. Admission free. Spitz projector. Director, Paul S. Watson.

BLOOMFIELD HILLS, MICH.: *McMath Planetarium.* Cranbrook Institute of Science, Bloomfield Hills, Mich.

SCHEDULE: Saturday and Sunday, 2:30 and 3:30 p.m.; Wednesday, 4 p.m. Spitz projector. In charge, James A. Fowler.

BOSTON: *Charles Hayden Planetarium.* Museum of Science, Science Park, Boston 14, Mass., Richmond 2-1410.

SCHEDULE: Tuesday through Friday, 11 a.m. and 3 p.m.; Friday, 8 p.m.; Saturday, 11 a.m., 2 and 3:30 p.m.; Sunday, 1:30, 2:45, and 4 p.m. Korkosz projector. Director, John Patterson.

CHAPEL HILL: *Morehead Planetarium.* University of North Carolina, Chapel Hill, N. C.

SCHEDULE: Daily, 8:30 p.m.; also at 11 a.m. and 3 p.m. Saturday, 3 and 4 p.m. Sunday. Zeiss projector. Manager, A. F. Jenzano.

CHARLESTON, W. VA.: *Hillis Townsend Planetarium.* Children's Museum, Public Library Building, Charleston, W. Va.

SCHEDULE: Saturday, 11 a.m. Admission free. Spitz projector. Director, Mrs. R. L. Sullivan.

CHATTANOOGA, TENN.: *Clarence T. Jones Observatory.* University of Chattanooga, Brainerd Rd., Chattanooga, Tenn., MA 2-5733.

SCHEDULE: Friday, 8 p.m. Admission free. Jones projector. Astronomer in charge, Karel Hujer.

CHICAGO: *Adler Planetarium.* 900 E. Achsah Bond Dr., Chicago 5, Ill., Wabash 2-1428.

SCHEDULE: Monday through Saturday, 11 a.m. and 3 p.m.; Tuesday and Friday, 8 p.m.; Sunday, 2 and 3:30 p.m.; Tuesday through Friday, 10 a.m., special school program. Zeiss projector. Acting director, Robert I. Johnson.

COLORADO SPRINGS: *Academy Planetarium.* U. S. Air Force Academy, Colorado Springs, Colo., Granite 2-2779.

SCHEDULE: Wednesday, 8 p.m.; Saturday, 2:30 p.m. (except November 14th, 21st, and 28th); Sunday, 2 and 3:15 p.m. Admission free. Spitz Model B projector. Director, Maj. Richard J. Pfrang.

DALLAS: *Dallas Planetarium.* Dallas Health Museum, Fair Park, Dallas 10, Tex., HA 8-8351.

SCHEDULE: Saturday and Sunday, 3 p.m. Spitz projector. Planetarium educator, Mrs. Claudia Robinson.

DENVER: *Denver Museum of Natural History Planetarium.* City Park, Denver, Colo., East 2-1808.

SCHEDULE: Saturday and Sunday, 1 to 4:30 p.m. Spitz projector. Curator, W. R. Van Nattan.

FLINT, MICH.: *Robert T. Longway Planetarium.* Flint Junior College, 1310 E. Kearsley St., Flint 3, Mich., Cedar 8-1631.

SCHEDULE: Tuesday through Sunday, 8 p.m.; Saturday and Sunday, 2 p.m. Spitz Model B projector. Director, Maurice G. Moore.

FT. WORTH: *Charlie M. Noble Planetarium.* Ft. Worth Children's Museum, 1501 Montgomery, Ft. Worth, Tex., PE 2-1461.

SCHEDULE: Tuesday through Friday, 4:15 p.m.; Saturday, 11 a.m., 2:30 and 3:30 p.m.; Sunday, 2:30 and 3:30 p.m. Spitz projector. Supervisor, Norman C. Cole.

INDIANAPOLIS: *Holcomb Planetarium.* Butler University, Indianapolis 7, Ind.

SCHEDULE: Saturday and Sunday, 4 and 8 p.m. Spitz projector. Director, H. Crull.

KANSAS CITY: *Kansas City Museum Planetarium.* 3218 Gladstone Blvd., Kansas City 23, Mo., Humboldt 3-8000.

SCHEDULE: Saturday and Sunday, 3 p.m. Spitz projector. Director, Wilber E. Phillips.

LANCASTER, PA.: *North Museum and Planetarium.* Franklin and Marshall College, Lancaster, Pa.

SCHEDULE: Tuesday and Thursday, 8 p.m.; Saturday and Sunday, 2 and 3 p.m. Admission free. Spitz projector. Curator, John W. Price.

LAQUEY, MO.: *Tarbell Planetarium.* Inca Cave Park, Laquey, Mo.

SCHEDULE: Sunday, 1 to 6 p.m., continuous. Spitz projector. Director, E. D. Tarbell.

LOS ANGELES: *Griffith Observatory and Planetarium.* Griffith Park, P. O. Box 27787, Los Feliz Station, Los Angeles 27, Calif., Normandy 4-1191.

SCHEDULE: Daily (except Monday), 3:30 and 8:30 p.m.; also 2 p.m. Saturday and Sunday. Zeiss projector. Director, C. H. Cleminshaw.

MINNEAPOLIS: *Science Museum.* Minneapolis Public Library, 1001 Hennepin Ave., Minneapolis 3, Minn.

SCHEDULE: Saturday, 10 a.m. and 2 p.m. Admission free. Spitz projector. Planetarium director, Mrs. Maxine B. Haarstick.

NASHVILLE: *Sudekum Planetarium.* Children's Museum, 724 2nd Ave. S., Nashville 10, Tenn., Chapel 2-1858.

SCHEDULE: Sunday, 2:45, 3:30, 4:15 p.m. Spitz projector. Director, Jacqueline Avent.

NEWARK: *Newark Museum Planetarium.* 49 Washington St., Newark 1, N. J., Mitchell 2-0011.

SCHEDULE: Saturday, Sunday (except 1st Sunday of month), and holidays, 2:30 and 3:30 p.m. Admission free. Spitz projector. Supervisor, Raymond J. Stein.

NEW YORK CITY: *American Museum-Hayden Planetarium.* 81st St. and Central Park West, New York 24, N. Y., Trafalgar 3-1300.

SCHEDULE: Monday, 2 and 3:30 p.m.; Tuesday through Friday, 2, 3:30 and 8:30 p.m.; Saturday, 11 a.m., 1, 2, 3, 4, 5 and 8:30 p.m.; Sunday and holidays, 1, 2, 3, 4, 5 and 8:30 p.m. Zeiss projector. Chairman, J. M. Chamberlain.

PHILADELPHIA: *Fels Planetarium.* Franklin Institute, 20th St. at Benjamin Franklin Parkway, Philadelphia 3, Pa., Locust 4-3600.

SCHEDULE: Tuesday through Sunday, 3 p.m.; Saturday, 11 a.m.; Saturday, Sunday, and holidays, 2 p.m.; Wednesday and Friday, 8 p.m. Zeiss projector. Director, I. M. Levitt.

PITTSBURGH: *Buhl Planetarium and Institute of Popular Science.* Federal and West Ohio Sts., Pittsburgh 12, Pa., Fairfax 1-4300.

SCHEDULE: Daily, 2:15 and 8:30 p.m.; also at 11 a.m. Saturday and 4:15 p.m. Sunday. Zeiss projector. Director, Arthur L. Draper.

PROVIDENCE: *Roger Williams Planetarium.* Roger Williams Park Museum, Providence 5, R. I., Williams 1-5640.

SCHEDULE: Saturday, 3 p.m.; Sunday and holidays, 3 and 4 p.m. (Oct. 1-May 30). Admission free. Spitz projector. Director, Maribelle Cormack.

SAN FRANCISCO: *Morrison Planetarium.* California Academy of Sciences, Golden Gate Park, San Francisco 18, Calif., Bayview 1-5100.

SCHEDULE: Daily (except Monday and Tuesday), 3:30 and 8:30 p.m.; also 2 p.m. Saturday, Sunday, and holidays. Academy projector. Curator, George W. Bunton.

SAN JOSE, CALIF.: *Rosicrucian Planetarium and Science Museum.* Park and Naglee Aves., San Jose, Calif.

SCHEDULE: Sunday and Wednesday, 2 and 3:30 p.m. Spitz projector. Director, Rodman R. Clayson.

SANTA BARBARA, CALIF.: *Gladwin Planetarium.* Museum of Natural History, 2559 Puesta del Sol Rd., Santa Barbara, Calif., WO 6-6720.

SCHEDULE: 1st and 3rd Monday, 3 p.m.; 2nd and 4th Thursday, 8 p.m. Admission free. Spitz projector. Lecturer, C. Adair.

SPRINGFIELD, MASS.: *Seymour Planetarium.* Museum of Natural History, Springfield 5, Mass.

SCHEDULE: Tuesday, Thursday, and Saturday, 3 p.m.; also 8:30 p.m. Tuesday; special star stories for children, Saturday, 2 p.m. Admission free. Korkosz projector. Director, F. Korkosz.

STAMFORD, CONN.: *Edgerton Planetarium.* Stamford Museum and Nature Center, Stamford, Conn., Davis 2-1646.

SCHEDULE: Saturday, 11 a.m.; Sunday, 4 p.m. Spitz projector. Director, Ernest T. Luhde.



"About a month ago an astronomical group had a number of telescopes set up for a public viewing of the moon and planets. There were several refractors of 4 and 5 inches and a 12-1/2-inch reflector. I looked through them all and then set up my Questar. I was very pleased with what I saw. None of the other instruments seemed to produce the image my Questar did. I am very pleased with its performance, and I am sure no other telescope is so easy to carry and set up and use."

—S. Paul Jones, Louisville, Kentucky

It is very pleasant to receive a phone call from a Questar owner several hundred miles distant who says nice things like this about our product. It happens, however, with a frequency we find most gratifying. On this agreeable occasion last June we made bold to ask Mr. Jones if we could quote him in some way, to which he replied that he would put it in writing at once, and would be most pleased to have us use it as a signed testimonial.

That is how this ad commenced. Mr. Jones became a Questar owner early this year, and very quickly found himself taking excellent pictures, like the one above of the central portion of the moon. The exposure was 4 seconds, on 35-mm. Panatomic film, developed in Ethol. An inexpensive Praktica camera body was used, with eyepiece projection by our standard dual-purpose coupling which permits great increases in either photographic focal length or visual power. This coupling is shown on page 26 of the Questar booklet.

At right is Questar, with a Hexacon camera, attached to a black-anodized auxiliary base plate, which is, in turn, supported by the new Linhof Heavy Duty Pro Tripod and Deluxe Pan Head. This is the wonderful new tripod about which we are so enthusiastic, for it is the first one we can endorse wholeheartedly. Old friends will recall our reservations about the stiffness of the previous Linhof tripod when fully extended. This one is adequately steady at all heights, looks as though it were made for Questar, and costs only $139.50, which is $30 less than its predecessor. Weighs only 17 pounds, has new quick leg adjustments, removable braces, and folds to only 36 inches. Big rubber tips screw down to keep the leg spikes from hurting people, floors, or cars.

The new Deluxe Pan Head is better than ever, and costs $59.95. Note how entire instrument is offset, with nothing near the camera. In southerly latitudes, where the polar axis nears the horizontal, such a support is hard to beat.

Questar is priced at $995, postpaid, complete with English leather case and luggage cover. It is mailed in a shockproof reusable Leverpak drum. Literature will be sent upon request. Extended payments are available.

Questar

NEW HOPE · PENNSYLVANIA

"See-thru"

STAR MAPS

Luminous stars printed on clear plastic make star identification easy by direct comparison of "See-thru" map and sky. Twelve maps systematically cover entire sky. All navigational stars included from the *American Ephemeris* (1960 co-ordinates). Unique constellation figures using straight lines. Gnomonic projection.

Set of 12 maps ..... $4.95 ppd.

BRODER MAPS

1368 McAllister St., San Francisco 15, Calif.

THE ASTRONOMER KIT

★ FOUCAULT TESTER, in its entirety, with light source, knife-edge, measuring scale
PYREX MIRROR BLANK
★ NEW, NONVITREOUS TOOL, faster, smoother, more accurate
SEVEN ABRASIVES for greater safety and speed
CERIUM OXIDE for fast, clean polishing
SPECIAL OPTICAL PITCH to minimize danger of turned-down edge
MAGNIFYING LENS, essential in mirror grinding
★ DETAILED, ILLUSTRATED INSTRUCTIONS, "Mirror Making in the Kitchen"

(Starred items exclusive with The Astronomer Kit)

6-inch kit.....$15.00 ppd.
4¼-inch...$11.50 ppd.; 8-inch...$24.50 ppd.
Foucault tester only.....$6.00

Money-back guarantee.
New catalogue available soon.

ASTRONOMY, INC.
Suite 305, American National Bank Bldg.
Denver 2, Colorado

The NEW FI-NIS Off-Axis Catadioptric Telescope

for the discriminating amateur astronomer

utilizes a Schmidt-type, off-axis correcting lens finished to the highest telescopic accuracy

To achieve unexcelled optical resolution, with image quality matching that which is theoretically possible for its aperture.

This new telescope, embodying a design never before commercially available, enables the realization of a rugged, closed-tube construction whose parts are solidly and permanently aligned — including the built-in wide-field finder.

Its 4¼-inch aperture is *entirely free* of obstructions — its image *completely achromatic*. The flawless color fidelity and brilliant contrast are striking at first glance — a result of its well-made off-axis construction. This is a *really different* telescope.

The price, without mounting, is presently $400.00 subject to quotation. Write us for detailed information.

Other products: **SOLA-FIL** — an unusual solar filter

TELEWIN — a precision window for reflecting telescopes

OPTRON LABORATORY

Box 25, D. V. Station
Dayton 6, Ohio

OBSERVER'S PAGE

Universal time (UT) is used unless otherwise noted.

Observing the Moon — Parry and Bonpland

LOCATED within the broad level expanse of Mare Nubium, Parry and Bonpland are members of a chain of ruined craters extending from Fra Mauro on the north to Guericke on the south. This group is seemingly older than much of the lunar surface as we now see it. While showing some traces of their former imposing magnificence, all of these formations are in some degree dilapidated.

Parry is 28 miles in diameter, according to J. Young's measurements, and is noticeably hexagonal in shape. Its walls for the most part are low, and are broken by several passes from the interior to the surrounding plain, the widest being on the west and north. The eastern wall is highest, forming a ridge that separates the floor from that of Bonpland. This ridge has several peaks reaching heights of perhaps 5,000 feet or more; a small, deep crater at its southern end is labeled Parry E in the International Astronomical Union's lunar atlas.



The lunar craters Parry (left) and Bonpland, drawn by Alika K. Herring on June 26, 1958, at 4:50 Universal time. He used a 12½-inch reflector with a power of 228x. Sunrise on this portion of the moon's surface had occurred about 16 hours before. South is at the top, east to the right.

Bonpland, 37 miles across, is somewhat larger than Parry and has apparently suffered more erosion. The walls are lower, particularly on the east, where they degenerate into a broken chain of little hills. The approximate circularity of the formation (its outline has indications of polygonal structure) is broken by a large bay on the southwest. This was perhaps an ancient crater that once formed a complete ring, but the north and east walls are now missing, and I have been unable to detect the least trace of them.

The interiors of Parry and Bonpland are crossed by many rills that form part of a much more extensive system. One easily seen cleft enters Parry through a pass in the west wall, runs diagonally across its southern floor, and bends sharply southward to emerge through an opening in the south wall.

Another conspicuous cleft crosses this crater in a north-south direction near its east wall, and passes through Parry E. A northward extension cuts through the wall, and runs for some distance across the floor of adjoining Fra Mauro. The southward extension across the bay in Bonpland was hidden by the shadow of the wall at the time my drawing was made. South of Bonpland, this cleft continues some distance across the plain to end at a small crater which is designated A on H. P. Wilkins' map.

These are easy rills, and should be readily visible in 3- or 4-inch telescopes. Another, more difficult cleft traverses the eastern side of Parry's floor in a wide semicircle, concentric with the wall. To be well seen, this very delicate cleft requires fairly large apertures, optimum seeing and illumination. It is a test object for an 8-inch telescope.

The most conspicuous cleft in Bonpland crosses the floor centrally from south to north, beginning in the plain to the south at a craterlet marked C on the Wilkins map. North of the midpoint of Bonpland the cleft narrows and becomes difficult to see, but can be traced as far as the north wall. Near there, it is crossed by a delicate cleft originating



The area of Parry, Bonpland, and Fra Mauro is shown here in a reproduction of part of H. P. Wilkins' 300-inch map of the moon. South is at the top, and the grid spacing is 0.02 of the moon's radius. Compare the cleft pattern with that in the drawing opposite.



White WINDMASTER

No. 409. Two matching dials, with 5½" mounting diameter, show wind speed up to 120 miles per hour and wind direction around full 360°. Speed dial has HI-LOW range switch and is self-powered. Direction dial has ON-OFF switch and requires either 110 volts a.c. or 6-volt battery. Complete with 5' mast, 60' cable, and hardware. Available in brass or chrome.

$122.50

Units available separately:

Speed, $55.00 **Direction, $67.50**

Write for complete catalogue.

SCIENCE ASSOCIATES

Instruments/Weather • Astronomy/Teaching Aids

194 Nassau St., P. O. Box 216, Princeton, N. J.

YEAR-END SALE!

151 POWER SPACE SCOPE

Now . . . a 2.4-inch refracting telescope with the precision qualities of much larger instruments, that will enable you to explore the depths of space, to see the rings of Saturn, the craters of the moon, the belts of Jupiter and many double stars! When you observe terrestrially with the Space Scope, you can make out human figures at distances up to 24 miles, distinguish a person's general appearance at 6 miles, and make an individual identification at 2½ miles.

OBJECTIVE LENS . . . hard-coated Fraunhofer type. Over-all diameter, 62.5 mm.; clear aperture, 60 mm. Astronomical eyepieces: 151x Huygenian-Mittenzwey, 73x Huygenian-Mittenzwey, 35x Kellner, 28x Achromatic Huygenian. Terrestrial eyepiece: 30x.

COMBINATION ALTAZIMUTH-EQUATORIAL MOUNT enables scope to be used as an altazimuth or an equatorial instrument. It has three axes, giving horizontal motion if desired, as well as slow motions in right ascension and declination.

DIAGONAL PRISM (right angle) permits greater ease in astronomical observations.

ACCESSORIES include sunglass, 6x finder with crosshairs, 10-cm. sun projection screen, equipment tray. Scope is about seven feet high when fully extended. Complete with hardwood carrying case and 10-page instruction folder.

REGULARLY $89.50 **$69.50**

TO ORDER: Send check or money order. Shipped freight collect. For C.O.D. send 1/3 deposit or more. Calif. resid. add 4% state tax. 10-day money-back guarantee applies to everything.

ALL LENSES OF HIGHEST JAPANESE OPTICAL QUALITY

the akron

4394 SUNSET BOULEVARD • LOS ANGELES 29, CALIFORNIA

MASTERPIECES

of precision performance!

This Deluxe 6-inch model has removable tripod inside its massive 45 lb. pier

You'll find these fully professional features in every Custom DYNASCOPE®!

- Exquisite optical systems, fully achromatic and corrected to meet the most critical research standards.
- Massive equatorial mounts that provide exceptional stability and embody the latest mechanical refinements for ease of operation.
- Full sets of matched orthoscopic and achromatic eyepieces with wide flat fields sharp to extreme edges.
- Needle bearings carefully fitted on substantial shafts to assure movements so fine that they equal the ease and precision of a fine watch.
- Lifetime materials throughout, for atmospheric resistance and dimensional stability.
- Fully rotating tubes, with counterpoise to provide perfect balance during rotation.
- Rotary secondary supports to minimize diffraction.
- Exclusive Dyna-mesh rack and pinion focus for smoother, more accurate focusing action.
- Two-way pier tripods (Pat. Pending) in 6-inch and 8-inch models. 45-pound pier can be mounted permanently. Portable tripod is instantly removable for field trips.
- Highly accurate electric drives, free from backlash, with fully compensating clutch.
- 8 x 50 coated achromatic finderscopes with crosshairs and extra-large 6° fields.
- 7"-diameter setting circles spoked for easy setting. Graduated with verniers.
- Fully adjustable to all latitudes.

CUSTOM DYNASCOPES®

in your choice of
6"-8"-10"-12"-16" models

Every Custom DYNASCOPE – from 6" to 16" – is designed and built to meet such high standards and exacting specifications that it is the overwhelming choice of schools, colleges, universities, and many well-known professional astronomers. Each of these magnificent instruments incorporates features that are years ahead in optical engineering, yet based on time-tested principles. Each is painstakingly produced and assembled by craftsmen who put careful workmanship before speed. Each is required to pass, flawlessly, a long series of rigid performance and operational tests. Nothing is overlooked to make sure that these superb telescopes are the finest in their class, regardless of price. Whichever size you select, we guarantee that it *must* measure up to this standard or your purchase price will be refunded in full. You can order with absolute confidence . . . and with the exciting expectation of new thrills and new experience ahead.

Ask About Our "Easy-To-Own" Payment Plan

YOURS FREE: This Colorful Brochure

Before you decide on *any* telescope, send today for this fascinating new booklet that gives you the full specifications and prices of all Custom DYNASCOPES, plus helpful information to guide you in selecting your next telescope. A postcard will bring your copy FREE!

6-INCH STANDARD **DYNASCOPE®**
As Above But Without Electric Clock Drive, Setting Circles and Pier
f.o.b. Hartford, Shipping Wt. 80 lbs.
$265 complete

6-INCH DELUXE **DYNASCOPE®**
$475 complete f.o.b. Hartford, Shipping Wt. 150 lbs.

CRITERION Manufacturing Co. · MANUFACTURERS OF QUALITY OPTICAL INSTRUMENTS
Dept. D-31, 331 Church St., Hartford 1, Connecticut

© Copyright 1959 Criterion Mfg. Co.
®TM Registered U.S. Pat. Office

on the northeast part of the floor. After the intersection, this second one makes a sharp bend to the northwest, passing through the wall onto the floor of Fra Mauro.

Another easily seen cleft starts just south of Parry E and extends southeastward across the bay in Bonpland, but before reaching the wall it narrows and disappears on the floor.

Many other fine details are contained inside Parry and Bonpland, but are well seen only when thrown into strong relief by grazing solar illumination. Besides a number of small craterlets and hills, there are some very low ridges, generally oriented north and south. Of special interest is the curiously curved row of several saucerlike depressions, lying midway between the center and northeast wall of Bonpland.

ALIKA K. HERRING
1312 Arlington Ave.
Anaheim, Calif.

NEW VARIABLE STARS

Two recently discovered variable stars are bright enough for profitable observation in binoculars. One of them is HD 186776, in Cygnus at $19^h\ 43^m.1$, $+40°\ 36'$ (1950 co-ordinates), a 6th-magnitude object which varies by 0.7 magnitude. The other is HD 203378, in Cepheus at $21^h\ 18^m.2$, $+55°\ 14'$, which varies by 0.5 magnitude around an average value of 7.

Both stars are red, of spectral type *M*, and have light curves like that of Mu Cephei, according to the discoverer of their variability, W. Strohmeier of Remeis Observatory at Bamberg, West Germany. He first noted them while comparing patrol photographs. The two objects are shown in the Skalnate Pleso *Atlas of the Heavens*, and the first is noted as red in Norton's *Star Atlas*.

OBSERVATIONS OF A NOVA IN OPHIUCHUS

Discovery of an 8th-magnitude nova in Ophiuchus by G. Haro, of Tonanzintla Observatory in Mexico, was reported on page 343 of the May, 1958, issue of Sky and Telescope. With the aid of the photograph of the region published with that announcement, I found it easy to locate the star and follow its light changes visually.

Between June and November, 1958, the nova showed well-marked oscillations in brightness, between magnitudes 11 and 12½. By February, 1959, when the star could again be observed in the morning sky, it was magnitude 13, fading very gradually to 13½ by the beginning of September. These observations will be published in detail in a *Circular* of the variable star section of the Royal Astronomical Society of New Zealand, after accurate magnitudes for the comparison stars become available.

ALBERT F. JONES
40 Trafalgar St.
Timaru, New Zealand

Read This Advertisement

Here is a combination of a Barlow and a particular ocular which gives outstanding results. It consists of our new Barlow and our 16.3-mm. (⅝" focal length) Erfle eyepiece. While the Barlow was not specifically designed to work with this eyepiece, it does so to an astonishing degree. All images are sharp and hard to the very edge of the field.

The Barlow gives magnification up to slightly over three times that of the ocular alone. It is achromatic, coated, and mounted to the U. S. standard size of 1.250 inches.

The modified Erfle eyepiece has a field of 75 degrees with excellent eye relief. The combination gives the equivalent focal length of slightly under 6 millimeters. Many users state it is far superior to any shorter focal length ocular of equivalent magnification.

The Barlow sells for $16.00 postpaid, and the Erfle for $14.75 postpaid. Both are guaranteed to perform as stated above or money refunded.

ORTHOSCOPIC OCULARS—All hard coated, standard 1¼-inch outside diameter.

28-mm.	$15.00	10.5-mm.	$16.25	4-mm.	$17.25
16.3-mm. (Erfle)	$14.75	7-mm.	$17.25	Barlow 3x	$16.00

Telescopics
1000 North Seward Street
Los Angeles 38, Calif.

Warranted to equal or surpass any oculars obtainable anywhere or money refunded.

Finished mirrors, mirror kits, spiders, elliptical flats, focusing devices, aluminizing.

Send for catalogue.

EXTRA-THICK PYREX ELLIPTICAL DIAGONALS

Aluminized with quartz over-coat

Fine annealed Pyrex. Minor- and major-axis tolerance, 45 degrees ± 5 minutes. Angular dimensions, +0%, −½%. Flatness, ⅛-wave-length sodium light. Surface quality, pitch polished, free of significant chips, scratches, pits, or bubbles.

Minor Axis	Thickness	Polished blank	Unpolished blank
1½"	½"	$ 8.00	$2.15
1¾"	½"	10.00	3.33
2"	½"	14.95	4.50
2½"	⅝"	17.95	5.40
2¾"	¾"	23.95	7.25

Discounts to quantity purchasers.

SCHOTT-JENA OBJECTIVE BLANKS

Made by Jena Glass Works, Germany, each pair consists of one crown-glass blank, 1.516 refractive index for sodium light and 64.0 dispersion, and one flint-glass blank, 1.620 refractive index and 36.3 dispersion.

Diam.	Grade A	Grade B	Diam.	Grade A	Grade B
4"	$23.10	$12.10	6½"	$ 84.70	$46.20
4½"	29.70	15.90	7"	104.00	57.20
5"	38.50	20.90	7½"	242.00	—
5½"	52.80	28.60	8"	297.00	—
6"	63.80	35.20			

Grade A blanks are fine annealed and tested for striae — the best grade they make. Grade B are cast glass which may contain some fine striae.

Four formed steel tools cut to your radius specifications on order for approximately $30 in 5" diameter to $75 in 8" diameter.

SLIDE FOCUSER

Completely new, never offered before. Highly recommended for small reflectors, economical yet durable and light. Frame is made of accurately spot-welded steel which is plated and has mounting holes. Aluminum tube takes 1¼" eyepieces — is highest quality obtainable commercially with extra thickness. Long-lasting construction. 15-day-refund guarantee.

$2.95 ppd.

COATED ACHROMATIC TELESCOPE

40-mm. aperture, 20x magnification with 2-degree field. Fainter than 10th-magnitude stars visible — better than 3 seconds of arc resolution! Makes a superb spotter for the beginner — better than a field glass. Mounts and dismounts on small tripod with ease. Readily adaptable for use as finder scope. Complete with erecting prisms and tripod.

$16.95 ppd.

BAUSCH & LOMB RED ROUGE

This is a fine red rouge which is sold only in factory-sealed cans. For clubs, makers of large mirrors, and commercial use, where a high-quality polish is desired. 5-lb. can, $8.00 packed, ppd.

CERIUM OXIDE

Optical grade for fast-polishing action. Use 3 or 4 ounces for 6" mirror. In sealed plastic bag.

25¢ per ounce

MIRROR CELLS

Solid cast aluminum, fully adjustable, painted, complete and ready to use. Shipped postpaid.

6" mirror cell for	7"	or larger tube	.....	$ 6.75
8"	"	9¼"	"	..... 8.95
10"	"	11¾"	"	..... 15.95
12½"	"	14½"	"	..... 20.95

BULK GRINDING ABRASIVES

This fast-cutting aluminum oxide has shaped grains that bite into the work for minimum grinding effort. It is tougher and stands up longer under severe usage. Excellent for both hand- and machine-grinding. Individually packaged quantities for shipment.

Grain size	5 lbs.	25 lbs.	50 lbs.	100 lbs.
#60	$2.25	$10.00	$16.00	$25.00
#150	2.50	11.25	19.00	35.00
#220	2.85	13.00	23.00	40.00

All quantities shipped express collect.

Write for free catalogue.

NYE OPTICAL CO.
2100 Cherry Avenue, Long Beach 6, Calif.
Telephone: GEneva 9-1373

Store hours: Daily, 9 a.m.-5:30 p.m.; Saturday, 9 a.m.-2 p.m.

Frontiers are extended by the practical visionary

It is the practical visionary who has given us much of what we enjoy today. And it will be the visionary—the man with ability to seek concepts beyond the existing limitations of science—who will guide our developments of tomorrow.

The Applied Physics Laboratory (APL) of The Johns Hopkins University seeks men who will be engaged in advanced research problems—who will find solutions to problems yet to be posed. Their findings will provide guidelines for the space and missile hardware research of the future.

Your endeavors will be heightened by the professional atmosphere of APL. This atmosphere, created by men dedicated to the furtherance of science, has earned APL a reputation as a leader in programs vital to the national security.

Appointments at APL offer exceptional opportunities. For detailed information, address your inquiry to:

Professional Staff Appointments

The Johns Hopkins University Applied Physics Laboratory

8635 Georgia Avenue, Silver Spring, Maryland

Astronomical Seeing

EVERY OBSERVER with a 3-inch telescope or larger is uncomfortably aware that on some nights the moon, planets, or close double stars do not yield satisfactory images. The user of binoculars, however, generally has sharp, clear views, even when his colleague with a powerful telescope complains of poor seeing.

The term *seeing* is used to indicate the departure of the telescopic image from its ideal form. A commonly employed seeing scale ranges from 0 (hopelessly blurred) to 10 (perfect in every way). Thus, seeing 5 characterizes average conditions, in which a fair-sized amateur telescope will show considerable detail on Mars or Jupiter, but with enough atmospheric disturbance to prevent fully efficient use of a good optical system. However, all experienced planetary observers will bear poor-to-average seeing for hours on end in order to enjoy the brief moments when the image steadies down, permitting full exploitation of the telescope.

Ideally, a lens or mirror brings rays from a point on a distant object to a point in the focal plane of the telescope. Air, being a refracting medium, affects the path of the light rays according to its density, and is as much a part of the optical system as the telescope itself. Anything that alters the homogeneity of the long column of air through which the telescope looks will distort the path of the light rays.

For the purposes of the visual or photographic observer, three distinct types of seeing (I, II, and III) may be distinguished, but they can also occur in combination. This threefold classification is helpful in analyzing what can be done about seeing, for it is one of the knottier problems facing the practical observer.

In poor seeing of type I, the image changes rapidly, and small objects such as stars or Jupiter's satellites occasionally appear double or triple. On the moon or planets, the disk may show two or more distinct boundaries rapidly moving or vibrating, and surface detail seems jumpy. In addition, an out-of-focus image of a bright star shows boiling or moving streakiness.

The cause of these phenomena is air currents within or very close to the telescope, with a wave length of about one-tenth to one-half the diameter of the mirror or objective. These moving inhomogeneities within the optical path prevent rays from all parts of the mirror or objective from reaching the focus simultaneously. The effect is much as if the telescope consisted of several small mirrors in relative motion. When this trouble is present, photographic work of any delicacy is out of the question, no matter how short the exposure.

Type-II seeing is characterized by images that are steady but blurred or fuzzy. There is poor contrast on extended surfaces, and star images appear swollen, often to several times their theoretical size.

Usually type-II poor seeing results from air currents outside the telescope tube, but low-lying and often in fairly rapid motion. The wave lengths of these disturbances are small, perhaps less than half the diameter of the mirror or lens. These conditions may arise during the night cooling of air, when stable temperature layers dissipate in the presence of natural or artificial objects having different cooling rates (houses, rocks, pavement). A change of wind direction with increasing altitude may produce the same effect, if the vertical temperature gradient is steep. The passage of weather fronts often produces violent type-II disturbances. Under such conditions, lunar or planetary photography would be hopeless for any observer.

Finally, in seeing of type III the images are sharp but somewhat unsteady. A star image is crisp and clear, yet in motion; the edge of the moon ripples to some degree, although it is sharply defined. These are the effects of high-level atmospheric inhomogeneities, whose wave lengths are large compared with the aperture of the telescope, perhaps three or four times as large, or more. When

SEE MORE WITH YOUR TELESCOPE

Recently a customer purchased a new Cal-Astro 8" f/7 mirror and diagonal to replace those in the telescope he was then using (a widely advertised American make). After testing he writes in part:

> . . . looking at Saturn last night, using a 4-mm. eyepiece (about 360 power) . . . Cassini's division was so black and sharp it could have been painted on with black paint. . . .
>
> . . . Some of the amateur astronomers I have talked to have the opinion that a mirror of extra-high quality is only of value when the seeing is unusually good. . . . I can positively say that this is not the case. The performance is better regardless of the seeing, and I think that anyone who is really interested in astronomy and does not get the very best mirror he can get is doing himself an injustice. *(Name and address of writer upon request.)*

If you are not satisfied with the present performance of your telescope, let Cal-Astro help you to better viewing.

WRITE TODAY FOR ADDITIONAL DETAILS.

Cal-Astro
Optical Laboratories
Since 1947. Owned and operated by Leland S. Barnes.

30 North San Gabriel Blvd.
PASADENA, CALIFORNIA
Phone: SYcamore 2-8843

A time service for your telescope

HAINES ELECTRIC SIDEREAL CLOCK

The minute circle is 4 inches in diameter. The case is machined cast aluminum, polished and buffed; and the bronze bezel is polished and lacquered. Hour and minute hands are set with the knob at the bottom of the case. Computed accuracy is within 3.75 seconds a year slow (assuming the frequency constant). Please specify your current: 60-cycle, 110-volt; 50-cycle, 110-volt; or 60-cycle, 220-volt.

$53.00

Two other sidereal clocks available: **A.** 8-inch-diameter dial (dimly illuminated from behind), which may be bolted to your telescope pier. **B.** 14-inch-diameter minute circle mounted in a 17-inch wooden case.

USED AS A CHRONOGRAPH

Record your observations of celestial phenomena with the Sidereal or Solar Clock and your tape recorder. Signals of every second, every minute, and every hour can be broadcast as desired, by means of additional devices built in the clock. Write for estimates.

SPECIAL: Your Local Sidereal Time will be computed at your request free of charge for any time and place. Tell us your longitude, or write for instructions.

FOR THE STUDY OF CELESTIAL CO-ORDINATES and **QUICK** location and identification of celestial bodies, equip your telescope mounting with large co-ordinate circles. YOU SUPPLY THE CIRCLES, from 10 to 21 inches in diameter (preferably of aluminum). We graduate them by AUTOMATIC MACHINE, accurate to within 2 or 3 seconds of arc, to mark single minutes of right ascension and sidereal time and 15 minutes of arc in declination. Our 16-inch circles with verniers will give single minutes of arc for the declination and 4 seconds for sidereal time and right ascension. Write for prices and instructions on how to proceed. The 21-inch circles are readily adaptable to radio telescopes. Circles supplied if required.

Haines **SCIENTIFIC INSTRUMENTS**
Box 171, Englewood, New Jersey

U.S. SURPLUS BARGAINS

IN HIGH-QUALITY OPTICAL PARTS AND EQUIPMENT

BAUSCH & LOMB F/6 AERO-TESSAR LENS

Focal length 24". Completely mounted with iris and shutter. Excellent for wide-field telescope. All in excellent condition. Price **$25.00**

WIDE-ANGLE ERFLE EYEPIECE (68° FIELD)

Coated optics, 3 achromatic lenses. 1¼" focal length, 1-13/16" aperture. Price **$12.50**



16-mm. AERIAL GUN CAMERA, MODEL AN-N6

Takes standard 16-mm. magazine load (50'). Operates at 16, 32, or 64 frames per second. Complete with Wollensak Type V lens, 1⅜" (35-mm.) f/3.5. Excellent condition. Operates on 24-volt d.c. Price **$29.95**

TELESCOPE MOUNT

Rotating yoke with micrometer positioning tightens onto any ½" vertical shaft with a thumbscrew. Levels with built-in bubble; orients with built-in magnetic compass. Turned by knurled wheel on one end of worm shaft. A 1"-diam. micrometer-calibrated wheel on other end divides each coarse azimuth unit into hundredths. The coarse scale divides each 180° into 32 parts, which, with the micrometer, makes the standard gunnery circle of 6,400 mils. A direct-reading conversion graph of mils to degrees is furnished. Quick-disconnect lever disengages the worm for turning platform quickly. Yoke will mount ⅜" shafts 3¼" apart in split pillow blocks 3½" above platform. Bronze worm on a shaft with a knurled wheel is mounted below the outside of one pillow block. If the shaft through the block had a 2"-pitch-diam., 32-pitch single-thread, 64-tooth right-hand gear, the worm would mate with it for controlled tilt. Price **$5.00**

5" SCHMIDT ULTRA-HIGH-SPEED OBJECTIVE LENS SYSTEM

Eastman Kodak infrared receiver, formerly known as U. S. Navy Metascope, Type B. 7" long with 5" Schmidt ultra-high-speed objective lens (approx. f/0.5). Elaborate optical system, many coated lenses. Uses two penlight batteries. Gov't. cost, approx. $300.00. Factory-new. Shipping wt. 9 lbs. Price **$19.95** Waterproof carrying case extra. Shipping wt., 3 lbs. Price **$3.00**

NAVY INFRARED RECEIVER, TYPE C-3

Produces visible image from infrared image in complete darkness. For seeing in dark with infrared light source. Nothing to wear out. Operates on two flashlight batteries. Has high-voltage power supply and image tube (1P25) and elaborate optical system. Weighs only 6¼ pounds and is only 11" long, 8" high, 5-12" wide. Manufactured by R. C. A. Objective f/0.9 Schmidt system, 2.4" focal length, 3¼" aperture. Excellent condition, with waterproof carrying case. Price **$35.00**

INFRARED FILTERS

5⅛" diam. x ⅛" thick. Price **$2.95**



SIMPLE DIFFERENTIAL WITH BALL-BEARING SUN GEARS

The 1:1 reverse ratio spur gears are 48-tooth, 32-pitch brass with 3/16" available face. On one side the shaft is 23/64" diam. for 11/16" and has a pinhole, then increases in diam. to 0.377" for the remaining 3/16" of length. On the other side, the shaft is 0.377" diam. ⅛" long. 2-13/16" diam. is required to clear the body. Stock No. A6-115. Price **$15.00** each.



SIMPLE DIFFERENTIAL WITH SPACED-OUT SUN GEARS

1:1 reverse ratio spur gears are aluminum, 3/32" face, 32-pitch, 32-tooth on one side, 48-tooth on the other. The body is ¾" thick, but the sun gears are spaced out so that they are 1½" apart. ¼"-diam. shaft on each side is 23/32" long. Over-all length, 3⅛". Requires 1-23/32" diam. to clear the body. Stock No. A6-124. Price **$4.50** each.

VARIABLE-SPEED BALL DISK INTEGRATORS (All shafts ball-bearing supported)

No. 145. Forward and reverse 2¼-0-2¼. Input shaft spline gear 12 teeth 9/32" diam., ⅜" long. Output shaft 15/64" diam. x 15/32" long. Control shaft 11/32" x ⅜" long. Cast-aluminum construction. Approx. size 3" x 3" x 2¾". Price $17.50

No. 146. Forward and reverse 4-0-4. Input shaft 5/16" diam. x ¾" long. Output shaft 15/64" diam. x 9/16" long. Control shaft 11/64" diam. x 11/16" long. Cast-aluminum construction. Approx. size 4½" x 4½" x 4". Price $18.50 each.

Send full amount with order. All prices net f.o.b. Pasadena, Calif. No C.O.D.'s, please.

C&H SALES CO.

2176 East Colorado St.
Pasadena 8, Calif. MUrray 1-7393

poor seeing of this type is present, there can be appreciable positional errors on photographs, since image points separated by only a few seconds of arc may be differently refracted.

Something can be done to combat the first two varieties of inferior seeing, but not the third. Seeing of type I generally results from faulty telescope design (for example, an uninsulated metal tube), heat from the observer's body, or air currents in or just above the tube. With a reflector, on very still nights, it is well to open the vents at the bottom of the tube for a while to get an approximation to thermal equilibrium, then close them to avoid slow convection currents in the interior of the tube. On windy nights, leave the vents open, as the gusts will prevent the formation of appreciable temperature gradients in the interior of the tube.

In any event, telescope tubes are best made of material with low heat conductivity and low heat capacity, to keep rapid local variations of outside air temperature from being transmitted to the tube interior, and to minimize the amount of heat so transferred. It has often been found that the performance of a telescope with a metal tube is improved by applying an interior layer of insulation, a favorite material being sheet cork.

An open or latticework tube may be helpful if it is rotatable, so that if the wind is behind the observer on a cool night, the tube can be turned to prevent his body heat from causing poor seeing of type I. However, the asymmetry of such a tube invites temperature gradients, as its material loses heat by radiation, becoming cooler than the surrounding air.

Poor seeing of type II can sometimes be cured by moving the telescope to a better location. Removal of the instrument from a downslope site will avoid the typical night drainage of turbulent cold air. Getting away from a heated building may help. Other remedies will occur to the reader who considers his own observing situation.

While nothing can be done about inferior type-III seeing, it often does not hinder delicate visual work. Photography is possible when these high-level disturbances are not too pronounced, if the exposure time is kept short compared to the period of rippling.

In summation, it is plain that seeing of type I is to a large extent remediable, that II may sometimes be ameliorated, and that III is frequently not serious enough to impair visual and photographic observation. Of course, it will often happen that the seeing, despite all efforts, is just not good, in which event it is as well to go to bed and get a good night's sleep!

FRANK VAUGHN
5801 Hammersley Rd.
Madison, Wis.

The only thing more amazing than its low price is its breathtaking performance!

See how this truly portable RV-6

6-inch DYNASCOPE®

can bring you new viewing pleasures!

Model RV-6
Complete with Dyn-O-Matic Electric Drive and All Features Pictured and Described on This Page

$194.95

f.o.b. Hartford, Conn.
Shipping Wt. 55 lbs.
Express Charges Collect

This new addition to the Dynascope line has won immediate recognition from schools, colleges, amateurs and professionals, as an outstanding achievement in a 6-inch telescope. Here is large aperture in a quality instrument which sells for a price that compares with many 4-inch telescopes.

You will find that this fine new portable 6-inch offers every feature necessary for serious astronomical viewing. The superb optical system resolves difficult objects with definition that is absolutely startling. The close tolerances of the precision construction assure an accuracy and smoothness of operation once associated only with the finest custom models. The heavy-duty mount, complete with electric drive, provides the stability so essential for satisfactory viewing, yet there is easy portability because in a matter of minutes the entire telescope can be dismantled into three easy-to-handle sections.

Only Criterion's engineering ingenuity, coupled with volume production and modern manufacturing methods, makes this handsome 6-inch model available at such reasonable cost. You can order it with complete confidence that it will live up to your expectations in every way, for this assurance is *guaranteed* under our full-refund warranty. Send your check or money order today. Or use our liberal time-payment plan and take months to pay.

You Could Pay $100 More Without Getting All These Superior Features (Except on Another Dynascope)

1. **EXQUISITE OPTICAL SYSTEM INCLUDES F/8 6-INCH PARABOLIC PYREX MIRROR** accurate to better than ⅛ wave, zircon-quartz coated. Teamed with elliptical diagonal, mounted in 4-vane adjustable spider.

2. **NEW DYN-O-MATIC ELECTRIC DRIVE** with smooth self-acting clutch that engages and disengages automatically as you seek different objects. Will not interfere when manual operation is desired. Entire drive housed in trim aluminum case. Plugs into ordinary household socket.

3. **CHOICE OF ANY 3 EYEPIECES** from the following: **65X**-18-mm. Achromatic Ramsden; **100X**-12.7-mm. Achromatic Ramsden; **150X**-9-mm. Achromatic Ramsden; **220X**-6-mm. Orthoscopic; **320X**-4-mm. Orthoscopic. All are precision-threaded, give sharp images to extreme edges.

4. **SOLID NEW EQUATORIAL MOUNT** extra-reinforced design to provide pillarlike stability. No annoying side play or wobble. Adjusts easily to any latitude.

5. **SETTING CIRCLES** for both right ascension and declination. Accurate to 0.0001 inch. Handsomely finished in fine aluminum.

6. **6 x 30 FINDERSCOPE, ACHROMATIC, COATED,** with accurate crosshairs and fine focus. Durable cast-aluminum bracket with 6 adjusting screws allows positive collimation.

7. **ROTATING RINGS FOR MAXIMUM VERSATILITY AND VIEWING COMFORT.** Solid-cast, chrome finish, rings are generous 1" wide with felt lining. Newly designed construction, with over-sized knurled adjusting knobs, affords maximum rigidity and allows for quick disassembly and portability, with or without tube.

8. **STRIKINGLY HANDSOME, WHITE, 50" BAKELITE TUBE** with porcelainized Duralite finish, durable yet light. Walls are ⅛" thick, completely insulated and anti-reflective blackened inside.

9. **BRASS RACK-AND-PINION EYEPIECE HOLDER** has precision-cut worm and gear for easier, smoother, more positive focusing. Takes standard 1¼" oculars, negative or positive.

10. **STRONG, VIBRATION-FREE, ALL-METAL TRIPOD** with easily removable legs. Provides sure, steady support, plus lightweight portability.

CRITERION MANUFACTURING CO.

331 Church St., Hartford 1, Conn.

Manufacturers of Quality Optical Instruments

ENJOY IT NOW — take up to 24 months to pay under our Easy Terms plan!

Just a small down payment delivers this big 6-inch DYNASCOPE. Pay the balance at your convenience over 6, 12, 18, or even 24 months! Write for details, or order today by sending your check or money order with coupon below.

Sound too good to be true? Then read what these delighted DYNASCOPE owners have to say:

"I never dreamed such a brilliant, clear image could be had with a telescope costing under $200. It is one of these rare bargains that you find only once or twice in a lifetime."
B.S. — New Jersey

"A friend of mine has a 6-inch telescope without electric drive, for which he paid over $300. He agrees that my new DYNASCOPE not only outperforms it, but is a much better-looking and better-built instrument." **J.L. — New York**

"Congratulations on the excellence of your workmanship. The optics are truly amazing. I never expected such wonderful performance."
L.H.N. — Mass.

Criterion Manufacturing Co.
Dept. STR-4, 331 Church St., Hartford 1, Conn.

☐ Please send me, under your MONEY-BACK guarantee, the RV-6, 6" Dynascope. Full payment of $194.95 is enclosed.

☐ Send complete information describing easy payment plans.

☐ Send FREE ILLUSTRATED LITERATURE of all the telescopes in the Dynascope line.

Name .....

Address .....

City ..... **State** .....

GOTO
TOKYO

Established in 1926, Japan's oldest and largest firm specializing in astronomical telescopes.

The Goto Planetarium Hall at the 1959 Tokyo International Trade Fair.



Goto's Model M-1 has been acclaimed by many visitors to the Fair, including specialists.

GOTO'S MEDIUM-SIZE PLANETARIUM

Goto Optical Mfg. Co., Japan's foremost maker of astronomical telescopes, is proud to present its revolutionary medium-size planetarium.

FEATURES:

1. Projects the entire celestial sphere and can show the heavens above any point in the world. Therefore, the starry sky over various places south of the equator, particularly the Antarctic, can be projected to the very last detail.
2. Projection of the apparent motions of the moon, sun, and five planets is provided. The peculiarities of the motion of each body can be shown.
3. Precession phenomena can be demonstrated.
4. Automatic explanations can be given with tape recordings.
5. Music and imitation sounds can be produced by a record player.
6. Remote operation from the control desk is possible.
7. By using auxiliary projection apparatus various types of celestial phenomena may be demonstrated.

Catalogues will be sent upon request.

GOTO OPTICAL MFG. CO.
1-115 Shimmachi, Setagaya-ku, Tokyo, Japan

Cable Address:
GOTOPTIC TOKYO

ALDEBARAN OCCULTATION

On the night of September 22-23, the 1st-magnitude star Alpha Tauri was covered by the moon. The phenomenon was watched at the Robert T. Longway Planetarium, Flint, Michigan, by R. N. Watts, Jr., and D. Splane. They used 7 x 50 binoculars and, with the aid of a sweep-second watch they had compared with time signals, found that the star disappeared at 11:16:26.0 p.m. Eastern standard time, and reappeared at 0:19:37.5. High cirrus clouds may have affected these results.

At New Orleans, Louisiana, W. P. Searcy, III, obtained photographs of Aldebaran soon after it had emerged from behind the dark edge of the moon. His pictures were taken with a 4-inch reflector.

OCCULTATION OF NEPTUNE

The sky was very clear when I observed the occultation of Neptune by the moon on September 6th, using my 5-inch f/15 Mellish refractor. With a 1¼-inch Erfle eyepiece, the bright part of the moon was kept out of the field, but the dark side was clearly seen.

First contact occurred at 7:12 p.m. Eastern standard time. Neptune lost brightness during a little more than one second, then disappeared suddenly. The fading may have taken longer, but with the low power the planet showed only a small disk. I am anxious to hear about other observations of this event.

RENE DOUCET
650 Des Grandes Prairies Blvd.
Cap de la Madeleine
Quebec, Canada

SUNSPOT NUMBERS

The following American sunspot numbers for July and August have been derived by Dr. Sarah J. Hill, Whitin Observatory, Wellesley College, from AAVSO Solar Division observations.

July 1, 163; **2**, 121; **3**, 145; **4**, 118; **5**, 121; **6**, 113; **7**, 113; **8**, 102; **9**, 100; **10**, 84; **11**, 102; **12**, 123; **13**, 152; **14**, 152; **15**, 150; **16**, 197; **17**, 209; **18**, 178; **19**, 154; **20**, 143; **21**, 109; **22**, 104; **23**, 106; **24**, 117; **25**, 133; **26**, 156; **27**, 176; **28**, 178; **29**, 184; **30**, 197; **31**, 191. Mean for July, 141.6.

August 1, 183; **2**, 214; **3**, 204; **4**, 179; **5**, 194; **6**, 167; **7**, 173; **8**, 141; **9**, 147; **10**, 161; **11**, 144; **12**, 136; **13**, 106; **14**, 106; **15**, 134; **16**, 145; **17**, 122; **18**, 141; **19**, 162; **20**, 160; **21**, 164; **22**, 195; **23**, 192; **24**, 153; **25**, 193; **26**, 228; **27**, 253; **28**, 259; **29**, 273; **30**, 285; **31**, 261. Mean for August, 179.8.

Below are provisional mean relative sunspot numbers for September by Dr. M. Waldmeier, director of Zurich Observatory, from observations there and at its stations in Locarno and Arosa.

September 1, 290; **2**, 256; **3**, 202; **4**, 161; **5**, 148; **6**, 144; **7**, 135; **8**, 136; **9**, 157; **10**, 141; **11**, 155; **12**, 170; **13**, 148; **14**, 151; **15**, 168; **16**, 130; **17**, 87; **18**, 100; **19**, 120; **20**, 143; **21**, 132; **22**, 155; **23**, 136; **24**, 155; **25**, 105; **26**, 106; **27**, 92; **28**, 87; **29**, 80; **30**, 76. Mean for September, 142.2.

Finest Christmas Gift Ever . . .

ASTROLA Reflecting Telescopes

In 1953, we at Cave Optical Co. conceived the idea of building reflecting telescopes of really adequate size and optical quality seldom found before in commercial reflecting telescopes. Fully realizing the amateur astronomers' telescopic requirements and limited budgets, we priced our ASTROLA Reflectors within the reach of all.

Since the first introduction of ASTROLA Reflecting Telescopes, increasing demand has remained ahead of our expanding productive capacity. We have always placed performance quality ahead of all other considerations in manufacturing the ASTROLA. Although appearance may have nothing to do with a telescope's performance, ASTROLA Reflectors are certainly pleasing to look at, but they are even more delightful to look through.

Thus, more than six years ago we pioneered the larger, but sensibly priced Newtonian reflecting telescope field. Success brings competition and imitation from others. Therefore, it is well to remember that ASTROLA was first and is still far ahead in quality and performance. In recent months we have greatly expanded our facilities for producing not only more ASTROLA Reflectors but much larger instruments. We are now doing all our own instrument machine work, so that we may give much more rapid delivery on each and every ASTROLA, but we also have even closer quality control.

Here are a few reasons why ASTROLA sales remain ahead of production:

1. ASTROLAS are by no means mass-produced, each instrument being carefully hand-assembled and each telescope completely field-tested before leaving our shops.

2. Each complete ASTROLA has been painstakingly engineered and meets every requirement of discriminating telescopists.

3. Every complete ASTROLA and every telescope mirror that we make or refigure are guaranteed unconditionally to give identical inside- and outside-of-focus star images, your assurance of ultimate optical quality.

All-New De Luxe Model "B" 8-inch ASTROLA, f/7, with rotating tube, clock drive, setting circles, and 3 oculars (85x, 210x, 360x, or 3 of your choice). **$590.00** plus $12.50 packing and crating charges.

The Completely New De Luxe Model "C" 10-inch ASTROLA, f/7, complete with rotating tube, clock drive, setting circles, and 4 oculars (74x, 111x, 222x, 444x, or 4 of your choice). **$750.00** plus $15.00 packing and crating charges.

	STANDARD	DE LUXE
STUDENT ASTROLA, 6-inch, f/7 or f/9	$200.00	—
ASTROLA MODEL "B", 8-inch, f/6, f/7, or f/8	$390.00	$590.00
ASTROLA MODEL "C", 10-inch, f/6, f/7, or f/8	$495.00	$750.00

NEW PYREX TELESCOPE MIRRORS AND REFIGURING

If your present telescope is not giving truly outstanding performance, let us refigure your mirror to our standard quality, and furnish you an elliptical diagonal plus aluminizing, return postage prepaid.

	NEW PYREX MIRRORS f/5 through f/10	REFIGURING f/5 through f/10
6-inch	$ 60.00	$ 35.00
8-inch	$ 92.50	$ 50.00
10-inch	$160.00	$ 75.00
12½-inch	$250.00	$120.00

Immediate delivery on new 6-, 8-, and 10-inch mirrors of normal focal ratio. Send for our newest 1960 full-color catalogue.

When in the Los Angeles-Long Beach area, visit our display room. Always in stock are many small reflectors and refractors, plus a large assortment of astronomical literature. Generous allowances on trade-ins.

CAVE OPTICAL COMPANY

4137 E. Anaheim St., Long Beach 4, Calif.
Phone: GEneva 4-2613

SINCE 1933

PANCRO mirrors, inc.

Reg. U.S. Pat. Off.

Let us give you the same hard aluminum coating that we are applying to the Cave Astrola mirrors. If your glass is properly polished, our coating will give a **minimum of 90% reflection.** Will not blister or peel. Check or money order, including return postage and insurance (Calif. res. add 4% sales tax), will guarantee **8 hours shipping service.** No C.O.D.'s please. Prices f.o.b. L.A.

6-inch.....$6.50 10-inch.....$11.25
8-inch.....$8.50 12-inch.....$16.00

PANCRO MIRRORS, INC.
Research and Production Laboratories
2958 Los Feliz Blvd., Los Angeles 39, Calif.

Do-It-Yourself
TELESCOPE KITS

Now, at last! All the mechanical parts you need to mount your finished optics. These matched items will save you money and guarantee you a professional-appearing instrument.

★ Finest Parks fiberglass tube
★ Completely adjustable 4-vane diagonal holder
★ 8 x 50 achromatic finder
★ Helical focuser
★ Cast-aluminum adjustable mirror cell

Prices: 6", **$57.50**; 8", **$71.50**; 10", **$109.50**; 12", **$129.95**

f.o.b. Canoga Park, California
Equatorial mounts also available.

ANRA MANUFACTURING ENGINEERS
7431 Canoga Ave., Canoga Park, Calif.

Deep-Sky Observers

An extremely fine, imported Zuiho binocular with a wide field of view — 10½°!

$35.00
Tax Paid

Complete with beautiful case. Delivered anywhere in the Continental United States.

AMERICAN TYPE WIDE ANGLE
7 x 35 Central Focus

SPECIFICATIONS

Magnification, 7x. Objective aperture, 35 mm. Eyepiece aperture, 24 mm. Exit pupil, 5 mm. Field of view at 1,000 yds., 551 ft. Field-of-view angle, 10.5°. Height, 128 mm. Breadth, 195 mm. Net weight, 950 grams. Weight of pigskin case, 400 grams.

We have sold hundreds of pairs of these binoculars with the utmost satisfaction. They are unconditionally guaranteed. If not pleased, return them and your money will be immediately refunded.

Be sure to write for our list of used telescopes.

THIS MONTH'S SPECIAL
One Unitron Model 155—$495
Trades accepted.

Before you buy or trade a telescope or radio amateur gear, see WARD W2FEU for the best deal.

Established 1936

ADIRONDACK RADIO SUPPLY
P. O. Box 88, Amsterdam, N. Y.
Ward J. Hinkle, W2FEU — Owner

DEEP-SKY WONDERS

AROUND the famous variable star Delta Cephei draw a circle 5½ degrees in radius. Then tally up the nebulae and clusters within the area. If you are using Norton's *Star Atlas* you will find five objects, while the de luxe edition of the Skalnate Pleso *Atlas of the Heavens* will give you six. But only one of these is common to both maps, though all but one have NGC numbers; it is evidently worth while to use more than one set of observing charts.



This chart, 11 degrees in diameter and with north at the top, is based on the Skalnate Pleso "Atlas of the Heavens," but has been simplified to emphasize the location of the clusters and nebulae discussed in the text. The objects plotted in Norton's "Star Atlas" have been added in their proper positions according to the "New General Catalogue." The symbols are the same as used on the Skalnate Pleso charts.

NGC 7380 (77^h) is the object found on both maps, at right ascension 22^h 44^m.9, declination +57° 49′ (1950 coordinates). It is a galactic cluster of some 50 stars, 10′ in diameter, and fairly faint. While it is not mentioned in observing handbooks, it can be seen in a 4-inch telescope, though the region is difficult because it lies in the Milky Way.

The rest of the clusters plotted in Norton's are: NGC 7261, 22^h 18^m.6, +57° 50′; 7281, 22^h 23^m.0, +57° 35′; 7352, 22^h 37^m.8, +57° 08′; and 7438, 22^h 55^m.4, +54° 05′. All of them were first described by John Herschel, and the *New General Catalogue* calls them pretty large or very large. Most are now generally regarded as merely brighter portions of the Milky Way. The writer, using a 10-inch reflector, found it very confusing trying to distinguish them among the starclouds in this region.

The Skalnate Pleso shows five other objects, two of them clusters: NGC 7245, 22^h 13^m.6, +54° 05′, a weak grouping 3′ in diameter and containing 40 stars; and NGC 7419, 22^h 52^m.6, +60° 34′, containing but 25 stars in a 2′ diameter.

There are three planetaries in the area, the brightest being IC 1470, at 23^h 03^m.2, +59° 59′, an 8th-magnitude object measuring 70″ by 45″, with a 12th-magnitude central star. This planetary is plotted as a very small diffuse nebulosity in the field edition of the *Atlas of the Heavens*.

NGC 7354, at 22^h 38^m.4, +61° 01′, is of the 13th magnitude and 32″ in diameter, apparently of such low surface brightness that it is virtually impossible to find visually with amateur telescopes. I failed to observe it in three nights of searching. The unnumbered third planetary is even more difficult, being only of magnitude 14, and 8″ in diameter. It is located at 22^h 54^m.4, +56° 54′, plotted on the accompanying chart.

WALTER SCOTT HOUSTON
Rte. 3, Manhattan, Kans.

Announcing the NEW . . .

MONTHLY EVENING SKY MAP

"The stars brought down to earth"

—detailed star and planet charts for each month of the year.
—informative articles for the beginner in astronomy.
—advance information on current events in the skies.

$3.00 per year Sample copy, 25¢

Published bimonthly by:

SKY MAP PUBLICATIONS, INC.
Box 213-S, St. Louis 5, Mo.

Astronomy Films
16-mm. sound, 400-foot reels
I THE SUN; II THE MOON;
III SOLAR SYSTEM; IV MILKY WAY;
V EXTERIOR GALAXIES.

2 x 2 SLIDES

35-mm. STRIPS OF SLIDES

THROUGH 200-INCH AND OTHER GREAT TELESCOPES
Catalogues on request.

International Screen Organization
1445 18th Ave. North, St. Petersburg 4, Fla.

Lafayette

14" TRANSPARENT CELESTIAL GLOBE

★ Basic Aid for Students and Teachers of Astronomy ★ Navigation ★ Aeronautics ★ Astronautics

59.50

only 5.95 down

FIRST TIME UNDER $100.00

AVAILABLE ONLY THROUGH LAFAYETTE IN THE UNITED STATES, CANADA, AND MEXICO

★ Adjustable satellite orbits

★ Satellites travel around earth automatically

★ Adjustable sun and moon

★ Adjustable lunar orbital inclination

★ 80 constellations — including all stars of 1st, 2nd, 3rd, and 4th magnitudes

★ Similar devices used by the Navy — Air Force — Marine Corps

A true "mechanical universe." Designed for and dedicated to those who recognize the challenge and adventure that lie on the frontiers of space. The concept of the celestiel sphere is the most convenient means of illustrating and discussing the common motions of the stars and the individual motions of the members of the solar system. It offers an easily used and easily understood working model of our universe. An invaluable aid in visualizing and identifying the stars and constellations and their relationship to each other, to terrestrial positions, and to times and dates. Permits an understanding of explorations in time and space beyond the earth's atmosphere.

The celestial sphere may be defined as a sphere of infinite radius with its center located at the center of the earth. The system of geocentric co-ordinates will illustrate the true geographic zeniths of the sun and stars at any time on any dey. All systems of astronomical spherical co-ordinates, such as the equatorial, galactic, ecliptic, which are based on the celestial sphere, rotate with the sphere. The apparent contra-rotation of the stars, actually due to the earth rotating about an axis, is clearly demonstrated in this well-made device.

The sphere shows how the sky will appear at any hour — on any date — from any point on earth. With it you can tell time by the stars — learn to tell time by the sky — clearly illustrate and help solve navigational problems. Traces the orbits of man-made satellites and finds their overhead positions. Serves as a superb star chart. Sun and moon positioned by external controls. Artificial satellites automatically circle around the terrestrial globe. The various stars, constellations and their identifications are permanently molded — not printed — on the celestial sphere.

The celestial globe is 14" in diameter, with the terrestrial globe centered within. Equipped with horizon ring — time ring — fixed meridian ring — swinging meridian ring — sun and moon pointers — fixed ecliptic — detachable "planet" indications — universal base — and complete instructions.

F-400 ..... Shpg. Wt. 36 lbs. ..... Net **59.50**

Lafayette Radio CORPORATION

165-08 LIBERTY AVE.
JAMAICA 33, N.Y.

USE OUR EASY PAYMENT PLAN
10% DOWN — EASY MONTHLY PAYMENTS

NEW YORK, N.Y., 100 Sixth Ave.
BRONX, N.Y., 542 E. Fordham Rd.
NEWARK, N.J., 24 Central Ave.
PLAINFIELD, N.J., 139 West 2nd St.
BOSTON, MASS., 110 Federal St.

LAFAYETTE RADIO P. O. BOX 1000 JAMAICA 31, NEW YORK

☐ Please Send F-400 CELESTIAL GLOBE.
I Enclose $..... Dept.
☐ Rush Free Catalog. SKK9

Name .....

Address .....

City ..... Zone ..... State .....

Star Atlases and Books on Astronomy

New: **OUR SUN**, by D. Menzel, 2nd edition.....$6.50
New: **ASTRONOMICAL PHOTOGRAPHY WITH THE TELESCOPE**, by T. Rackham.....$7.50
AMATEUR ASTRONOMER'S HANDBOOK, by J. B. Sidgwick....$12.50
OBSERVATIONAL ASTRONOMY FOR AMATEURS, by J. B. Sidgwick.....$10.00
THE OBSERVERS GUIDE.....$2.95
THE MOON, by Wilkins and Moore..$12.00
GUIDE TO MARS, by P. Moore.....$2.75
THE PLANET JUPITER, by B. Peek..$8.50
AMATEUR TELESCOPE MAKING Book 1, $5.00; Book 2, $6.00; Book 3, $7.00
Webb's ATLAS OF THE STARS.....$6.50
BONNER DURCHMUSTERUNG.....$100.00
1960 AMERICAN EPHEMERIS AND NAUTICAL ALMANAC—order now $4.00

Write for new free list on astronomical literature. Books on telescope making and optical glass-working. All books of any publisher — mail your wants. Out-of-print books located in a special search service.

HERBERT A. LUFT
69-11 229th St., Oakland Gardens 64, N. Y.

Again in Print!

Shapley-Ames CATALOGUE OF BRIGHT GALAXIES

"A Survey of the External Galaxies Brighter than the 13th Magnitude" gives NGC number, 1950 position, magnitude, size, and type for 1,249 objects.

This publication (Vol. 88, No. 2, of the Harvard *Annals*) was out of print for some time, but is now available in an offset-printed edition.
$1.50 postpaid

Order directly from

HARVARD COLLEGE OBSERVATORY
Cambridge 38, Massachusetts

CHRISTMAS SUGGESTIONS

Our past ads have listed the diversity of our astronomical products, all of which will make excellent gifts. It's really not too early to think of Christmas. You can't miss with an astronomical present.

SOLARSCOPE
Many of your friends will want one. Compact, yet loaded with 1,000 solar system facts and figures, at a twist of the disk, for every planet.
Only $2.25

THE SUN
By Giorgio Abetti. Written by a lifelong student of the sun. A valuable addition to any library. **$12.00**

THE SUN AND ITS INFLUENCE
By M. A. Ellison. **$4.50**

FABULOUS FIREBALL
By D. S. Halacy, Jr. A new approach on harnessing solar energy. A very different revelation. **$3.00**

ONCE AROUND THE SUN
By Ronald Fraser. The story of the International Geophysical Year. **$3.95**

SATELLITES AND SPACEFLIGHT
By Eric Burgess. **$3.95**

THE MILKY WAY
By Bart J. and Priscilla F. Bok. Not really a new book, but still one of the best illustrated ones available. **$5.50**

DISCOVERY OF THE UNIVERSE
By Gerard de Vaucouleurs. **$6.00**

THE GREEN FLASH
By D. J. K. O'Connell, S. J. Beautiful colored pictures and descriptions of this rarely seen phenomenon. **$6.00**

We haven't the room to list all our books. Our catalogue not only shows more but many other items to please that astronomically minded friend or relative. Write for your copy today.

ASTRONOMY CHARTED
33 Winfield St., Worcester 10, Mass., U.S.A.
Phone: PL 5-6992

BOOKS AND THE SKY

HANDBUCH DER PHYSIK, VOL. 53
ASTROPHYSICS IV: STELLAR SYSTEMS

S. Flügge, editor. Springer-Verlag, Berlin, 1959. 565 pages. DM 142.

ONE of the most overworked statements in the repertoire of a book reviewer is: "This book fills a long-felt need." It is, however, especially applicable to this attractive and stimulating volume on galactic structure and dynamics, galactic and globular clusters, galaxies and cosmology. So much has happened and is presently developing in these areas of research that three of the outstanding chapters in this book are on subjects that were scarcely known about a dozen years ago. One may hope that another *Handbuch* on these subjects can be written 10 years hence, for 25 years would be too long to wait. It is that time since the *Handbuch der Astrophysik* appeared.

Volume 53 should be an incentive to textbook authors to expand their offerings on galactic structure and the external galaxies. If there is any criticism to be voiced, it is that the names of H. Shapley, W. Baade, N. U. Mayall, and A. Sandage are not found in the list of authors. The first 12 chapters are in English; the last two, on cosmological theories, are in German.

F. K. Edmondson's chapter on the "Kinematical Basis of Galactic Dynamics" emphasizes that the basic kinematical principles, as stated by E. A. Milne, have been largely overlooked, with consequent troubles. The Oort double sine wave (second harmonic) in radial velocities is well known, but there is an often neglected first-harmonic term, increasing as the square of the distance from the sun, which in a typical case has about one-fourth the amplitude of the Oort term. Edmondson gives rather convincing reasons why the "low" Vyssotsky-Janssen "basic solar motion" of 15.5 kilometers per second should be used. A consistent set of galactic rotation parameters has not yet been derived. To obtain them, accurate luminosities and absorptions for distant supergiants are especially needed. These can almost certainly be obtained eventually with appropriate narrow- and broad-band photoelectric observations of such stars within and outside of selected galactic clusters.

B. Lindblad's contribution on "Galactic Dynamics" is a useful discussion of stellar mass motions, velocity distributions, and dispersions of stellar velocities with time. The final sections are concerned with spiral-structure problems and evolutionary effects. Much of the account is mathematical and not for the casual reader. The author refers to the importance of determining whether the spiral arms "trail" or "open up" and states: "From the observational point of view this question has not yet been definitely settled." The best that can be said for this statement is that it is true. Any interested reader can form his own judgment from the published observational material; the pertinent references may be found in the second de Vaucouleurs chapter in this book.

The study of galactic structure has been enormously accelerated by the development of radio astronomy. Leiden Observatory, under the leadership of J. H. Oort, is in the forefront of the spectacular 21-centimeter investigations of the dynamics and spiral structure of our galaxy. His chapter on radio-frequency studies of galactic structure is very readable but all too short, with his own contributions somewhat underemphasized. The spiral structure, derived from 21-cm. observations, is depicted in Fig. 4a (which is already out of date!), and the results from the continuous radiation observed at 1.5 meters for the whole sky are shown in Fig. 14.

The study of star clusters has always been peculiarly profitable. Because of this, the moderately long chapter by Helen S. Hogg has been written under difficulties — the literature is very extensive and many important topics need to be covered. The text is essentially nonmathematical and is delightful and informative reading for professional and amateur astronomers alike.

A comparison with Shapley's 1933 *Handbuch der Astrophysik* chapter on the same subject reveals how much has been learned in 25 years. The last decade has been especially fruitful, with the development and application of new photoelectric techniques in the Southwest and at Pacific Coast observatories. Mrs. Hogg gives only a short discussion of stellar associations, a subject that might well take a chapter of its own.

Very useful catalogues of 514 galactic clusters and 118 globular clusters are given in two appendices. The care with which such listings must be compiled is illustrated by the position of NGC 7790, correctly given here, but erroneously noted in the earlier literature. The new position "moves" this cluster to an area of the sky (it was there all the time!) that includes three classical Cepheids and one eclipsing binary, with a number of consequent important cosmological implications.

The next two sections are "Discrete Sources of Cosmic Radio Waves" by R. Hanbury Brown and "Radio Frequency Radiation from External Galaxies" by B. Y. Mills. Radio astronomy has always suffered from poor resolving power, which makes "radio star" positions inexact, identifications very difficult, and angular size usually impossible to measure. This condition is accentuated because many radio telescopes have a complex pattern

of side lobes. Furthermore, the radio energy drops rapidly toward shorter wave lengths where the resolving power is higher. There is marked disagreement, for example, between the Cambridge University and the Australian surveys over a common area of the sky, not only regarding positions of discrete sources but their numbers as one goes to smaller and smaller flux densities.

Of the relatively few discrete sources that have been optically identified and investigated, some have yielded results and implications so exciting as to earmark these sources as an extraordinarily fertile subject for research. These two chapters together, with some overlap, give an excellent account of the techniques, problems, often contradictory results, and future hopes in this rapidly changing radio work, which depends so much on enormous instruments and the most sensitive detectors.

The next two chapters, by G. de Vaucouleurs, concern the galaxies, their classification, morphology, and general physical properties. This is a field in which de Vaucouleurs himself has made many fundamental contributions, despite the handicap of the comparatively limited instrumentation available to him in the Southern Hemisphere. The pages are liberally sprinkled with beautiful reproductions from the soon-to-appear Hubble memorial volume. The main revisions in Hubble's classification system are given, as well as de Vaucouleurs' own three-dimensional plan, the additional third "dimension" being ring or spiral structure.

Two short contributions by F. Zwicky on "Multiple Galaxies" and "Clusters of Galaxies" are written in the vigorous style associated with this author. I could find no reference by Zwicky to the important and extensive investigations of George Abell on clusters of galaxies. Zwicky's work on multiple galaxies leads him into considerable disagreement with other astronomers concerning the effects of galactic collisions. A 200-inch photograph and a diagram of a cluster of galaxies are given in which the density approaches the somewhat fantastic value of 250,000 galaxies per square degree. Zwicky refers to galaxy red shifts as symbolic velocities of recession, calls attention to the mere handful of observers having access to large telescopes (with consequent slow accumulation of data and difficulty in obtaining independent observational checks), and correctly warns against drawing any too precipitous conclusions from often necessarily scanty data.

The next chapter, on the large-scale organization of the distribution of galaxies, by J. Neyman and Elizabeth L. Scott, is presumably concerned with some of the same problems as those in Zwicky's chapters, but the approach is very different. These authors' statistical analyses have been stimulated by the Shane-Wirtanen survey with the Lick 20-inch astrograph, which covers the sky from the north pole to declination −23°. Many of these six-by-six-degree plates have thousands upon thousands of extragalactic images on them; most of the counts have been made and already some of the analyses have been published.

G. C. McVittie's contribution concerns the observational basis of cosmology, and is refreshing because we find here a theorist who is properly skeptical of the observational data. The chapter is well written and deserves careful study. It was prepared three years ago, and since then many photoelectric investigations of clusters, Cepheids and the Magellanic Clouds have strengthened the observational picture. Continued photoelectric researches in the Magellanic Clouds are especially needed, but the number of photoelectric observers who are working at 19th and 20th magnitude in these uniquely important galaxies is very small.

The last two chapters concern various theoretical cosmologies. Written by O. Heckmann and E. Schücking, they require a knowledge of both German and tensor analysis, and will therefore be rather heavy going for the average astronomer trained in this country.

JOHN B. IRWIN
Goethe Link Observatory
Indiana University

Availability of ESSCO PUBLICATIONS

For several decades, the Eastern Science Supply Company, Boston, Massachusetts, provided teachers of astronomy, both amateur and professional, with publications and other study materials. We have acquired the remaining stock of star maps, charts, and booklets, of which the items listed below will be continued in print and brought up to date where necessary:

SC1	Equatorial constellation chart — with star designations
SC1	**Test** equatorial chart — **without** star or constellation names
SC2	Circumpolar constellation chart — with star designations
SC2	**Test** circumpolar chart — **without** star or constellation names
S508A	Ecliptic-based star map — with equatorial grid and names
S508	Ecliptic-based star map — with equatorial grid, **without** names
S508B	Ecliptic star map list — positions and magnitudes for 224 stars
S505	Nine-inch protractor on paper — for planet orbit drawings
S511	Inner planet chart — orbits of Mercury, Venus, Earth, Mars
S512	Outer planet chart — orbits of Mercury to Saturn
S501A	Special rectangular co-ordinate paper — for star maps
S502	Polar co-ordinate paper — for circumpolar star maps

Price for each item listed above: 1 to 9 sheets, 10 cents each; 10 to 24 sheets, 8 cents each; 25 to 99 sheets, 6 cents each; 100 or more, 5 cents each.

From Stetson's *Manual of Laboratory Astronomy*, the following chapter is available as a separate booklet, at 50 cents each: I, Star Chart Construction.

SPECIAL: DC5 — Large wall chart of the Draper spectral classes, printed in four colors and mounted on canvas. Limited quantity. (Slightly damaged along edges.) **$14.00 each**

SKY PUBLISHING CORPORATION
Cambridge 38, Massachusetts

Complete practical guide to telescope making—easy to use—scientifically accurate

For the first time, the amateur astronomer with no specialized knowledge of the mathematical theories of telescope making can produce a perfect mirror and a first-class telescope — simply by following the detailed step-by-step instructions and illustrations in this book. The author begins by explaining why the telescope he is going to build *with you* is the best kind for most amateurs, tells how to set up an efficient but inexpensive workshop, and proceeds from there to help you complete the job.

CLEAR TEXT, DETAILED DRAWINGS, SPECIAL PHOTOGRAPHS TEACH YOU HOW TO:

- Choose the right telescope to build
- Buy inexpensive equipment and tools
- Rough and fine grind the mirror
- Select the correct eyepiece
- Construct the tube and its fittings
- Mount your telescope
- Plan and build a complete observatory
- Photograph the stars

In addition, an invaluable appendix includes a glossary of terms, a list of suppliers of materials, tables for finding sidereal time, tables of films and exposures for celestial photography. $5.95

STANDARD HANDBOOK FOR TELESCOPE MAKING

by N. E. HOWARD, author of "Handbook for Observing the Satellites"

•

THOMAS Y. CROWELL COMPANY

Est. 1834 **New York 16, N. Y.**

Sky PUBLICATIONS *for every astronomical interest*

Norton's STAR ATLAS

This famous star atlas and reference handbook is particularly suited for amateurs and students who desire sky charts in book form. It covers the whole heavens, showing over 9,000 stars to magnitude 6½, nebulae, and clusters. There are descriptive lists of 500 interesting objects for viewing with small telescopes, and useful data are given for observers of the sun, moon, and planets. New 14th edition, 1959. **$5.25**

POPULAR STAR ATLAS

This compact, well-bound set of 16 maps is a simpler version of the Norton's *Star Atlas* described above. All stars down to magnitude 5½ are included. This edition is excellent for field use by constellation study groups and by meteor parties. **$2.00**

Elger's MAP OF THE MOON

The chart of the moon itself is 18 inches high by 17½ inches wide and identifies all the important lunar features. Below the map are notes by H. P. Wilkins on 146 of the more interesting areas. Printed on a sheet of paper 30 by 19½ inches and mounted on heavy canvas, the map is ideal for framing. **$3.00**

COLOR MAP OF THE NORTHERN HEAVENS

This is a large wall chart, 30 by 34½ inches, colorful as well as informative. The northern sky to −45° is shown on a polar projection, and each star is colored according to its spectral class. Stars brighter than magnitude 5.1 are included. The map is mailed unfolded in a heavy tube. **$2.00**

COLOR CHARTS OF THE MOON

Two striking color charts of the first- and last-quarter moon were drawn by Joseph Klepesta, the well-known Czechoslovakian amateur astronomer. Each lunar half is an impressive 27 inches in diameter; the overall size of a chart, 23 inches wide by 33 inches high. Printed in predominantly gold and black on heavy paper, the charts are shipped unfolded in a cardboard tube. **$3.00 per set of 2**

MOON SETS

18 pictures, showing the entire visible face of the moon, are made from unsurpassed Lick Observatory negatives of the first and last quarters. Each halftone print is 8½ by 11¾ inches. Key charts supplied. **$3.00 per set**

LUNAR CRESCENT SETS

10 pictures are a matching series to Moon Sets, but for the waxing crescent 4½ days after new moon, and the waning crescent about five days before new moon. Four prints are close-ups of the waxing crescent, four of the waning; two show each crescent as a whole. Key charts included. **$2.50 per set**

SKY SETS I

24 pictures of objects in the solar system and in the Milky Way, all celestial wonders of interest and beauty. Each halftone print is 8½ by 11¾ inches. Separate sheet of captions included. **$4.00 per set**

SKY SETS II

24 pictures of nebulae in our galaxy, portraits of other galaxies, many made with the 200-inch telescope, and four drawings of the 200-inch telescope by Russell W. Porter. Sheet of captions included. **$4.00 per set**

LUNAR MAP. In two colors and over 10 inches in diameter. **25 cents each; 3 or more, 20 cents each**

SPLENDORS OF THE SKY. 36-page picture book of our neighbors, near and distant, in the universe. **75c**

RELATIVITY AND ITS ASTRONOMICAL IMPLICATIONS, by Dr. Philipp Frank. **75c**

HOW TO BUILD A QUARTZ MONOCHROMATOR for Observing Prominences on the Sun, by Richard B. Dunn. **50c**

THE STORY OF COSMIC RAYS, by Dr. W. F. G. Swann, Bartol Research Foundation. **75c**

Special Christmas Suggestions

For Lunar Enthusiasts

Here is a special moon package of five items that will make a wonderful gift for anyone (including yourself) on your Christmas shopping list. **Moon Sets** and **Lunar Crescent Sets** show breathtaking views of the visible side of the moon. Our popular **Lunar Map** is ideal for reference at the telescope. **Elger's Map of the Moon** identifies all the important lunar features, while the **Color Charts of the Moon** are two striking maps to decorate your clubroom, den, or observatory wall. (Offer expires January 31, 1960.)

Moon Sets, Lunar Crescent Sets, Lunar Map
Elger's Map of the Moon, Color Charts of the Moon
A retail value of $11.25 for **$10.00**

For Armchair Scientists

With both amateur and professional scientists, **Sky Sets I** and **Sky Sets II** have long been favorites. Each set contains 24 large pictures of astronomical wonders, including objects in the solar system and in the Milky Way, portraits of other galaxies, and drawings of the Hale reflector by Russell W. Porter. (Offer expires January 31, 1960.)

Sky Sets I and **Sky Sets II**
A retail value of $8.00 for **$7.00**

And don't forget to give

SKY AND TELESCOPE

Here's a fine gift for all, from the budding young astronomer to the armchair scientist — a subscription to **Sky and Telescope,** the world's largest astronomical magazine, now printing 32,500 copies monthly. He will remember your thoughtfulness each month as he reads about our rapidly expanding science. A gift card, informing the recipient, will be sent in time for Christmas on orders received by December 10th. (Another way to give this magazine is through membership in an astronomical society that takes the group rate plan; consult your local club.)

Subscription in the United States and possessions: $5.00, one year; $9.00, two years; $13.00, three years. In Canada and Pan American Postal Union countries, $6.00, one year; $11.00, two years; $16.00, three years. In all other countries: $7.00, one year; $13.00, two years; $19.00, three years.

Two Editions Available

ATLAS OF THE HEAVENS

ATLAS COELI 1950.0

These 16 charts by Antonin Becvar and his coworkers at the Skalnate Pleso Observatory, Czechoslovakia, cover the entire sky to stellar magnitude 7.75, showing double, multiple, and variable stars; novae, clusters, globulars, and planetaries; bright and dark nebulae; the Milky Way and constellation boundaries.

FIELD EDITION. Each chart is reduced from the original Atlas Coeli and is printed on heavy, stiff paper 18 by 12¼ inches. Stars are white on black background. Charts are shipped flat and unbound, 16 to a set. Ideal for outdoor observing and use at the telescope.
$4.00 each; 2 for $7.50

DE LUXE EDITION. Printed in many colors, more than 35,000 celestial objects, including over 100 radio sources, are plotted. Bayer-letter and Flamsteed-number designations given. Transparent co-ordinate grid overlay included. The 16 charts are permanently bound in a heavy cloth cover, 16½ by 23 inches, with color chart key on a foldout flap. **$9.75**

ATLAS ECLIPTICALIS 1950.0

By Antonin Becvar

A bound set of 32 star charts, each 17½ by 24 inches, covers declinations between +30° and −30°, complete to magnitude 9.0, scale two centimeters per degree, epoch 1950.0. Spectral classes are shown by color. See page 161 of the January, 1959, issue for further description. Place your order with us and allow time for its transmission to the publisher in Czechoslovakia, from whom the atlas will be shipped directly to you. **$17.00**

MAKING YOUR OWN TELESCOPE

By Allyn J. Thompson. Tens of thousands of amateurs are using this basic book on telescope making. Here are complete step-by-step directions for making and mounting your own 6-inch reflecting telescope at low cost. This telescope can use magnifications up to 300 times. In easy-to-understand chapters, you will learn how to grind, polish, and figure the mirror, and how to make an equatorial mount that will provide a sturdy, solid support for your mirror.

211 pages, 104 illustrations (6th printing).... **$4.00**

THE HISTORY OF THE TELESCOPE

By Henry C. King. The first book to tell the full story of the evolution of the telescope. 456 pages, 196 illustrations, extensive references. **$7.50**

Spaceflight

Here is a popular yet authoritative magazine on rockets, astronautics, and space-travel astronomy, written especially for the layman, and edited by members of the British Interplanetary Society.

Spaceflight is printed during January, April, July, and October. Please specify the issue with which your subscription should start. Back issues to Vol. 1, No. 1, October, 1956, are available.

Subscription in United States and possessions, Canada, Mexico, and Central and South America: $2.50, four issues; $4.50, eight issues; $6.00, twelve issues. Single copy, 75 cents.

Write for descriptive catalogue. All items sent postpaid. Please enclose your check or money order.

SKY PUBLISHING CORPORATION

Harvard Observatory — Cambridge 38, Massachusetts

Films and the Sky

THE STORY OF THE UNIVERSE UNIT I: THE EARTH AND ITS MOONS

Harlan J. Smith. Films for Education, New Haven, Conn., 1959. Six filmstrips, $36.00.

WITH THE ADVENT of the age of space travel, astronomy is becoming an ever more popular subject and is assuming its proper place as an integral part of the elementary science curriculum. Science teachers, many of whom have little background in astronomy, are now required to conduct a unit in this subject. Planetarium demonstrations, television programs, and other popular presentations of astronomy are drawing larger audiences than ever before. These trends produce an increasing demand for quality teaching aids.

The Story of the Universe is a series of color filmstrips produced by Films for Education. While intended for an upper-grade school audience, the strips contain enough variety of pictorial material to be interesting to still younger groups. Also, the technical content and broad coverage make these films suitable for high school classes, adult laymen, and even college survey courses for nonconcentrators in astronomy.

Following the traditional approach to the study of astronomy, the series begins with the earth and proceeds outward. Unit I, "The Earth and Its Moons," contains six filmstrips.

1. "The Earth's Shape and Size" (56 frames) deals with several methods used by the ancients to prove that the earth is spherical and to measure its circumference. How their proofs and measurements are borne out by modern science is also shown.

2. "Motions of the Earth in Space" (65 frames) proceeds from a discussion of motion in general to a description of the rotation and revolution of the earth. As in the first strip, proofs from observed happenings are stressed.

3. "The Earth as a Planet" (67 frames) discusses the atmosphere's mass, density, and composition, and explains the radio-reflecting properties of the ionosphere. Causes of the seasons are also illustrated.

4. "Exploring the Space Around the Earth" (59 frames) first describes conventional aircraft propulsion and its inadequacies for travel beyond the earth's atmosphere. The section on rockets contains a thorough discussion of the law of action and reaction. Orbits and their properties are treated, as well as escape velocity.

5. "Information from Satellites" (63 frames) begins with the uses of artificial satellites. Radiation measurement raises the subject of the kinds of radiation man will encounter in space travel. Condi-



YOUR TELESCOPE IS NO BETTER THAN YOUR ABILITY TO FIND THE THINGS YOU WANT TO SEE

Now — for the first time

An entirely practical solution to the amateur's biggest problem. Now you can find celestial objects with an ease, speed, and accuracy never before approached.

To find any object you must know the sidereal time and **ACCUSTAR** will give it to you instantly. Having already set the outer disk to your longitude, turn the inner disk until your clock (standard) time corresponds with the date. The sidereal time is then indicated on the inner dial. It's as simple as that. Consistently accurate within two minutes or less.

ACCUSTAR will do many things: tell precisely where any star or constellation will be at any time or place, determine longitude, tell the solar time with one simple observation easily made with **ACCUSTAR.**

ACCUSTAR and the book, *Find the Stars* **$9.75**

A precision instrument, blue and gold in color, 10 by 12 inches, plastic coated, metal backed, weighs less than a pound. Nothing to get out of order.

Book *Find the Stars* has 23 sky maps and needed information on over 250 celestial objects of prime interest to the amateur.

ACCUSTAR will be your indispensable guide to the heavens and multiply your pleasure in exploring them. Order now for yourself or for the perfect Christmas gift.

ACCUSTAR P. O. Box 2806, Detroit 31, Michigan

tions in a spaceship – especially weightlessness – are touched upon, leading to an explanation of the difference between mass and weight. Finally, the space station and rockets of the future are mentioned.

6. "The Moon" (72 frames) reviews the composition of the moon, conditions on its surface, and its gravitational attraction. The moon's motion as a satellite – its distance and orbit – are explained along with its phases and influence on the tides. The question of why we always see the same side of the moon is answered, and some prominent lunar features are described.

Perhaps the most impressive aspect of the entire series is the excellent artwork in all the slides. The text is brief, compact, and well written. Clarity and continuity are stressed throughout. Frames of review questions are placed at intervals, providing an opportunity to test the effectiveness of the previous presentation. Coverage is as thorough as is required, and basic concepts of physics are introduced and explained where they apply to the astronomical topic presented.

The continuity of topics and text within the individual strips and among the films of the series is most helpful to the teacher with little knowledge of astronomy. For the lecturer already familiar with the fundamentals of astronomy, a series of separate slides would probably be preferable for talks on selected subjects.

The Story of the Universe is designed as an introduction to astronomy, and Unit I, "The Earth and Its Moons," makes an excellent beginning toward the fulfillment of this purpose.

Orders for films and requests for further information should be addressed to Films for Education, Audio Lane, New Haven, Conn. Prices are $7.50 for each strip, $36.00 for a unit of six filmstrips, and $144.00 for the entire set of four units. Units II, III, and IV are now in production.

VALERIE A. ROBERTS
Charles Hayden Planetarium

The latest
Spitz MODEL A-2
PLANETARIUM
has been installed at
Western Washington
College of Education
Bellingham, Washington
★
Spitz Laboratories
Inc.
YORKLYN, DELAWARE
Phone: CEdar 9-5212

U. S. ARMY RELEASE
PRISMATIC REFRACTING
TELESCOPES

Interchangeable 10X and 15X large-diameter, highly corrected wide-angle eyepieces. Solid bronze construction with tripod pivots, length 25 inches. Weight of scope 20 lbs.

Unusually well suited for astronomical, terrestrial, target-spotting and marine use. Easily altered to accommodate standard (1¼" O.D.) telescope eyepieces up to 100 power.

U. S. Government tripod, made of oak with bronze legs and fittings, collapsible, extends to eye level, Gov't. cost $66.00. Also special, aluminum offset yoke with locking device. Telescope can be mounted and dismounted in seconds. Tripod and yoke $25.00 additional.

Satisfaction guaranteed or money refunded.
No C.O.D.

BEN ROSENBERG
101 WEST 45TH ST., NEW YORK 36, N. Y.

EXCELLENT CONDITION
$49.00
Express Collect
Original cost
$500.00

Includes heavy-duty saddle-leather-fitted carrying case.

3¼" Achromatic Objective,
Built-in Filar Micrometer,
Huge Helical
Focusing Mount and
Prismatic Erecting System.

NEW BOOKS RECEIVED

How Old Is the Earth? *Patrick M. Hurley,* 1959, *Doubleday.* 160 pages. 95 cents, paper bound.

Modern knowledge of the age, origin, and structure of our planet is described by a leading geologist for the scientifically minded layman and for students. There is an explanation of dating rocks by their radioactivity.

This book is one of the new science study series being issued by the Physical Science Study Committee, formed in 1956 to improve secondary-school science teaching in the United States. Other books in the series are *The Neutron Story,* by Donald J. Hughes; *Magnets,* by Francis Bitter; *Echoes of Bats and Men,* by Donald R. Griffin; and *Soap Bubbles,* by C. V. Boys.

The Sun, *Karl Kiepenheuer,* 1959, *University of Michigan Press.* 160 pages. $5.00.

How the sun is observed and recent findings about solar phenomena are the themes of an illustrated book written in nontechnical language by an expert. This is a translation of a work first published in Germany in 1957.

Contemporary Geodesy, *Charles A. Whitten* and *Kenneth H. Drummond,* editors, 1959, *American Geophysical Union,* 1515 Massachusetts Ave., N.W., Washington 5, D. C. 95 pages. $5.50.

This volume contains the proceedings of a symposium on contemporary problems in the measurement of the size and shape of the earth. Satellites and other recent advances have accentuated the need for an international geodetic network. The symposium took place at the Smithsonian Astrophysical and Harvard observatories in December, 1958, with 107 scientists attending.

Astronomischer Jahresbericht, Vol. 57, 1959, *Astronomisches Rechen-Institut,* Heidelberg, West Germany. 525 pages. DM 60, paper bound.

The world's astronomical literature published in 1957 is comprehensively listed in this latest annual volume of a standard reference work. Titles are arranged by subject, with short abstracts in German for the more important references. The alphabetical subject index is in English.

The Realm of the Nebulae, *Edwin Hubble,* 1958, *Dover.* 207 pages. $1.50, paper bound.

This is a photo-offset reprint of Hubble's classic book of 1936, in which he reviewed the observational evidence that galaxies are other star systems, and summarized his findings as to their distances, sizes, and distribution in space.

While this work remains a landmark in the history of science, much of its contents is out of date. Hubble's scale of distances of galaxies, upon which most of his numerical conclusions depend, is now undergoing drastic revision. Hence the book should be read with caution, and cannot be recommended as a reference for numerical data.

Space Technology, *Howard S. Seifert,* editor, 1959, *Wiley.* 1,151 pages. $22.50.

This systematic textbook for graduate students covers the engineering basis of artificial satellites and space travel. The treatment is in five broad categories: flight dynamics, propulsion and structure, communications and guidance, psychophysical and biological problems of space travel, and research in space physics. The material is based on a course given by the University of California in co-operation with Space Technology Laboratories.

1. Moon – last quarter

2. Great Nebula in *Orion*

3. Spiral Nebula in *Triangulum*

4. Great Nebula in *Andromeda*

5. Saturn

6. Southern sector of Moon

8. Edge on view – Spiral Nebula in *Andromeda*

9. Spiral Nebula in *Canes Venatici*

7. Solar prominence

"The sky's the limit in decorating today. Now, outer space has been brought down to earth... Enlargements of astronomers' photographs of the Moon, Saturn ...and other heavenly objects brought down to wall size, prove to have an uncanny kinship with modern art.

"The pocked surface of the moon looks like a textural abstraction. The Whirlpool Nebula recalls the heavens as Van Gogh painted them."

from New York Times review of Astro Murals.

10. Moon – 14 days

$3.95 *(24" x 36" PRINT)*

THREE PRINTS FOR $11.
ENTIRE SET OF TEN PRINTS $35.

These superb reproductions are in use in leading universities and planetaria where they have received the endorsement of professional astronomers. A unique Xmas gift! Order now and avoid the Xmas rush.

astro murals 33 West 60th Street, New York 23, N. Y.

Gentlemen:

Please send the following circled items:

1 2 3 4 5 6 7 8 9 10

Name .....

Address .....

City ..... State .....

☐ Money order ☐ Check enclosed for sum of .....

Orders shipped postpaid in mailing tube. Sorry, no COD's.

★ ★ TRECKERSCOPE ★ ★

AMERICA'S NO. 1 TELESCOPE FOR AMATEURS AND PROFESSIONALS ALIKE . . .

THE COAST INSTRUMENT TRECKERSCOPE line has, over the years, established itself in amateur historical annals as one of the truly fine commercial instruments of all time. This by no means insinuates that our research and development has ceased for a second. Rarely has a precision, research-quality telescope such as the TRECKERSCOPE been in the price range of everyone. Here you buy the STANDARD OF COMPARISON of ALL telescopes!

Our own parabolic mirrors, NEW 1/20th wave or better, with matching elliptical diagonals, figured by one of the nation's finest opticians, from the finest, selected Pyrex, insure viewing second to none. Your choice of the world's largest selection of oculars (eyepieces) — yes, even including the superb ORTHO-STARS — your choice of any 3 oculars (except our 20-mm. Erfle ORTHO-STAR).

The second-to-none, super-rigid TRECKERMOUNT can be defined as the pinnacle of mechanical perfection. 1½" and 1¾" SOLID-STEEL shafts supported by massive aluminum equatorial head castings on thrust surfaces of Teflon produce velvet-smooth axis movement. This, in itself, gives sheer viewing enjoyment. The age-old problem of portable telescopes retaining their rigidity has been solved by modern engineering here at our factory. TRECKERSCOPES are easily assembled, or disassembled, in three minutes, and rock-steady performance is guaranteed.

8" DE LUXE TRECKERSCOPE • MADE IN U. S. A.

The time-proven HYDRO-GLIDE focusing device is used on all models. It eliminates all of the binds and backlashes of conventional rack-and-pinion movements — the HYDRO-GLIDE, and only the HYDRO-GLIDE, moves the eyepiece with a cushioned Neoprene roller, assuring satin-smooth focusing.

Our TRECKERFINDER is superb, producing a wide, flat field with resolution equal to high-priced 50-mm. refractors.

The following accessories are furnished with De Luxe models, and are available for the Standard: High-precision clock drive, giving effortless viewing, coupled with TRECKERSCOPE's exclusive manual flex-line slow-motion control for driving your instrument when 110-v. power is unavailable; fully rotatable tube system; sidereal time and declination setting circles (accurate to 0.001"); snap-out casters for convenience in moving your telescope from storage to viewing location.

AVOID LATER DISAPPOINTMENTS — BUY TRECKERSCOPE!

Our treatise, "WHAT YOU SHOULD KNOW, LOOK FOR, AND DEMAND BEFORE BUYING ANY TELESCOPE," now included in our free catalog — send for it!

TRECKERSCOPE PRICES

STANDARD 6"	$425.00	DE LUXE 6"	$675.00
" 8"	$525.00	" 8"	$775.00
" 10"	$775.00	" 10"	$1025.00
" 12½"	$1135.00	" 12½"	$1385.00
10" CASSEGRAIN "SKY-GIANT"			$1795.00

EXTENDED PAYMENT PLAN ON SCOPES AND MOUNTS

MIRRORS, PARABOLIC, GUARANTEED UP TO 1/20TH WAVE OR BETTER FROM COAST'S OWN OPTICAL SHOP, ALUMINIZED AND QUARTZ OVERCOATED

	6" f/10	6" f/8	8" f/7.5 or f/6	10" f/7.5 or f/6	12½" f/7.5 or f/6
1/20th WAVE		$82.50	$115.00	$255.00	$362.50
1/10th WAVE		$62.50	$95.00	$195.00	$282.50
1/8th WAVE	$47.50				
REFIGURING, NO CHANGE IN YOUR OWN FOCAL LENGTH					
1/20th WAVE		$47.50	$65.00	$110.00	$175.00
1/10th WAVE		$37.50	$55.00	$80.00	$125.00

ALL PRICES ABOVE INCLUDE MATCHING ELLIPTICAL DIAGONALS AND ALUMINIZING WITH QUARTZ OVERCOATING.

TREAT YOURSELF TO THE FINEST — Words alone cannot express the superb quality of COAST mirrors — they will positively amaze you with their "out-of-this-world" resolving power. This we guarantee you! Dawes' limit so easily!

TELESCOPE OPERATING MANUAL
by Raymond Dudley

Explains amateur telescope operation, including setting circles, sidereal time, alignment, orientation. Shows how to find 52 most beautiful objects in sky as well as general "how-to-observe" instructions. 44 pages of information every amateur needs.

$1.00 postpaid

HELAJUST FOCUSING MOUNT

Now at a new low price, but unexcelled in construction. Both push-pull for coarse adjustment, and helical movement for fine, precise focusing. Beautifully finished in black crackle and satin chrome, with a total extension of main tube and drawtube of 6", and engineered to fit all sizes of tubes without modification. Has single-vane-spider holding boss for those desiring this method of using diagonal. Guaranteed far superior to any other low-priced focusing mount, or **money back.** (Takes 1¼" eyepieces.) **$9.95**

ORTHO-STAR OCULARS

DUST SEALED — ALL COATED BEAUTIFUL FINISH

Guaranteed to be the finest you ever used— or return for full refund! Outstanding features: wide flat field, sharp to the very edge; extra-long eye relief; parfocalized for easy change of power; sealed-in optics, never need interior cleaning; hard coated, magnesium fluoride; boldly marked for easy identification; striking chrome and black-velvet finish, beautifully machined, 1¼" O.D.

ORTHO-STAR oculars are available in the following focal lengths, giving, for example, the indicated powers when used in conjunction with an 8" f/8 mirror: 27 mm.—61x; 20 mm.—81x; 16 mm.—102x; 10 mm.—163x; 7 mm.—233x.

$19.50 each ppd.

NOW AVAILABLE: 20-MM. ERFLE ORTHO-STAR OCULAR, 66°.7 apparent field. **$29.50 each ppd.**

TRECKER-PATHFINDER EQUATORIAL MOUNT

$89.50 complete

This mount will accommodate 4-inch to 8-inch telescopes. Specify your tube size when ordering.

Standard 36-inch height — massive 1½-inch steel shafting, in oil-impregnated bronze bearings.

This amazing EQUATORIAL MOUNT is just what the doctor ordered for mounting that homemade telescope you labored so hard to finish. Now you can purchase a beautifully constructed, highly rigid equatorial mount, COMPLETE, for your own telescope as economically as if you had built it yourself. This terrific mount is made entirely of metal; all of the moving equatorial parts are polished to work with maximum ease. Legs, head, and counterweight are all removable for easy storing. The saddle allows complete rotation of your tube. One of the more important features in this mount is that the polar axle is extended for ease in attaching a clock drive and/or setting circles, which may be added at any time. The TRECKER-PATHFINDER mount also has a beautiful, chip-resistant finish. Taking all of these unusual features into consideration, this is truly one of the best DOLLAR-FOR-DOLLAR values.

USE OUR EXTENDED PAYMENT PLAN ON ALL SCOPES AND MOUNTS



HYDRO-GLIDE Precision Focusing Device

Coast Instrument's own "Hydro-Glide" (type formerly referred to as "rack and pinion"). Now you can have WHISPER-SMOOTH control with absolutely no high and low spots as geared units have. This has never before been available in **any** focusing mount. Standard in appearance, yet utilizing an entirely **new** concept. We guarantee you will be astonished at its unbelievable superiority. If this isn't the absolute "smoothest," **return it for full refund.** **$18.50 patent pending**

MIRROR CELLS

Skeleton type

6"	$6.95	10"	$17.95
8"	$10.95	12½"	$21.00

4-VANE SPIDERS

6"	$11.50	10"	$14.95
8"	$11.50	12½"	$16.95

TRECKER FINDER

7x, 50-mm. objective, helical focusing, with mounts and crosshairs. Same as used on TRECKERSCOPES. **$18.50**

Send for Coast Instrument's Professional MIRROR CLEANING KIT. **$2.35 ppd.**

SEND FOR OUR NEW 68-PAGE CATALOG showing many of the other telescope items manufactured or distributed by COAST INSTRUMENT. It includes our treatise, "WHAT YOU SHOULD KNOW, LOOK FOR, AND DEMAND, BEFORE BUYING **ANY** TELESCOPE," an impartial, informative article that will give you information you've probably been looking for to help make up your mind as to just what telescope you should buy!

All prices, unless otherwise indicated, are at our warehouse in Long Beach, California, from where shipping charges will be added, and are subject to change without notice. Nominal crating charge added for all telescopes and mounts. California residents: Add 4% sales tax to all prices. NOW APPOINTING DEALERSHIPS . . . INQUIRIES INVITED.

TRECKERSCOPES ARE CUSTOM MADE IN AMERICA BY PROFESSIONAL OPTICIANS AND ENGINEERS.

COAST INSTRUMENT, INC.

"IN OPTICS SINCE 1933"

4811 Long Beach Blvd., Long Beach 5, Calif.

Phone: GArfield 2-3411 or NEvada 6-7683

GLEANINGS FOR ATM's

CONDUCTED BY ROBERT E. COX

A Low-Power Microscope System for High Magnifications

DURING the 1958 opposition of Mars, for my evening astronomy class at the Wilke Observatory of Macalester College, I tried projecting the planet's image directly from the telescope onto a white screen. The 8-inch f/8 reflector and a good 1-inch eyepiece gave a much larger and brighter image than I had thought possible.

Since the 8-inch reflector is Springfield mounted for the observing convenience of visitors and students, the focal plane is not extended very far from the last prism in the optical train. This helps reduce the size of the prisms and lowers diffraction and light loss. However, it prevents the use of a Barlow lens for increased magnification, and all high powers have heretofore been obtained using short-focal-length eyepieces with little eye relief.

Dr. Sherman Schultz, Jr., demonstrates his method of amplifying telescope magnification, on the 8-inch reflector at the Wilke Observatory of Macalester College. He uses a positive lens at the diagonal eyepiece holder, and the ocular is at the end of the long extension tube. The author has supplied all photographs with this article.

While looking at the projected image, the thought suddenly occurred to me that here was half of the optical system of a low-power microscope. If another eyepiece were placed behind the image, in place of the projection screen, then it would be viewing the enlarged image with a great increase in magnification.

In an ordinary microscope, an objective of short focal length is used to form an enlarged image of an object by projection, the image then being examined by a magnifier or eyepiece. While conventional microscopes have a standard separation between the optics, here that spacing could be varied over a wide range to give a corresponding range in amplification. Experience has shown that this arrangement is practical.

The explanation of the method is simple. In a telescope, an eyepiece with an EFL (equivalent focal length) of 1″ has its principal plane 1″ from the image formed by the main objective, and the emerging beams of light are parallel. If the eyepiece is withdrawn twice its focal length from the primary image, in this case 2″, it will project a secondary image equal in size to the primary one, but inverted and 2″ behind the eyepiece. This system would have no amplification, only transferring the image to a position some 4″ or 5″ from the primary focal plane (the thickness of the eyepiece assembly must be added, too).

If the eyepiece is slowly brought closer to the primary image, the projected image swiftly moves to a greater distance and its size increases. In a 1-inch projection system, the amplification is half the distance between the eye lens of the eyepiece and secondary image. Thus, if the image is 4″ away, its size will be doubled; 6″, tripled. The use of another 1-inch eyepiece to



A 1-inch eyepiece placed to act as a transfer lens with no amplification. Note that the new image has been erected.

Precision Diagonals

You will get the best possible performance from your telescope with one of our clear **fused quartz** diagonals. Accuracy guaranteed 1/20 wave.

Ellipse 1.25" x 1.77" . . $12.00
Ellipse 1.5" x 2.12" . . $15.00

Pyrex diagonals, 1/8-wave accuracy.

Ellipse 1.25" x 1.77" . . $ 6.00
Ellipse 1.5" x 2.12" . . $ 9.00

Without aluminum coating, deduct $1.00.

Send for our complete list of supplies, quartz mirrors, blanks, oculars, coatings, and accessories.

E & W OPTICAL CO.
2406 E. Hennepin Ave.
Minneapolis 13, Minn.

OPTICAL FLATS

Pitch Polished, Beral Coated

Rectangular shape
1⅝" x 1⅞"
½ wave $1.75 each
¼ wave $4.50 each
Postpaid.

Elliptical shape
Heavy edges to minimize temperature effects.
1¼" minor axis.
½ wave $4.00 each
¼ wave $6.50 each
Postpaid.

BERAL COATINGS — same optical characteristics as aluminum — mechanically more durable — not over-coated — may be removed without harming glass surface. Prices for Beral coating telescope mirrors: 4" diameter - $2.75, 6" - $3.50, 8" - $4.50, 10" - $6.50, and 12½" - $9.75 each, f.o.b. Skokie.

LEROY M. E. CLAUSING
8038 Monticello Ave. Skokie, Ill.

TELESCOPE MAKERS

Everything for the Amateur

Mirror-Making **KITS . . . . . $4.50 up**

Thousands of our customers with no special training have built powerful precision telescopes with our quality supplies, instructions, and guidance.

Send for Complete Telescope Making Instructions, 10c

ALUMINIZING

Superior Reflecting Surface.
Guaranteed not to peel or blister.
Mirrors Made to Order — Mirrors Tested Free

EYEPIECES
CELLS
PRISMS
BINOCULARS

PRECISION TRI-TURRET

Holds 3 standard 1¼" O.D. eyepieces. Smooth turn to grooved notch aligns eyepiece precisely, ready to focus for various powers. Suitable for reflectors or refractors. **$15.75**

ORTHOSCOPIC EYEPIECES
HIGHEST QUALITY • MADE IN U.S.A.

Special four-element design, with fluoride-coated lenses, gives a wide flat corrected field. Standard 1¼" O.D. — E.F.L. 6-8-12-16-24-mm. ..... Postpaid **$15.95**

Write today for FREE Catalog.

Precision Optical Supply Co.
1001-T East 163rd St., New York 59, N. Y.



For extreme amplifications one or more 1¼″ tubes may be coupled together. Here two of them are joined by a piece of metal tubing. With a 1-inch projection lens at the left, and a 1-inch eyepiece at the right, this arrangement increases the magnification by a factor of about six.

view the projected image will yield respective powers two and three times normal for the telescope used. If a ½-inch eyepiece is used to view the projected image, the power will be increased four and six times, respectively.

Since eyepieces are designed primarily to emit parallel light, I decided to try a system that was specifically designed for projection purposes, and selected a good f/1.6 8-mm. motion-picture projector lens of 1″ focus. But no perceptible difference in image quality could be detected in comparison with a good orthoscopic eyepiece. Therefore, if you have an excellent ocular that can be used as a projection lens, it is unnecessary to buy any extra optics.

The table shows the amplification effect with various eyepiece combinations and separations, using a 1-inch eyepiece for projection.

TABLE OF AMPLIFICATION

Distance of projected image	*Viewing eyepiece*		
	1-inch	*½-inch*	*¼-inch*
2″	1x	2x	4x
4″	2x	4x	8x
6″	3x	6x	12x

For assembling the low-power microscope unit, chrome plumbing tubing is quite suitable. It comes with an outside diameter of 1¼″ along most of its length, and the flared-out section at one end has an inside diameter of 1¼″. To prevent internal reflections the inside should be painted black. If necessary, several tubes can be fastened together by couplings, as shown in the picture above.

The method of mounting the unit will depend on the construction of the projection eyepiece. If the eye cap can be removed without disturbing the eye lens, then the eyepiece can be slipped into the 1¼″-inside-diameter section of tubing. If this cannot be done, then a little more ingenuity must be used. Be sure that the



If a 1-inch eyepiece is used, it must be placed less than two inches from the primary image to project an image that is enlarged—two times in this example.

Astronomical

featuring our

SPACE K-4 DELUXE

TELESCOPES

by SPACEK INSTRUMENT CO.

CONSIDER THESE PROFESSIONAL-TYPE FEATURES

- **Optical system:** 4¼-inch mirror, f/11, spherically corrected with surface accurate to ⅛ wave, aluminized and quartz over-coated.
- **Mounting:** Equatorial and adjustable for latitude. Massive aluminum castings with 1-inch-diameter shafting on both axes and Tensalloy bearings. Clamps provide any degree of tension desired. Two lock-nuts on saddle permit simple removal of tube. All steel parts chrome plated and carefully machined to give maximum rigidity.
- **Tube:** Seamless aluminum, painted black inside, with hammer-tone green finish.
- **Rack-and-pinion focusing:** Accommodates standard 1¼-inch eyepieces.
- **Finder:** 6 power, with achromatic objective, and mounted in adjustable bracket.
- **Pedestal:** Heavy-wall steel tube with collapsible legs. Guaranteed free from torque and wobble. Design allows tube to be swung in any position without bumping into legs. Pedestal, mount, and legs painted black crinkle finish.

$89.50

- **Three eyepieces:** 50x Kellner, 100x and 200x Ramsdens.
- **Instructions.**

WANTED

Professional and Amateur Astronomers

Put your ideas and experience to work. Extensive expansion program by Spacek Instrument Co. requires individuals with a knowledge of telescopes, optics, and astronomy.

Proven design, low cost, and excellent performance of our SPACE K-SERIES telescopes have met with the approval of many schools, observatories, and clubs.

Here is an excellent opportunity for responsible persons, who are interested in supplementing their income, to demonstrate and sell our telescopes.

Write for complete information about our Individual Dealer Plan.

All prices subject to change.
Write for free literature.
DEALER INQUIRIES INVITED

OTHER SPACE K-SERIES TELESCOPES AND ACCESSORIES

SPACE K-6 (6-inch Equatorial)	$249.50
SPACE K-3 (3-inch Equatorial)	$ 39.50
SPACE K-SATELLITE (3-inch Altazimuth)	$ 29.50
RACK AND PINION (fits any size tube)	$ 8.50
6-POWER ACHROMATIC FINDER	$ 7.25
MIRROR CELLS: 6-inch, $6.50; 8-inch, $10.50; 10-inch, $19.50	

Prices are f.o.b. Pottstown, Pa.
Pennsylvania residents add 3% sales tax to all prices.

SPACEK INSTRUMENT CO.

1130 Sembling Ave.
Pottstown, Pa.
Telephone: FAculty 3-2825

eye lens of the projection eyepiece is turned toward the new image plane.

Has this system any advantages over the use of a Barlow lens? First, if you already have good eyepieces, it is not necessary to purchase any more optical parts. Second, since the usual Barlow's diameter is smaller than that of the incoming cone of light from the primary mirror, there is a decided narrowing of the field and also vignetting. Because the low-power microscope system is placed behind the focus at a position where it intercepts the full cone of light from the primary, it has a minimum light loss and vignetting. Third, from two or three good eyepieces almost any reasonable power can be secured with a modest collection of extension tubes. These eyepieces need not be high-power ones, probably no shorter than a ½-inch.

It is most important to keep the tubes mechanically stable and rigid, as bending can cause serious image distortion and poor focus. The Springfield mounting, with its stationary eyepiece location, is especially suited for this system. On German- or fork-mounted Newtonians, care should be taken not to overbalance the telescope, setting up unwanted vibrations.

SHERMAN W. SCHULTZ, JR.
944 Cobb Rd.
St. Paul 17, Minn.



Fig. 1. A reduction of almost one half can be made in the size of a Newtonian diagonal by the introduction of a transfer lens working at unity power, as explained in the text below.

SOME ADVANTAGES OF A TRANSFER LENS

DR. SCHULTZ'S method of projecting the prime-focus image of a telescope has important advantages when the object and image are equidistant from the projection lens. Then it simply transfers the image from one focal plane to another, without change in magnification. If the lens is of good quality, no serious aberrations are introduced by the transfer.

With this system, a smaller diagonal can be used in a Newtonian reflector. Fig. 1 shows a 6-inch reflector fitted with a 1-inch orthoscopic eyepiece as a transfer lens. The diagonal would normally be placed 5.5″ inside the primary focus, and it would have a minor axis of 1.11″. With the transfer lens, the diagonal need be only 1.5″ inside the primary focus, and its shortest diameter can be 0.65″. Only 1.2 per cent of the incoming light is blocked by this diagonal, instead of 3.4, and there is less diffraction from the shorter circumference. It is also possible to place a field stop near the new focal plane to remove stray light during daytime observing.

In designing a Newtonian that uses a positive lens to reduce diagonal size, care should be taken that the focus is either in front of or entirely behind the diagonal itself. Were any part of the flat mirror to lie in the transfer-image plane,

MAGNUSSON TELESCOPES AND ACCESSORIES

- New mountings as low as $29.00 and $59.00.
- New low-priced clock drives.

Parts are sold separately. Write for prices.

O. MAGNUSSON
14570 W. 52nd Ave., Arvada, Colorado

MAKE A FINE NEWTONIAN TELESCOPE FOR LITTLE MONEY

No need for months of tedious grinding. Telescope success is assured you. For those who wish to make their own, we supply COMPLETED (machine-made) MIRRORS, ALUMINIZED, with lenses for eyepieces and flat, as follows:

3″ Kit, 39″ f.l., 40 and 80 power
$7.00 + 25¢ postage
4″ Kit, 54″ f.l., 60 and 120 power
$12.00 + 40¢ postage
6″ Kit, 63″ f.l., 65 and 130 power
$20.50 + $1.00 postage
8″ Kit, 85″ f.l., 85 and 170 power
$55.00 + $4.00 postage



Send for your telescope kit NOW.
Free instructions with each kit.

HARRY ROSS Telescopicals
59 to 63 Reade Street
New York 7, New York

DE LUXE
SKYSCOPE

As illustrated — with new plastic tube, 6x finder, heavy all-metal tripod, equatorial mounting, and 60x eyepiece.

Price **$44.75**

Unconditionally guaranteed.
100% American made.

Write for free brochure showing our standard model which has been sold world-wide for over 20 years. Skyscope still remains outstanding in value by any comparison.

THE SKYSCOPE CO., INC.
P. O. Box 335, Brooklyn 28, N. Y.

Mounted Eyepieces

The buy of a lifetime at a great saving. Perfect war-surplus lenses set in black-anodized standard aluminum 1¼" O.D. mounts.

F.L.	TYPE	PRICE
6 mm. (¼")	Ramsden	$ 4.75
12.5 mm. (½")	Ramsden	4.50
12.5 mm. (½")	Symmetrical	6.00
16 mm. (⅝")	Erfle (wide-angle)	12.50
16 mm. (⅝")	Triplet	12.50
18 mm. (¾")	Symmetrical	6.00
22 mm. (27/32")	Kellner	6.00
27 mm. (1-1/16")	Kellner	4.50
32 mm. (1¼")	Orthoscopic	12.50
35 mm. (1⅜")	Symmetrical	8.00
55 mm. (2-3/16")	Kellner	6.00
56 mm. (2¼")	Symmetrical	6.00

COATED LENSES 75 cents extra.

"Giant" Wide-Angle Eyepieces

ERFLE EYEPIECE (65° field) contains 3 coated achromats. 1½" E.F.L., clear aperture 2⅛". Has a focusing mount with diopter scale. Will make an excellent 35-mm. Kodachrome viewer. Magnifies seven times ..... **$12.50 ppd.**

Same as above without diopter scale **$9.95**

1¼"-diam. Adapter for eyepiece above **$3.95**

WIDE-ANGLE ERFLE (68° field) **EYEPIECE.** Brand new; coated 1⅛" E.F.L. Focusing mount. 3 perfect achromats, 1-13/16" aperture **$13.50**

1¼"-diam. Adapter for eyepiece above **$3.95**

WIDE-ANGLE ERFLE 1½" E.F.L. Brand new; contains Eastman Kodak's rare-earth glasses; aperture 2"; focusing mounts; 65° field **$12.50**

1¼"-diam. Adapter for eyepiece above **$3.95**

Astronomical Mirrors

These mirrors are of the highest quality, polished to ¼-wave accuracy. They are aluminized, and have a silicon-monoxide protective coating. You will be pleased with their performance.

	Diam.	F.L.	Postpaid
Plate Glass	3-3/16"	42"	$ 9.75
Pyrex	4¼"	45"	13.50
Pyrex	6"	60"	25.00

Mirror Mount

Cast aluminum. Holds all our mirrors firmly with metal clips. Completely adjustable. Assembled.

3-3/16"	Mount fits our 4½"	tubing	$4.00 ppd.
4¼"	Mount fits our 5"	tubing	4.00 ppd.
6"	Mount fits our 7"	tubing	6.50 ppd.

Rack-&-Pinion Eyepiece Mounts



Here is a wonderful opportunity for you to own a most mechanically perfect Rack-&-Pinion Focusing Eyepiece Mount with variable tension and adjustment. Will accommodate a standard 1¼" eyepiece, positive or negative. The body casting is made of lightweight aluminum with black-crackle paint finish, focusing tube of chrome-plated brass. Focusing tube for refractors has a travel of 4", for reflectors 2", and will fit all size tubing.

REFRACTOR TYPE for 2⅛" I.D. Tubing	$12.95 ppd.
" " for 3¼" I.D. Tubing	12.95 ppd.
" " for 4⅛" I.D. Tubing	12.95 ppd.
REFLECTOR TYPE (less diagonal holder)	8.50 ppd.
DIAGONAL HOLDER	1.00 ppd.

Special Coated Objective

BIG 2⅛" DIAM. — 15" F.L. — $6.00

These achromatic objective lenses are tested and have the same high quality as "Big Lenses" described above, except they are seconds for slight edge chips or small scratches only. Quality guaranteed. **ONLY $6.00 ppd.**

AIRSPACED OBJECTIVES

MOUNTED IN ALUMINUM CELLS f/15

We offer the lowest priced, hand-corrected, precision, American-made astronomical objective, mounted in a black-anodized aluminum cell. Our reputation for high-quality astronomical lenses has established us as the most reliable source in the industry.

"Those in the know" BUY FROM US BECAUSE:

Each lens is thoroughly tested by us and is guaranteed to resolve two seconds of arc or better. They are corrected for the C and F lines (secondary chromatic aberration). The zonal spherical aberration and the chromatic variation of spherical aberration are negligible. The cell is machined to close tolerances so that it will fit directly over our standard aluminum tubing, eliminating any mounting problems.

3¼" diam., 48" F.L. (uncoated)	$28.00	4⅛" diam., 62" F.L. (uncoated)	$60.00
Same as above with coating	$32.00	Same as above with coating	$69.00

We can supply ALUMINUM TUBING for the lenses above.

"BIG" ACHROMATIC TELESCOPE OBJECTIVES

We have the largest selection of diameters and focal lengths in the United States available for immediate delivery. These are perfect magnesium-fluoride coated and cemented Gov't. surplus lenses made of finest crown and flint optical glass. Not mounted. Fully corrected. Tremendous resolving power. They can readily be used with eyepieces of only ¼" focal length, thereby producing high powers. Guaranteed well suited for astronomical telescopes, spotting scopes, and other instruments. Gov't. cost up to $100.

Diameter	Focal Length	Each	Diameter	Focal Length	Each
54 mm. (2⅛")	254 mm. (10")	$12.50	83 mm. (3¼")	660 mm. (26")	$28.00
54 mm. (2⅛")	300 mm. (11.8")	12.50	83 mm. (3¼")	711 mm. (28")	28.00
54 mm. (2⅛")	330 mm. (13")	12.50	83 mm. (3¼")	762 mm. (30")	28.00
54 mm. (2⅛")	390 mm. (15.4")	9.75	83 mm. (3¼")	876 mm. (34½")	28.00
54 mm. (2⅛")	508 mm. (20")	12.50	83 mm. (3¼")	1016 mm. (40")	30.00
54 mm. (2⅛")	600 mm. (23½")	12.50	102 mm. (4")	876 mm. (34½")	60.00
54 mm. (2⅛")	762 mm. (30")	12.50	108 mm. (4¼")	914 mm. (36")	60.00
54 mm. (2⅛")	1016 mm. (40")	12.50	110 mm. (4⅛")*	1069 mm. (42-1/16")	60.00
54 mm. (2⅛")	1270 mm. (50")	12.50	110 mm. (4⅛")	1069 mm. (42-1/16")	67.00
78 mm. (3-1/16")	381 mm. (15")	21.00	128 mm. (5-1/16")*	628 mm. (24¾")	75.00
80 mm. (3⅛")	495 mm. (19½")	28.00	128 mm. (5-1/16")	628 mm. (24¾")	85.00
81 mm. (3-3/16")	622 mm. (24½")	22.50	*Not coated		

● We can supply ALUMINUM TUBING AND CELLS for the lenses above. ●

NEW! 6" LENSES

Our 6" objective does not need high-pressure salesmanship. Its sparkling performance speaks for itself. Test one, or have any qualified person test it; we are certain that you will be satisfied. If not, take advantage of our money-back guarantee.

6"-DIAM. AIR-SPACED TELESCOPE OBJECTIVE

Hard-coated on 4 surfaces

f/10 - 60" focal length	MOUNTED	$175.00
	UNMOUNTED	150.00
f/15 - 90" focal length	MOUNTED	175.00
	UNMOUNTED	150.00

We can supply ALUMINUM TUBING for the lenses above.

Coated Binoculars

Beautifully styled imported binoculars. Precision made.

LOW, LOW PRICES!

American Type "Zeiss" Type

Complete with carrying case and straps. American type offers a superior one-piece frame and clean design.

Size	Field at 1,000 yards	Type	Center Focus	Ind. Focus
6 x 15	360 ft.	Opera	—	$12.75
6 x 30	395	"Zeiss"	$18.75	16.75
7 x 35	341	"Zeiss"	20.75	17.95
7 x 35	341	American	23.50	—
7 x 35	378	Wide-Angle 11°	35.00	—
7 x 50	372	"Zeiss"	24.95	22.50
7 x 50	372	American	32.50	—
8 x 30	393	"Zeiss"	21.00	18.25
10 x 50	275	"Zeiss"	28.75	26.75
20 x 50	185	"Zeiss"	33.75	31.75

All prices above plus 10% Federal tax.

8-POWER ELBOW TELESCOPE

This M-17 telescope has a brilliant-image 48° apparent field — 325 feet at 1,000 yards. The telescope can be adjusted for focusing 15 feet to infinity. It has a 2" objective, focusing eyepiece 28-mm. focal length with an Amici erecting system. Turret-mounted filters: clear, red, amber, and neutral. Lamp housing to illuminate reticle for nighttime use. Truly the biggest bargain you were ever offered. Original Gov't. cost $200.

BARGAIN PRICE	Uncoated	$13.50 ppd.
	Coated	$17.50 ppd.

GIANT "3" inch TELESCOPE

40 POWER

ALL LENSES COATED

HIGH-POWER SCOPE AMERICAN MADE

Big 3"-diameter, achromatic, coated objective will give you the brightest crystal-clear images. Micrometer spiral focusing drawtube, lightweight aluminum construction throughout, tough black crackle finish. Length open 22", closed 15½". Upright image. Guaranteed to perform flawlessly. **$57.50 ppd.**

TRIPOD FOR TELESCOPE ABOVE

Sturdy, lightweight, all-metal, black crackle finished tripod. Ball-swivel joint allows easy directional adjustments. Equipped with a standard ¼ x 20 stud which will fit any photo tripod. Stands 11" high, weight 1½ lbs. Comes with web straps. **$8.95 ppd.**

● **"MILLIONS" of Lenses, etc.** ●
Free Catalogue

We pay the POSTAGE in U. S. — C.O.D.'s you pay postage. Satisfaction guaranteed or money refunded if merchandise returned within 30 days.

A. JAEGERS

691 S Merrick Road
LYNBROOK, N. Y.

dust or surface imperfections would be in focus in the final image.

Springfield-mounted telescopes can also benefit from transfer lenses. Instead of the large diagonal mirror conventionally employed to bring the light down the declination axis, a significantly smaller flat can be used. With careful design, a transfer lens of several inches focal length allows the primary diagonal to be reduced in size in the same proportion as in our example of the Newtonian reflector.

Horace E. Dall has designed a compound telescope in which the secondary focus is between the primary and secondary mirrors, but is brought to a new focus behind the primary by a high-quality triplet lens. (This plan is described in *Scientific American* for May, 1939; December, 1947; September, 1951; and April, 1952.) The image is erect and a field stop permits daytime viewing even with an open tube. I built an 8-inch reflector of this type, and obtained fine results with a war-surplus erecting system for a unity-power transfer lens. I use a 2-inch secondary mirror and a field stop, and for terrestrial viewing the brilliance and color rendition of the images leave little to be desired.

Perhaps the most interesting application of a positive transfer lens is a pancratic system — one with alternative effec-



Fig. 2. The principle of a pancratic lens system, part "a" showing the reversibility of the light paths, while "b" and "c" illustrate actual systems. If the focal planes are to remain fixed, the projection lens may be placed in only two positions.

tive focal lengths but having fixed focal planes. This is made possible by the action of conjugate foci and the reversibility of a light path in an optical system. In Fig. 2a, a triplet lens of about 1.2″ focal length is 2″ from an object and forms an image 3″ on the other side of the lens. If the object is transferred to the image plane, light passing in the reverse direction through the lens (dashed lines in the drawing) will form a new image 2″ from the lens. But the sizes of the images will be different, the one nearer the lens being the smaller.

Figs. 2b and 2c show the application of such a lens to the Newtonian reflector of Fig. 1. If the focal length of the primary mirror is 48″ and the image size 1/100

When you build your telescope build a good one.

CLEVELAND PARTS

are designed to match and machined to fit

YOU'LL put a lot of time and work into your telescope before it's finished. If you want a fine, professional-looking instrument at low cost, and smooth, trouble-free performance, use guaranteed Cleveland parts. They're first choice of ATM's because they go together easily without special tools. You'll be proud of your Cleveland.

CLEVELAND EQUATORIAL MOUNT

1. Rotating base permits you to set up anywhere, then easily align and lock the polar axis on north. Makes accurate locating and tracking much easier.
2. Quick-adjusting, positive-locking tube clamp with concealed stop permits rotation of tube in saddle, keeps tube in correct balance, prevents sliding.
3. Husky, vise-grip latitude adjustment is easy to change, stays where you put it. We set it to your latitude before shipping.
4. Rigid, high-strength aluminum construction eliminates excessive weight, prevents rust.
5. All parts accurately machined to close tolerances for smooth, trouble-free operation. Designed for easy addition of accessories.
6. Accurately ground and polished steel axles, extra-long bearings and large brakes provide free movement and precise control.
7. Sturdy, cast-aluminum, tapered channel legs open to fixed position, require no folding braces. Close easily for carrying.

SPECIFICATIONS: Heavy-duty saddle, tube clamp, axle bearings, rotating base, tripod top, tapered channel legs, and extra pier top are cast aluminum. Axles are 1¼″ ground and polished steel. Bearings are 5½″ long. Large-area brakes, knurled bronze-aluminum alloy adjusting knobs. Brass thrust washers at all friction points. Knurled knobs lock rotating base in V groove in tripod top, release for easy transfer to permanent pier. Latitude is adjustable 0° to 55°. Latitude bolt, leg bolts, nuts, washers cadmium plated. Weight 23 lbs.

PRICE LIST

EQUATORIAL MOUNT Complete, for 7″ tube	$ 79.50	f.o.b.
Without legs	69.50	"
Complete, for tubes to 9″ x 60″	89.75	"
Without legs	79.75	"
Pier top for permanent mount	5.95	"
COUNTERWEIGHT 12½ lbs.	4.95	"
Extra weight, per pound	.50	"
ALUMINUM TUBE 7″ O.D. x 60″	14.85	"
6″ MIRROR CELL for 7″ tube	8.35	ppd.
DIAGONAL HOLDER for 1¼″ x 1¾″ x ¼″ elliptical or rectangular mirrors or 1¼″ prism. (Specify diagonal and tube size)		
For 7″ diameter tube	7.95	"
For tubes up to 10″ diameter	9.15	"
1¼″ EYEPIECE HOLDER	7.95	"
SETTING CIRCLES 1″ to 2″ bore. (Specify shaft diameter) Per pair	12.75	"
COMPLETE PARTS and instructions for building the 6″ Cleveland, except optical parts	129.75	f.o.b.
6″ PYREX MIRROR f/8 1/10 wave, parabolic, beryllium coated, with matching 1/10 wave, 1.25″ x 1.77″ diagonal	67.50	ppd.

Add 5% sales tax on Ohio orders.

COUNTERWEIGHT Lead and antimony. 3¼″ diam. x 3″ with 1″ or 1¼″ bore. Weight 12½ lbs.

TELESCOPE TUBE Extruded aluminum. 7″ O.D. x 0.072″ wall x 60″ length. Not machined.

SETTING CIRCLES 5″ cast aluminum with machined faces. ½″-wide scales are matte-white aluminum with large, legible figures and accurate black graduations. 1″ to 2″ bore. Specify shaft size.

6″ MIRROR CELL Solid plate protects mirror. Ring housing reduces convection currents. Improved clips hold mirror without pressure. Shockproof, cushioned adjustments prevent vibration, keep mirror in collimation. No springs. Cast aluminum, machined for 7″ O.D. tube.

DIAGONAL HOLDER Fully adjustable, shockproof design. One turn moves mirror position 0.050″. Three screw adjustments give you perfect control of mirror angle, make collimating easy. Made for 1¾ x 1¾ elliptical or rectangular mirrors, or 1¼″ prisms, tubes up to 10″. Specify mirror type, size, and tube I.D. when ordering.

1¼″ EYEPIECE HOLDER Acme-threaded brass sleeve in accurately machined, cast-aluminum housing gives you smooth, micrometer-sharp focusing. One turn moves eyepiece 0.125″. Stays where you set it. Only one moving part.

CLEVELAND ASTRONOMICS

BOX 209, CHAGRIN FALLS, OHIO

DELUXE PYREX
Reflecting Telescope Kits

Our kits have PYREX mirror blank, PYREX tool the same thickness, ample supply of optical quality abrasives, fast-polishing cerium oxide, red rouge and pitch. Packed in metal cans.

Size	*Thickness*	*Price*
4¼"	¾"	$ 6.00
6"	1"	$10.50
8"	1½"	$18.75
10"	1¾"	$33.65
12½"	2⅛"	$59.95

ADD POSTAGE: 1st and 2nd postal zones from Detroit, add 5%; 3rd and 4th, add 10%; 5th and 6th, add 15%; 7th and 8th, add 20%. Or we will ship C.O.D.

Send for free catalog of supplies, accessories, and refracting telescopes.

ASHDOWNE BROS.
18450 Grand River Detroit 23, Michigan

NOW!
Use BOTH Eyes for Your Visual Observing

Your telescope virtually becomes a giant binocular when used in conjunction with our newly developed double ocular attachment (one eyepiece for EACH eye).

Fits standard 1¼" focus tube unless otherwise specified. Complete price, including matched wide-angle oculars

only **$95.00**
postpaid anywhere in the world

Order your double eyepiece holder today. Checks or money orders accepted.

BRAYTON OPTICAL EXCHANGE
3343 Park Vista Dr., La Crescenta, Calif.

TRIGARTH TURRET *and* **Eyepiece Attachment with Rack and Pinion**

Just turn the Trigarth Turret and easily improve the performance of your telescope. It holds three eyepieces of standard 1¼" O.D. The Trigarth Turret sells for **$15.95 postpaid.**

Also priced at **$15.95 postpaid,** the Eyepiece Attachment with Rack and Pinion takes standard 1¼" O.D. eyepieces. The rack and pinion is machined from solid aluminum castings, precisely fitted for smooth performance. The main tube is 1¾" long; sliding tube adds 2"; total movement 3¾". Choice of gray or black crinkle finish.

Both Turret and Eyepiece Attachment, **$31.90.**

ELLIPTICAL DIAGONAL HOLDER

Accurately machined from solid brass to fit 1¼" minor axis elliptical diagonal. Fully adjustable for rotational and longitudinal movement. Guaranteed to keep your diagonal precisely and securely in proper position at all times. **$10.50 ppd.**

MIRROR CELLS
6" — $7.00; 8" — $11.50; 10" — $35.00

BUILD YOUR OWN TELESCOPE
Prisms Lenses Eyepieces
Mirrors ground to your order
Aluminizing—with Quartz Coating
Satisfaction Guaranteed

Write for Free Catalog. Instructions, 10¢.

GARTH OPTICAL COMPANY
P. O. Box 991 Springfield 1, Mass.

of this or 0.48", it will cover 34.3 minutes of arc. A pancratic transfer lens, if 3" from this image (Fig. 2b), forms a new focus 2" behind the lens, and the image size is 0.32". A 1-inch eyepiece that gave a power of 48x on the original image now gives 32x, only ⅔ as much.

When the lens is moved to 2" from the primary image, the new focus is 3" behind the lens. Now the image that was originally 0.48" in diameter becomes 0.72", an increase of 1½ times; the 1-inch eyepiece will give 72x. Shifting the projection lens by 1" more than doubles the image size. With the flick of a finger, the observer can change the power of any eyepiece by a factor of 2¼. It is startling to see this change in the field of view, especially when observing the moon. In practice, the pancratic arrangement should be restricted to small movements of the lens.

To determine the focal length, f, of a lens suitable for a particular set of object and image distances, o and i, we may use

$$1/f = 1/o + 1/i.$$

In our example, o is 2 and i is 3, giving 1/0.833 for $1/f$, whereby $f = 1.2$, the focal length of the lens in inches.

The projected image size, I, is given by $I = O\,i/o$, where O is the size of the object, that is, of the primary image before projection.

Because of the short focal lengths involved, an extremely high-quality transfer lens is required in a pancratic system. Two achromats of twice the desired focal length might work if placed crown lens to crown lens. Hastings triplets, designed for magnifiers, are generally of low quality, as they are mass-produced.

Hold the primary-focus image plane diameter to about 1/100 the telescope's focal length, or curvature of field and distortion may make the projection system impractical.

R. E. C.

TELESCOPE CAMERA HOLDER

A HOLDER for securing a 35-mm. camera over the eyepiece of a reflecting telescope can be made quite inexpensively — perhaps for only two dollars. The basic parts are angle iron, a photoflash unit arm, and nuts and bolts.

A 2" piece of angle iron is cut from a length of 2"-by-3" stock. In the 2"-by-2" leg, a triangle of ¼" holes is drilled, and in the other leg two 1¼"-by-¼" slots are cut 1½" apart, as seen in the picture. This iron bracket is attached to the telescope tube so that the camera, when mounted, will be squarely over the eyepiece.

The photoflash unit arm is held to the bracket with 1" bolts, the slots allowing it to be moved sideways. The camera is then fixed to the arm with a tripod-screw attachment.

This unit is best suited to a single-lens reflex camera. When taking pictures the lens is set at infinity; the eyepiece is focused sharply by eye; then the camera is eased down until the image is clearest.

GEORGE D. CARTER, JR.
1885 Glendale Dr.
Denver 15, Colo.



George Carter's arrangement for supporting a particular camera. Adjustment for different cameras would be provided by slots in the angle iron where it is fastened to the telescope tube (lower right in the picture).

NEW WOOD TRIPOD

BEN ROSENBERG
101 WEST 45th ST. NEW YORK 36, N. Y.

HIGH-QUALITY
ALUMINIZING

Our minimal thickness coatings are guaranteed not to alter the figure of your mirror, and are guaranteed not to peel or blister. No charge for removal of old coatings. Only the purest aluminum is used for maximum reflectivity. This same high-quality coating will be applied free to your diagonal, if it is shipped with mirror.

3½" — $2.50	8" — $5.00
4¼" — $3.00	10" — $7.50
6" — $4.00	12½" — $10.50

All prices f.o.b. New York. Send for our FREE list of telescope making supplies.

C. HEIN
Box 14, Wakefield Station, New York 66, N. Y.

FIRST STEP TO FINER PERFORMANCE!

Easy, Low-Cost Way To Enjoy Thrills Of Astrophotography

Convert your present 35-mm. camera with this new adapter

Developed by Criterion especially for single-reflex 35-mm. cameras with interchangeable lens. Simply remove present lens mount and add this precision-made achromatic lens that adapts camera to eyepiece of any telescope — refractor or reflector. When ordering, *be sure to specify make of your camera.*

For 35-mm. cameras with SCREW-TYPE lens mount
Cat. #CPA-36 for 1¼" eyepiece holders..**$19.00**
Cat. #CPA-37 for 4" standard Dynascopes **$17.50**

For 35-mm. cameras with BAYONET-TYPE lens mount
Cat. #BPA-36 for 1¼" eyepiece holders..**$22.50**
Cat. #BPA-37 for 4" standard Dynascopes **$21.00**

New!

Dyn-o-Astro 35-mm. Camera

at less than regular price of camera alone!

No need to be an expert photographer to take exciting astrophotos with this *single-reflex*-type, precision-made 35-mm. camera. No complicated settings. No plates or filmholders to load and unload. No worries about missing target. Large focusing screen shows you *exactly* what you are shooting, right up to moment you take the picture. Includes such advanced features as:

- — No accidental double exposures. Winding knob automatically advances film, positions mirror, winds shutter, counts exposures.
- — Dual-speed setting knob controls both fast and slow shutter speeds. Color coding makes mistakes impossible.
- — Integrated magnifier for critical focusing even on dim objects.

Takes time exposures and also has speeds up to 1/500 second. Guaranteed for 2 years. Complete, ready for use.

Model CP-35 fits 1¼" eyepiece holder **$89.00 postpaid**

Model CP-36 fits all standard 4" Dynascopes **$85.00 postpaid**

Dyn-o-Swing Camera Support

fits any 4" or 6" telescope without drilling!

Takes guesswork out of astrophotography by providing *precise centering* of camera over eyepiece. Smooth rack-and-pinion adjustment. Easily attached or detached from tube, with rigid clamp for maximum stability. Special arm lets you swing camera away from eyepiece for visual observation, with instant return for photographing. Can be used with almost any camera. Postpaid.

Cat. #CS-4 to fit all 4" Dynascopes..... **$17.50**
Cat. #CS-47 to fit all 4¼" O.D. tubes.. **$18.95**
Cat. #CS-67 to fit custom 6" Dynascope and 7¼" O.D. tubes..... **$19.95**

Catalog F, describing other accessories and parts, cheerfully sent on request. Satisfaction guaranteed, or money refunded. All items sent postpaid. We pay all postage costs. No shipping, crating or packing charges. Send check, cash, or money order for immediate delivery.

Paraboloidal Mirrors

The most important part of a reflector telescope is the precisely figured mirror. A mirror with a spherical surface suffers from spherical aberration, so it must be altered to a paraboloid to focus all the light rays in each bundle to the same point. Considerable skill is required to parabolize a fine mirror properly. Criterion Custom mirrors are made of the best pyrex glass, selected for freedom from internal stress and strain, and of the correct thickness for each size, parabolized by craftsmen and tested by Roochi and Foucault tests, as well as by diffraction rings and resolution of double stars. They are aluminized and overlaid with zircon quartz. Each is guaranteed unconditionally, and to perform to the limit of resolution for its size.

4" pyrex, f.l. approx. 40"	**$31.00**
6" pyrex, f.l. approx. 54"	**$45.00**
8" pyrex, f.l. approx. 64"	**$89.00**
10" pyrex, f.l. approx. 80"	**$179.00**
12" pyrex, f.l. approx. 96"	**$275.00**

A tolerance of 5% in focal length is customary. A deposit of ⅓ is required with orders for 8" to 12" mirrors.

Rack-and-Pinion Eyepiece Mount

The most mechanically perfect focusing is by rack and pinion. This mount takes standard 1¼" eyepieces. Full 3½" of travel — more than ever before. Accommodates almost any type of eyepiece — positive and negative. Two knurled focusing knobs, variably tensioned and positioned. Solid cast-metal sliding brass tube — close tolerance prevents looseness. Mount aligns itself to any type tube. Four mounting holes, nuts and bolts included. Eye mount has square-rod-type diagonal holder which prevents loose alignment and vibration. Rod tempered to minimize temperature changes. Adjustable for 3" to 8" scopes, also 12" scopes if so specified at no extra cost. Order one or more of the complete eyepieces described below at the same time you send for this rack-and-pinion device, which accommodates any of our eyepieces perfectly.

Cat. #SU-38 **$7.95 postpaid**

New Model Eyepiece Mount

Same features as above but has wider base that is contoured to match the curve of a 7" to 8" diameter tube. Makes professional appearance. Furnished without Diagonal Rod #SU-9R **$9.95**

Diagonal Rod — Cat. #SU-9R..... **$1.00**

Reflecting Telescope Mirror Mounts

Mounting the mirror to your scope correctly is most important. Criterion mounts are especially well designed, and are made of cast aluminum with brass mounting and adjustment screws. One section fits tube, other section holds mirror. Alignment accomplished by three spring-loaded knurled adjusting nuts. Outer cell designed to fit into or over your tube. Sufficient space left between the two cells. All drilled and tapped. Complete with holding clamps, springs, nuts, etc. Ready for use. Will prevent vibration and hold alignment once set. Will hold mirror without distortion of surface figure.

3"	**$3.00**	6"	**$6.00**
4"	**3.50**	8"	**12.50**
5"	**4.00**	10"	**14.75**

Complete Eyepieces

Finest quality. They are precision machined, mounted in standard 1¼" outside diameter barrels; ⅞" O.D. also available at no extra cost. Can be taken apart for cleaning. Designed to give sharp flat field clear to edge.

Huygens 18-mm. f.l. (¾")	**$ 7.50**
Kellner 9-mm. f.l. (⅜")	**7.90**
Kellner 7-mm. f.l. (9/32")	**8.50**
Kellner 12.7-mm. f.l. (½")	**9.50**
Orthoscopic 6-mm. f.l. (¼")	**12.50**
Orthoscopic 4-mm. f.l. (5/32")	**14.50**

Revolving Turret

The Criterion Revolving Turret holds three eyepieces so that, as desired, the power of the telescope can be changed by merely turning the turret to a different ocular. Click stop insures positive and accurate positioning of each eyepiece. Turret holds eyepieces of standard 1¼" outside diameter. Fits into the holder of any refractor or reflector telescope that uses 1¼" eyepieces. Requires no alteration or adjustment and can be attached as easily as putting eyepiece into scope. Made of brass and aluminum with polished chrome-plated barrels.

Cat. #SRT-350..... **$14.50**

Custom Achromatic Objectives and Sunshade

Criterion Custom Series Achromatic Objectives are of the highest precision quality, tested and approved and acclaimed to be superb in every respect. Equipped with sunshade and dewcap.

FEATURES

- Perfect color correction at the wave lengths of the C and F spectral lines, best for visual observing. Scientifically designed, corrected by master craftsmen.
- Delicately hand-corrected to fullest minimization of residuals, and to eliminate spherical aberration.
- Air-spaced to very critical optical formula.
- Magnesium-fluoride coated on all surfaces for maximum light transmission.
- Cell engraved with effective focal length and completely threaded.
- Sunshade extension with dewcap on all sizes.

Cat. #	Clear Aperture	Focal Length	Price
S-210	60 mm. (2.4")	910 mm. (35.8")	**$19.95**
S-220	3"	1250 mm. (49.2")	**52.00**
S-230	4"	1600 mm. (62.9")	**95.00**

Four-Vane Diagonal Holders

Criterion 4-vane diagonal mountings are fully adjustable, will hold elliptical diagonals in perfect alignment. Made of brass, chemically blackened. Precision adjusting screws center flat and vary its angle so that primary and secondary mirrors can be set in perfect alignment. Thin vanes with special adjustable studs.

Cat. #	Minor-Axis Size	For Tubes	Price
S-51	1.25"	6½" to 7½"	**$10.00**
S-52	1.30"	6½" to 7½"	**10.00**
S-53	1.50"	8½" to 9½"	**10.00**
S-54	1.75"	9½" to 10½"	**12.50**
S-55	2.00"	11" to 11½"	**14.95**
S-56	2.50"	Specify tube I.D.	**19.95**

CRITERION MANUFACTURING CO.

Manufacturers of Quality Optical Instruments

Dept. STP-18, 331 Church St., Hartford 1, Conn.

ITEMS ON THIS PAGE ARE ESPECIALLY SUITABLE FOR CHRISTMAS GIFTS!

3-inch Astronomical Reflector

60 to 180 Power

An Unusual Buy!

Assembled — ready to use! See Saturn's rings, the planet Mars, huge craters on the moon, star clusters, moons of Jupiter, double stars, nebulae, and galaxies! Equatorial-type mounting with locks on both axes. Aluminized and over-coated 3"-diameter f/10 primary mirror, ventilated cell. Telescope comes equipped with a 60x eyepiece and a mounted Barlow lens, giving you 60 to 180 power. A finder telescope, always so essential, included. Sturdy, hardwood, portable tripod.

FREE with Scope: Valuable **STAR CHART** plus 272-page **"HANDBOOK OF THE HEAVENS"** plus the book **"HOW TO USE YOUR TELESCOPE."**

Stock #85,050-Y.....$29.95 ppd.

See the Stars, Moon, Planets Closeup!

4¼" ASTRONOMICAL TELESCOPE

REFLECTOR TYPE — Up to 270 Power!

With this scope you can see the craters on the moon, the rings of Saturn, double stars. Mirror guaranteed to give theoretical limit of resolution. Rack-and-pinion focusing, removable mirror mount, counterweight. Real equatorial mounting — only one adjustment follows stars! Aluminum tube. 6-power finder telescope. 2 standard-size eyepieces and mounted Barlow lens give you powers of 40x, 90x, 120x, 270x. Mirror f/11, and corrected to better than ¼ wave length. FREE with Scope: — Valuable **STAR CHART** plus 272-page **"HANDBOOK OF THE HEAVENS"** plus **"HOW TO USE YOUR TELESCOPE"** book.

Stock #85,006-Y.....$74.50 f.o.b. Barrington, N. J.

NEW! STATIC ELECTRICITY GENERATOR

Educational and Lots of Fun!

See a thrilling spark display as you set off a miniature bolt of lightning. Absolutely safe and harmless — perfect for classroom experimentation — ideal for science clubs. Sturdily made — stands 14" high. Turn the handle and two 9" plastic disks rotate in opposite directions. Metal collector brushes pick up the static electricity, store it in the Leyden-jar-type condenser until discharged by the jumping spark. Countless tricks and experiments. 24-page instruction booklet included.

Stock #70,070-Y.....$12.95 ppd.

50-, 150-, 300-POWER MICROSCOPE

Amazing value — equal of a $75.00 instrument! 3 achromatic objective lenses on revolving turret! Imported. The color-corrected, cemented achromatic lenses in the objectives give you far superior results to the single lenses found in other microscopes selling in this range. Results are worth the difference! Fine rack-and-pinion focusing.

Stock #70,068-Y.....$14.95 ppd.

MOUNTED 500-POWER OBJECTIVE

. . . Threaded for easy attachment on microscope above. Achromatic lenses for fine viewing. 3-mm. focal length.

Stock #30,197-Y.....$5.00 ppd.

NEW BINOCULAR-TO-CAMERA HOLDER

For Exciting Telephoto Shots

Will fit any camera.

Bring distant objects 7 times nearer with a 35-mm. camera, 7 x 50 binocular, and our NEW BINOCULAR-TO-CAMERA HOLDER. Ideal for photographing the constellations, star clusters, the moon, as well as cloud formations, wild life, vistas. Camera and binocular attach easily. Use any binocular or monocular — any camera, still or movie. Take color or black-and-white. Attractive gray crinkle and bright chrome finish, 10" long. Full directions for making telephotos included.

Stock #70,223-Y.....$11.50 ppd.

Take Pictures Through Your Telescope with the EDMUND CAMERA HOLDER for TELESCOPES

Bracket attaches permanently to your reflecting or refracting telescope. Removable rod with adjustable bracket holds your camera over scope's eyepiece and you're ready to take exciting pictures of the moon. You can also take terrestrial telephoto shots of distant objects. Opens up new fields of picture taking!

SUN PROJECTION SCREEN INCLUDED

White metal screen is easily attached to holder and placed behind eyepiece. Point scope at sun, move screen to focus . . . and you can see sunspots!

All for the low, low price of $9.95

Includes brackets, 28⅜" rod, projection screen, screws, and directions. Aluminum . . . brackets black crinkle painted.

Stock #70,162-Y.....$9.95 ppd.

NOTICE!

EDMUND IS NOW HEADQUARTERS FOR MATH LEARNING AND TEACHING AIDS!

Play This New Game — MATH MAGIC . . . The Fun-Way to Math Skill!

Educator-approved! 3 fascinating games in one! Great fun for the whole family. Increases skill at addition, subtraction, multiplication, division. Includes dial and spinner, numbered cards, plastic tokens, etc. — also rules and directions.

Stock #70,204-Y.....$3.00 ppd.

D-STIX Construction Kits

Shape Your Ideas Fast!

Scientists, executives, planners, designers — all find D-STIX simplify visualization and demonstration of complex ideas. D-STIX — colored wood sticks ⅛" thick and "easy-on" rubber joints approx. 3/16" diam. — help you work out molecular structures, geometric figures, structural members, shapes, models of scientific apparatus. Used by professional men, educators, hobbyists. Big basic kit contains 230 or 370 pieces. Additional parts in supplementary kit. Far superior to other kits of wood or metal. Money-back guarantee.

Stock #70,209-Y....(230 pcs.).....$3.00 ppd.
Stock #70,210-Y....(370 pcs.).....$5.00 ppd.
Stock #70,211-Y....(452 pcs.).....$7.00 ppd.

Ideal for Beginners . . .

STARMASTER ASTRONOMY SET!

Bring the thrills and wonders of outer space right into your home with this fascinating, educational Astronomy Set. De Luxe, 37-piece Set includes separate film projector and action pictures of a trip to the moon, from blast-off to return. Exciting views of the moon, planets, comets, distant galaxies, sun explosions, star clusters, other space wonders.

Also star projector and constellation projector (each approx. 4½" high). Projectors show many great constellations and stars, outline forms of Big Bear, Sagittarius, Southern Cross and others, for easy identification.

Also includes 10x telescope (single-element lens, meniscus, nonachromatic) with 1" aperture, suitable for viewing the moon.

Power source, flashlight pointer, and instructions included. Lower-priced Junior Set with single projector also available, but without telescope.

Stock #70,234-Y....De Luxe.....$10.00 ppd.
Stock #70,233-Y....Junior.....$ 5.00 ppd.

TELESCOPE ROLL-FILM CAMERA

This model uses rolls of #127 film. Picture area will be a circle 1-9/16" in diameter.

The advantage of this model is the ease of using roll film. With each camera you get a piece of ground glass. Before loading film in the camera, you focus the telescope. Then lock it in this position. For positions other than infinity, you can scribe a mark on your tube.

Stock #70,182-Y.....$29.50 ppd.

SHEET-FILM CAMERA

Uses sheet film 2¼" x 3¼" size. Camera box size is 3" x 4" x 5".

Stock #70,166-Y.....$39.50 ppd.

ASTRO COMPASS and STAR FINDER

Gov't. Cost $75 — Price $14.95 ppd.

Determines positions of stars quickly. Shows various celestial co-ordinates. An extremely useful star finder which can be rotated through 60° angles along calibrated degree scale. Has single eye lens with viewing stop, two spirit levels for aligning, tangent screw with scale for fine precision readings, azimuth scale graduated in two-degree intervals, adjustable tilting azimuth scale for angle reference of stars on distant objects. War surplus. Gov't. cost $75.00. Instructions, carrying case included.

Stock #70,200-Y.....Only $14.95 ppd.

"EASY-CARRY" SLIDE PROJECTOR

Now you can show your favorite slides ANYWHERE to friends, classes, groups, without lugging cumbersome equipment. "EASY-CARRY" PROJECTOR folds to 2½" wide, 4½" high, 4½" long — opens to 8" long. Weighs less than 3 lbs. SHOWS and PROJECTS 2-x-2 35-mm. slides or 35-mm. filmstrips, single or double frame. 100 watt, 120 volt. 3-element f/3.5 lens with 80-mm. focal length. 6' cord and plastic leatherette carrying case included. Ideal for travel or occasional use.

Stock #70,232-Y.....$22.95 ppd.

Terrific Buy! American Made!

OPAQUE PROJECTOR

Projects illustrations up to 3" x 3½" and enlarges them to 4 ft. wide. No film or negatives needed. Projects charts, diagrams, pictures, photos, lettering in full color or black-and-white. Operates on 115-volt a.c. current. 6' extension cord and plug included. (Operates on 60-watt bulb, not included.) Size 12" x 8" x 4½" wide. Weighs 1 lb., 2 oz. Plastic case with built-in handle.

Stock #70,199-Y.....$7.95 ppd.

EDMUND SCIENTIFIC CO.

SALE! TERRIFIC WAR-SURPLUS BARGAINS!

AERIAL CAMERA LENSES

Big variety . . . at a fraction of Gov't. cost! f/6, 24" f.l., with diaphragm and lens cone. Used. Weight 25 lbs.
Stock #85,059-Y..$39.50 f.o.b. Utah

Same as above, but new. Weight 25 lbs.
Stock #85,060-Y..$59.50 f.o.b. Utah

f/8, 40" f.l., no mount or shutter. Weight 6⅛ lbs.
Stock #70,186-Y.....$49.50 ppd.

f/3.6, 20" f.l., telephoto with shutter and diaphragm. Weight 6⅛ lbs.
Stock #70,187-Y.....$65.00 ppd.

f/4.5, 6⅜" f.l., with shutter and diaphragm. Weight 1 lb., 6 ozs.
Stock #70,189-Y.....$24.50 ppd.

These lenses are being successfully used for wide-aperture Moonwatch telescopes to see the small and fainter satellites. For eyepiece use our GIANT ERFLE.

DOUBLE AND TRIPLE YOUR TELESCOPE'S POWER WITH A BARLOW LENS



WHAT IS A BARLOW? A Barlow lens is a negative lens used to increase the power of a telescope without resorting to short focal length eyepieces, and without the need for long, cumbersome telescope tubes. Referring to the diagram above, a Barlow is placed the distance P inside the primary focus of the mirror or objective. The Barlow diverges the beam to a distance Q. This focus is observed with the eyepiece in the usual manner. Thus, a Barlow may be mounted in the same tube that holds the eyepiece, making it very easy to achieve the extra power. The new power of the telescope is not, as you might suppose, due to the extra focal length given the objective by the difference between P and Q. It is defined as the original power of the telescope times the quotient of P divided into Q.

Beautiful chrome mount. We now have our Barlow lens mounted in chrome-plated brass tubing with special spacer rings that enable you easily to vary the power by sliding split rings out one end and placing them in other end. Comes to you ready to use. Just slide our mounted lens into your 1¼" I.D. tubing, then slide your 1¼" O.D. eyepieces into our chrome-plated tubing. Two pieces provided, one for regular focal length eyepieces and one for short focal length ones.
Stock #30,200-Y Mounted Barlow lens.....$8.00 ppd.

UNMOUNTED 3X BARLOW LENS

These lenses are made for telescopes that have smaller diameter eyepieces than the standard 1¼" size. Mount one between the eyepiece and objective, and triple your power. Instructions included. Single-element lens, focal length −1-5/16", unmounted.
Stock #30,185-Y....0.932" diam.....$3.50 ppd.
Stock #30,328-Y....0.912" diam.....$2.50 ppd.

3X ADJUSTABLE-DIAMETER BARLOW LENS

For telescopes with eyepieces smaller than the standard 1¼" outer-diameter size. Prongs on mount can be opened or closed to fit tubes from 13/16" to 1" outer diameter. Directions for using included.
Stock #30,339-Y....$5.00 ppd.

WAR-SURPLUS TELESCOPE EYEPIECE

Mounted Kellner Eyepiece, Type 3. Two achromats, focal length 28 mm., eye relief 22 mm. An extension added. O.D. 1¼", standard for most types of telescopes. Gov't. cost $26.50.

Stock #5223-Y.....$7.95 ppd.

SPECIAL SALE OF OPTICAL PERISCOPES

We made a lucky buy so here is another famous Edmund war-surplus bargain. It is a $500.00 tank periscope with over $200.00 worth of optics in it for only $18.00 postpaid. At one spot, you look through and see up and out of the prisms at unit power; or you can look through the built-in telescope system and see a wide-angle view at 6x. Simply remove the top prism and you have an 11°.3 6-power satellite telescope. *Brand new in original packages.* Over-all size is 14" long by 7" wide by 2½" deep.

Now just for the bit of work involved in disassembling, you get a treasure-trove of precision optical parts for only $18.00. Glance at this list of parts and you'll realize what a value we are offering. You get a wide-angle Erfle eyepiece with a full 68° field and 1¼" focal length. This is the same eyepiece used in Moonwatch telescopes, our Stock No. 5160. Also included are a 38-mm.-diam. collimator lens, 131-mm. focal length; 48-mm.-diam., mounted, coated objective lens, 189-mm. f.l.; prism cluster containing two light flint glass right-angle prisms (A, 41 mm.; B, 91 mm.; C, 64 mm.; and A, 41 mm.; B, 57 mm.; C, 40 mm.); two mounted right-angle prisms (A, 23 mm.; B, 34 mm.; C, 25 mm.; and A, 40 mm.; B, 42 mm.; C, 30 mm.); two silvered tank prisms (A, 102 mm.; B, 53 mm.; C, 38 mm.; and A, 167 mm.; B, 69 mm.; C, 49 mm.). (*See our catalog for explanation of these prism sizes.*) Also some other optical elements and reticles. Every amateur will find this a real value. Don't wait — order today. The last time we had these they sold for $30.00 — now on sale at only $18.00.

Stock #70,227-Y.....$18.00 ppd.

EQUATORIAL MOUNT and TRIPOD with CLOCK DRIVE

Heavy-duty mount. Drive operates on 110-volt, 60-cycle, a.c. house current. Follows motion of stars smoothly. 32" tripod legs included.
Stock #85,081-Y.....$76.50 f.o.b. Barrington, N. J.

Same mount as above, without clock drive, for 8" or smaller reflectors and for 4" or smaller refractors.
Stock #85,023-Y.....New Low Price.....$39.50 f.o.b. Barrington, N. J.

8-POWER ELBOW TELESCOPE

War Surplus — Amazing Buy!
$200 Gov't. Cost—Only $13.50

Big 2" objective, focusing eyepiece 28-mm. focal length, Amici erecting system, turret-mounted filters of clear, red, amber, and neutral, reticle illumination. Sparkling, clear, bright image — 6° field (325 ft. at 1,000 yards). Focus adjusts 15 ft. to infinity. Eyepiece alone, 28-mm. focal length, is worth more than $12.50.
Stock #70,173-Y.....$13.50 ppd.

GIANT MAGNETS! TERRIFIC BARGAINS!



War Surplus — Alnico V type. Horseshoe shape. Tremendous lifting power. 5 lb. size. Its dimensions: A — 3¼"; B — 2¾"; C — 4-3/16"; D — 1⅛"; E — 1¼"; F — 2⅝". Strength is about 2,000 gauss. Will lift over 125 lbs.
Stock #70,183-Y.....$8.50 ppd.

15¾-lb. size magnet. Approximately 5,000-6,000-gauss rating. Will lift over 250 lbs.
Stock #85,088-Y.....$22.50 f.o.b. Barrington, N. J.
Shipping wt. 22 lbs.

Mounted Ramsden Eyepieces

Standard 1¼" Diameter

Our economy model, standard-size (1¼" O.D.) eyepiece. We mounted two excellent quality plano-convex lenses in black anodized aluminum barrels instead of chrome-plated brass to save you money. The clear image you get with these will surprise you. Directions for using short focal length eyepieces are included with both the ⅛" and ½" models.
Stock #30,204-Y....⅛" focal length....$4.75 ppd.
Stock #30,203-Y....½" focal length....$4.50 ppd.

"MAKE-YOUR-OWN" 4¼" MIRROR KIT

The same fine mirror as used in our telescopes, polished and aluminized, lenses for eyepieces, and diagonal. No metal parts.
Stock #50,074-Y.....$16.25 ppd.

GIANT ERFLE EYEPIECE

Here is an exciting bargain. We have obtained a large lot of these eyepieces reasonably — so down goes the price to $9.95 for a real sale. Lens system contains 3 coated achromats over 2" in diameter. Gov't. cost over $100.00. Brand new, weight 2 pounds. The value will double when this lot is all sold, and triple and quadruple as years pass. If we didn't need to reduce our inventory, we'd be tempted to hold onto these eyepieces. Their wide apparent field is 65°. The focal length is 1½". Lenses are in a metal cell with spiral threads; focusing adapter with 32 threads per inch is included; diameter is 2-11/16". If you don't order now and you miss out on a hundred-dollar eyepiece for only $9.95, you can't say that we didn't try to impress you with its value. You can make some super-duper finders with these eyepieces. They are also ideal for rich-field telescopes, which are becoming more popular daily, particularly in the Sputnik age.
Stock #50,178-Y.....$9.95 ppd.

8" SETTING-CIRCLE SET

Stock #50,133-Y....Complete set.....$3.00 ppd.
Stock #60,078-Y....360° declination circle only $1.60 ppd.
Stock #60,079-Y....24-hour right-ascension circle only $1.60 ppd.

5¼" SETTING-CIRCLE SET

Stock #50,190-Y....Complete set.....$2.50 ppd.
Stock #60,080-Y....360° declination circle only $1.35 ppd.
Stock #60,081-Y....24-hour right-ascension circle only $1.35 ppd.

6X FINDER TELESCOPE

Has crosshairs for exact locating. You focus by sliding objective mount in and out. Base fits any diameter tube — an important advantage. Has 3 centering screws for aligning with main telescope. 20-mm.-diameter objective. Weighs less than ½ pound.
Stock #50,121-Y.....$8.00 ppd.

BE SURE TO GET FREE CATALOG "Y"

128 Pages! Over 1000 Bargains!

Fantastic variety — never before have so many lenses, prisms, optical instruments, and components been offered from one source. Positively the greatest assembly of bargains in all America. Imported! War Surplus! Hundreds of other hard-to-get optical items. Many science and math learning and teaching aids. Write for Free Catalog "Y."

ORDER BY STOCK NUMBER . . . SEND CHECK OR MONEY ORDER . . . SATISFACTION GUARANTEED!

BARRINGTON • NEW JERSEY

Christmas . . .
VERNONscope

Make this an extra-special Christmas with a highest-quality De Luxe VERNONscope — these professional-type reflecting telescopes are now selling for amazingly low prices.

The 4¼" De Luxe VERNONscope

- ★ 4¼" parabolic mirror, f/10, ground and polished to better than ⅛ wave, aluminized and over-coated with quartz for maximum optical protection, guaranteed to perform to Dawes' astronomical limit of resolution.
- ★ NEW, rugged PAR Jr. Equatorial Mount, designed to give utmost rigidity at the points of greatest stress (note tapered aluminum housings). Includes latitude adjustment, large knurled locking screws and close-tolerance machined bearings with ¾" chrome-plated precision-ground steel shafting. Base of mount is fully rotatable for altazimuth viewing and can be quickly disassembled from the tripod. Steel legs fold for easy portability. Mount has black primer and 3-dimensional space blue marbleized finish.
- ★ Seamless aluminum tube, blackened inside, with 3-D space blue finish.
- ★ Finest 14-mm. wide-angle Kellner eyepiece and Barlow lens mounted in black-anodized tubing, giving magnifications of 80x and 160x.
- ★ New achromatic-type 4x finder with adjustable mounting bracket.
- ★ Smooth, precision helical focusing unit permits micrometer accuracy to give sharpest images possible.
- ★ Machined aluminum end caps and adjustable mirror mounting.
- ★ Covers for all optical components.
- ★ 44-page telescope operation manual included.

4¼" VERNONscope $69.50
Shpg. wt., 25 lbs. f.o.b. Candor, N. Y.

The 3" De Luxe VERNONscope

- ★ 3" parabolic mirror, f/8, pyrex, aluminized and over-coated with quartz, corrected to better than ⅛ wave.
- ★ 14-mm. wide-angle Kellner eyepiece and Barlow lens give magnifications of 40x and 80x.
- ★ Same features and components as in 4¼".

3" VERNONscope $49.50
Shpg. wt., 16 lbs. f.o.b. Candor, N. Y.

Telescopes are completely assembled, thoroughly tested, and correctly aligned before leaving the factory. They are carefully packaged to insure safe shipment.

We unconditionally guarantee each VERNONscope to be exactly as described and illustrated, and to surpass even your greatest expectations, or return it within one month and your money will be immediately refunded. VERNONscope & Co. is the sole manufacturer of the 100%-American-made VERNONscope, selling to our customers direct from the factory. Send your order today for immediate delivery. C.O.D. orders accepted with one-half payment.

VERNONscope & Co.
Candor, New York

CELESTIAL CALENDAR

Universal time (UT) is used unless otherwise noted.

OBSERVABLE ALGOL MINIMA

CHECKING whether Algol, the famous eclipsing variable in Perseus, is below normal brightness is a good habit for one who frequently scans the sky. But just what are the chances of finding Beta Persei at or near minimum brightness?

The minima occur every two days, 20 hours, 49 minutes, usually on one date in three. About 10 hours are required for the change from magnitude 2.2 to 3.5 and back again. During the first two and last two hours the loss in light could escape a casual glance, so for only one-quarter of a day would the change attract attention. Thus, Algol is noticeably faint about 1/12 of the time.

But some minima occur in the daytime, halving the chances even in winter, to about 1/24. Considering cloudy nights, the expectation of casually finding Algol faint is, therefore, very poor, and predictions of minima are practically essential to the observer.

As the listing here shows, the most convenient times this month for American amateurs to see Algol dimmed will be during the minima predicted for the 9th at 2:38 a.m. Eastern standard time; the 11th, at 11:27 p.m.; and the 14th, at 8:16 p.m.

NOVEMBER METEORS

Beginning in late October, the Taurid meteor shower reaches its maximum during the first week of November. On the 6th a single observer may see up to 15 meteors per hour, the radiant then being located in western Taurus. The moon, only a week past new phase, should not interfere seriously with observations of these meteors.

For the Leonid shower, with maximum on November 17th, observing conditions are very unfavorable, as it is shortly after full moon. W. H. G.

MINOR PLANET PREDICTIONS

Parthenope, 11, 9.6. November 28, 6:03.9 +18.25. December 8, 5:54.6 +18.30; 18, 5:44.0 +18.39; 28, 5:33.4 +18.49. January 7, 5:24.1 +19.02; 17, 5:17.0 +19.17. Opposition on December 18.

After the asteroid's name are its number and the magnitude expected at opposition. At 10-day intervals are given its right ascension and declination (1950.0) for 0ʰ Universal time. In each case the motion of the asteroid is retrograde. Data are supplied by the IAU Minor Planet Center at the University of Cincinnati Observatory.

MINIMA OF ALGOL

November 3, 14:01; 6, 10:50; 9, 7:38; 12, 4:27; 15, 1:16; 17, 22:05; 20, 18:54; 23, 15:43; 26, 12:32; 29, 9:21.

December 2, 6:10; 5, 2:59; 7, 23:48.

These minima predictions for Algol are based on the formula in the 1953 *International Supplement* of the Krakow Observatory. The times given are geocentric; they can be compared directly with observed times of least brightness.

UNIVERSAL TIME (UT)

TIMES used in Celestial Calendar are Greenwich civil or Universal time, unless otherwise noted. This is 24-hour time, from midnight to midnight; times greater than 12:00 are p.m. Subtract the following hours to convert to standard times in the United States: EST, 5; CST, 6; MST, 7; PST, 8. If necessary, add 24 hours to the UT before subtracting, in which case the result is your standard time on the day preceding the Greenwich date shown. For example, 6:15 UT on the 15th of the month corresponds to 1:15 a.m. EST on the 15th, and to 10:15 p.m. PST on the 14th.

SKY - GAZERS EXCHANGE

Classified advertising costs 30 cents a word, including address; minimum charge, $4.00 per ad. Only one *for sale* ad per issue for each advertiser. *Remittance must accompany order.* Insertion is guaranteed only on copy received by the 20th of the second month before publication; otherwise, insertion will be made in next issue. We cannot acknowledge classified ad orders. Sky Publishing Corporation assumes no responsibility for statements made in classified ads, nor for the quality of merchandise advertised. Write Ad Dept., *Sky and Telescope,* Harvard Observatory, Cambridge 38, Mass.

FOR SALE: 4½" Wray refractor, all accessories, $450.00. Write John Sanford, 365 First St., Newburgh, N. Y.

METEORITES: Excellent for teaching, research, lectures, and demonstrations. $1.00, $2.00, $12.00, $15.00. Scientific Laboratory, 2846 Oakley Ave., Baltimore 15, Md.

6" PYREX parabolic mirrors, f/5.5 and f/7.5, $40.00 each. Heyman Optics, 19 Priscilla, New Bedford, Mass.

SIDEREAL DRIVE for telescope, track stars accurately, simple gearless design, plans, $1.00. "Space Chart," planets, sun, moon, asteroids, constellation maps, $1.00. L. Mussgnug, Box 74, Bethel, Conn.

FOR SALE: 8" reflecting telescope, with equatorial mounting, $285.00. Morgan Grace, Lawrenceville School, Lawrenceville, N. J.

FIBERGLASS TUBES: Ready to mount. White glass-smooth surface, lightweight, great strength, low thermal conductivity, blackened inside, polished end rings. W. R. Parks, 20942 S. LaSalle, Torrance, Calif.

ALUMINUM TUBING: 17 sizes, 1" through 8", Pesco-A, Box 363, Ann Arbor, Mich.

SOUTHWEST OBSERVERS: Finest refractor telescopes — 2.4", 3", and 4" altazimuth and equatorial models shipped from Dallas, Texas. Terms and details on request. Melton Industries, 1901 Levee St., Dallas 7, Tex. Phone: RI 8-4769.

FOR SALE: 6" f/9 altazimuth reflector, $135.00. Also aluminized Pyrex mirrors: 4¼" f/10, $20.00; 6" f/8, $40.00. Donald O'Toole, 2100 Eye St., Apt. 5, Sacramento, Calif.

COLOR SLIDES depicting other planets, paintings, space models. Catalogue and sample, 25¢. Morris Dollens, 4372 Coolidge Ave., Los Angeles 66, Calif.

TELESCOPES, aluminum tubing, eyepieces. Write Peninsula Scientific, 2421 El Camino Real, Palo Alto, Calif.

MIRRORS: If you are interested in buying or selling a used telescope mirror, consult Robert Wall, 1520 Mimosa Ave., Charlotte 5, N. C.

QUESTAR, de luxe, like new, practically unused, $825.00. J. E. Gifford, 404 E. Howell, Seattle 22, Wash.

FOR SALE: Refractor telescope. Good Sears model. Now $50.00. Lee Moore, 1823 Buckland Ave., Fremont, Ohio.

INTERESTED in astronomy as a career? *Vocational and Professional Monographs: Astronomy,* by Dr. Freeman D. Miller, describes personal qualifications, scholastic training, and job opportunities. $1.00 postpaid. Send to Box B, *Sky and Telescope,* Harvard Observatory, Cambridge 38, Mass.

CLOSEOUT SALE: "Satellite Pathfinder," an ingenious device designed at the American Museum-Hayden Planetarium to help predict Northern Hemisphere passages of artificial satellites. $1.00 postpaid. Send to Box C, *Sky and Telescope,* Harvard Observatory, Cambridge 38, Mass.



THE SUN, MOON, AND PLANETS THIS MONTH

The sun, on the ecliptic, is shown for the beginning and end of the month. The moon's symbols give its phase roughly, with the date marked alongside. Each planet is located for the middle of the month or for other dates shown. All positions are for 0^h Universal time on the respective dates.

Mercury reaches greatest eastern elongation on November 3rd, 23° 33′ from the sun, and may be seen very low in the southwest for about a week before and after this date. The planet is poorly placed for observation at this elongation, however, and sets about an hour after the sun. Inferior conjunction is on the 24th.

Venus attains greatest western elongation, 46° 37′ from the sun, on November 12th. It is a brilliant object, magnitude −4.0, rising about four hours before sunup. On the 11th its disk will be half illuminated and 24″.7 in diameter. The moon will be very near Venus on the morning of the 27th, conjunction occurring at 2:10 Universal time, with the planet 37′ north if viewed from the center of the earth. An occultation will be seen from South Africa and Australasia.

Mars is in the morning sky, but too near the sun to be observed this month.

Jupiter may be seen the first week in November, low in the southwest. On the 1st it is magnitude −1.3, setting about 1½ hours after the sun. Mercury and Jupiter will be in conjunction on the 7th, with the former about 3½° south.

Saturn, of the 1st magnitude, is in Sagittarius, low in the southwest during the early evening. A telescope will reveal its disk, 13″.8 in polar diameter, and the rings, 34″.8 in extent. The moon will pass 4½° north of Saturn on the 4th.

Uranus, in western Leo, rises about midnight local time. Binoculars will reveal this 6th-magnitude object, at right ascension 9^h 33^m.9, declination +15° 12′ (1950 co-ordinates) on November 14th, the time of its western quadrature. Uranus is stationary in right ascension on the 27th, and then begins retrograde (westward) motion among the stars.

Neptune is too close to the sun to be seen this month. W. H. G.

MOON PHASES AND DISTANCE

First quarter	November 7,	13:23
Full moon	November 15,	9:42
Last quarter	November 23,	13:03
New moon	November 30,	8:46
First quarter	December 7,	2:11

November	Distance	Diameter
Perigee 2, 1^h	223,100 mi.	33′ 17″
Apogee 17, 7^h	252,600 mi.	29′ 24″
Perigee 30, 12^h	221,600 mi.	33′ 30″
December		
Apogee 14, 7^h	252,600 mi.	29′ 24″

VARIABLE STAR MAXIMA

November 2, S Carinae, 100661, 5.7; 4, T Normae, 153654, 7.4; 7, RS Librae, 151822, 7.7; 7, RS Cygni, 200938, 7.4; 19, R Sagittarii, 191019, 7.2; 21, T Herculis, 180531, 8.0; 22, R Cygni, 193449, 7.3; 27, T Aquarii, 204405, 7.9.

December 1, R Draconis, 163266, 7.6; 5, T Centauri, 133633, 6.1.

These predictions of variable star maxima are by the AAVSO. Only stars are included whose mean maximum magnitudes are brighter than magnitude 8.0. Some, but not all of them, are nearly as bright as maximum two or three weeks before and after the dates for maximum. The data given include, in order, the day of the month near which the maximum should occur, the star name, the star designation number, which gives the rough right ascension (first four figures) and declination (bold face if southern), and the predicted magnitude.

ONE QUALITY–1/20 WAVE

The finest optics, as well as mediocre or poor ones, never wear out, but each kind tells you its story *every minute* you observe with it over the years ahead.

1/20–WAVE PARABOLOIDAL MIRRORS

The new mirror price includes aluminizing, as well as a 1/10-wave diagonal mirror of the size listed. Add 10% for nonstandard focal ratios from f/5 to f/12.

Mirror Size	*Flat Size*	*Price*
6-inch f/9	1¼″	$ 95.00
8-inch f/8	1½″	155.00
10-inch f/8 or f/7	1¾″	255.00
12½-inch f/8 or f/7	2″	395.00

Retouching of defective mirrors, f/5 to f/10: 60% of prices above. In sending mirrors for retouching, include also your flat; if defective in figure but otherwise suitable, it will also be corrected, at no extra charge.

Terms: ¼ with order, ¼ when ready to ship, balance up to 6 months on arrangement. If full payment is included with order, a 2% discount may be taken.

Quotations on special curves or materials on request. We also offer telescopes less mountings.

Write for free booklet, *The Reflecting Telescope,* for a practical discussion of why 1/20 wave *is* necessary for the finest results, and also ways for obtaining the most from your reflecting telescope.

frank vaughn Optics

1806 South Park St., Box 2053
Madison 5, Wisconsin

NOW for a mere $55 down— YOU can own this UNITRON

Here is one of the most popular refracting telescopes for the amateur astronomer who wishes to combine both visual observation and astrophotography. Optically excellent, this instrument also has the very precise mechanical adjustments so essential for serious work. And the low 10% down payment is well within the average budget.

Model 145 is supplied complete with equatorial mounting and slow-motion controls, tripod, 2.4" photographic guide telescope with star diagonal and crosshair eyepiece, 10x 42-mm. view finder, 6 eyepieces, choice of UNIHEX Rotary Eyepiece Selector or Star Diagonal and Erecting Prism System, sun projection screen, sunglass, UNIBALANCE tube assembly, cabinets, and so forth, all for only $550. Extra eyepieces are available to permit a total of nine magnifications ranging from 30x to 300x.

If your interest lies only in visual observation, then we suggest UNITRON Model 142, the regular 3" Equatorial, priced at only $435 complete (down payment $43.50).

There is a UNITRON 3" Altazimuth model for as little as $265, complete, and other UNITRON Refractors are priced as low as $75. All may be purchased for only 10% down using our Easy Payment Plan. Whichever model you choose, you are assured of obtaining the finest instrument in its class.

After all, like thousands of others, you can place your confidence in UNITRON.

See the back cover.

3-inch PHOTO-EQUATORIAL

ACCESSORIES to Modernize Your Telescope

Many of the accessories and components pictured on Model 145 are available separately, to fit telescopes of various sizes.

ASTRO-CAMERA 220: complete with accessories	$ 69.50
PHOTOGRAPHIC GUIDE TELESCOPE: with brackets and accessories	75.00
VIEW FINDER: 10x, 42-mm. with mounting brackets	18.00
EQUATORIAL MOUNTING and TRIPOD: with cabinet	198.00
UNIHEX ROTARY EYEPIECE SELECTOR: for rapid change of eyepieces	24.75
UNICLAMP CAMERA BRACKET: for 3" refractor	3.75
OBJECTIVE LENS: 3", coated, air-spaced, in cell	69.00
RACK-AND-PINION MECHANISM: with drawtube	Prices from $12.50

UNITRON INSTRUMENT DIVISION of UNITED SCIENTIFIC CO.
204-206 MILK STREET BOSTON 9, MASSACHUSETTS

Does Santa know you want a UNITRON for Xmas?

UNITRON 6" AND 4" MODELS

UNITRON 6" and 4" Model 166V.

There are UNITRON 6" models priced from $5125 to $6075. Write for Bulletin 600.

HOW TO ORDER A UNITRON

Send check or money order in full or use our Easy Payment Plan, as described below. Shipments made express collect. Send 20% deposit for C.O.D. shipment. UNITRON instruments are fully guaranteed for quality, workmanship, and performance, and must meet with your approval or your money back.

Use Our EASY PAYMENT PLAN

UNITRON's popular Easy Payment Plan is a convenient and economical way to buy your UNITRON Refractor when you do not want to disturb your savings or when you haven't the total cost of the telescope immediately available. The down payment required is 10%. The balance due is payable over a 12-month period, and there is a 6% carrying charge on the unpaid balance. Your first payment is not due until 30 days after you receive the instrument, and if you should want to pay the entire balance due at that time, the carrying charge is canceled. (6" models available on special plan.)

NOW is definitely the time to reserve a UNITRON for that very special person on your Christmas list who perhaps may be none other than yourself. Precision craftsmanship cannot be hurried and the number of UNITRONS which Santa will have available is necessarily limited. To avoid disappointment, order your UNITRON now. Full payment or the down payment may be made at the time you wish to have the telescope delivered. Remember, there is no substitute for a UNITRON.

MANY Models To Choose From!

1.6" ALTAZIMUTH ($7.50 Down) with eyepieces for 78x, 56x, 39x — **$75**

2" SATELLITE ($7.50 Down) 6x, diagonal eyepiece, altazimuth mount with circles, stand — **$75**

2.4" ALTAZIMUTH ($12.50 Down) with eyepieces for 100x, 72x, 50x, 35x — **$125**

2.4" EQUATORIAL ($22.50 Down) with eyepieces for 129x, 100x, 72x, 50x, 35x — **$225**

3" ALTAZIMUTH ($26.50 Down) with eyepieces for 171x, 131x, 96x, 67x, 48x — **$265**

3" EQUATORIAL ($43.50 Down) with eyepieces for 200x, 131x, 96x, 67x, 48x — **$435**

3" PHOTO-EQUATORIAL ($55.00 Down) with eyepieces for 200x, 171x, 131x, 96x, 67x, 48x — **$550**

4" ALTAZIMUTH ($46.50 Down) with eyepieces for 250x, 214x, 167x, 120x, 83x, 60x — **$465**

4" EQUATORIAL ($78.50 Down) with eyepieces for 250x, 214x, 167x, 120x, 83x, 60x, 38x — **$785**

4" PHOTO-EQUATORIAL ($89.00 Down) with eyepieces for 250x, 214x, 167x, 120x, 83x, 60x, 38x — **$890**

4" EQUATORIAL with clock drive **($98.50 Down)**, Model 160V, eyepieces as above — **$985**

4" EQUATORIAL with clock drive and metal pier **($107.50 Down)**, Model 166V, eyepieces as above — **$1075**

4" PHOTO-EQUATORIAL with clock drive and Astro-camera **($117.50 Down)**, with eyepieces for 250x, 214x, 167x, 120x, 83x, 60x, 38x, 25x — **$1175**

4" PHOTO-EQUATORIAL with clock drive, pier, Astro-camera **($128.00 Down)**, with 10 eyepieces — **$1280**

Note These UNITRON Features

- LENSES are FULLY CORRECTED for spherical and chromatic aberration and are COATED for maximum brilliance and clarity of image. Optics especially designed for astronomical observation. AIR-SPACED OBJECTIVES.
- REFRACTOR type of design duplicates the performance of larger telescopes of other types. No mirrored surfaces to become oxidized. Superior definition to the very edge of the field.
- EYEPIECES of the HIGHEST QUALITY: Orthoscopic, Achromatized Symmetrical, Kellner, Huygens. Three to ten eyepieces included with each instrument.
- FINEST MATERIALS used throughout. DURALUMIN TUBE. Moving parts of BRASS carefully machined to close tolerances, and finished in CHROMIUM. No war-surplus components used.
- MODERN DESIGN based on time-tested engineering principles. HANDSOME APPEARANCE to which no illustration can do justice.
- EQUATORIAL MODELS have slow-motion controls for **both** declination and right ascension as well as convenient clamps on both axes. Sturdy TRIPOD (or PIER).
- ALTAZIMUTH MODELS have slow-motion controls for **both** altitude and azimuth as well as clamps for both co-ordinates. Sturdy TRIPOD.
- VIEW FINDER with crosshair eyepiece.
- RACK-AND-PINION FOCUSING.
- Choice of UNIHEX Rotary Eyepiece Selector or STAR DIAGONAL and ERECTING PRISM for TERRESTRIAL OBSERVATION. The same complete range of terrestrial magnifications as for celestial observation.
- SUNGLASS for solar observation.
- ADDITIONAL ACCESSORIES available to add further to your observing pleasure.

Higher- and lower-power eyepieces available for all models. Prices include basic accessories, tripod and mounting, fitted wooden cabinets, and operating instructions.

New UNITRON View Finders

UNITRON's popular view finders with newly designed optics and mechanical features are better than ever. From left to right: 23.5-mm., 30-mm., 42-mm.

1. **VIEW FINDER** (Used on UNITRON 2.4" Equatorials): 23.5-mm. (.93") achromatic objective, 6x eyepiece with crosshairs. Chromed brass tube. Mounting brackets with centering screws. **Only $8.50 postpaid**

2. **VIEW FINDER** (As used on UNITRON 3" Refractors): 30-mm. (1.2") coated achromatic objective and 8x eyepiece with crosshairs. Other details as in View Finder 3. **Only $10.75 postpaid**

3. **VIEW FINDER** (As used on UNITRON 4" Refractors): 42-mm. (1.6") coated achromatic air-spaced objective. 10x eyepiece with crosshairs. Duralumin tube finished in white enamel. Dewcap. Furnished with mounting brackets, centering screws for collimation, and mounting screws. This finder measures approximately 16" overall. It is light in weight, compact and small enough for use as a hand telescope furnishing spectacular wide-field views of the sky. **Only $18.00 postpaid**

UNITRON INSTRUMENT DIVISION OF UNITED SCIENTIFIC COMPANY
204-206 MILK STREET · BOSTON 9, MASSACHUSETTS

Get UNITRON's FREE Observer's Guide and Catalog on ASTRONOMICAL TELESCOPES

This valuable 38-page book is yours for the asking!

With artificial satellites already launched and space travel almost a reality, astronomy has become today's fastest growing hobby. Exploring the skies with a telescope is a relaxing diversion for father and son alike. UNITRON's handbook contains full-page illustrated articles on astronomy, observing, telescopes and accessories. It is of interest to both beginners and advanced amateurs.

Contents include—

- Observing the sun, moon, planets and wonders of the sky
- Constellation map
- Hints for observers
- Glossary of telescope terms
- How to choose a telescope
- Amateur clubs and research programs

UNITRON
INSTRUMENT DIVISION of UNITED SCIENTIFIC CO.
204-206 MILK STREET • BOSTON 9, MASS.

Please rush to me, free of charge, UNITRON's new Observer's Guide and Telescope Catalog 20-W.
Name _____
Street _____
City _____ State _____



SOUTHERN STARS

The sky as seen from latitudes 20° to 40° south, at 11 p.m. and 10 p.m., local time, on the 6th and 22nd of January, respectively; also, at 9 p.m. and 8 p.m. on February 6th and 21st. For other dates, add or subtract ½ hour per week.

Fifteen 1st-magnitude stars are above the horizon at latitude 30° south at chart time. Fomalhaut is just setting in the southwest, while Regulus is climbing the sky in the east. Orion, a summer constellation south of the Tropic of Capricorn, is almost on the meridian.

STARS FOR NOVEMBER

The sky as seen from latitudes 30° to 50° north, at 9 p.m. and 8 p.m., local time, on the 7th and 22nd of November, respectively; also, at 7 p.m. and 6 p.m. on December 7th and 22nd. For other dates, add or subtract ½ hour per week.

Running from east to west and crossing overhead this month, the northern Milky Way is an inspiring sight. Locate the bright constellations in it: Auriga, Perseus, Cassiopeia, Cygnus, and Aquila. Try also Lacerta, little Delphinus, Sagitta, and the faint stars of Vulpecula.

©

HOW TINSLEY DEVELOPED TWO TELESCOPES IN ONE FOR B. Y. U.

When Brigham Young University's Astronomy Department assigned Tinsley Laboratories to design and manufacture a new telescope for the University, they asked for an instrument that would perform two distinct jobs: 1) student instruction and visual use; 2) wide field photography for advanced research.

Tinsley combined both functions in a single powerful instrument for BYU: a custom-designed 24-inch reflector telescope with a Baker photographic corrector.

The basic Newtonian-Cassegrain design with a 24-inch-diameter f/4 main mirror fulfills BYU's requirements for visual instruction. The Cassegrain, which is f/15, has complete skylight shielding.

With the Baker photographic corrector the telescope performs its second job with unusual versatility. The corrector permits photography of a seven degree field—a picture equal in width to 14 full moon photos laid side-by-side!

BYU's telescope, now in operation at Provo, Utah, is an example of the precision craftsmanship and exacting specifications which have built Tinsley's reputation as the leader in custom optical instrumentation.

TINSLEY
LABORATORIES, INC.
2526 Grove Street, Berkeley 4, Calif.



are FOREMOST in our production

We point with pride to the fact that we have installed more than fifty major and dozens of smaller ASTRO-DOMES in most of the United States, the Arctic, Antarctic, and five foreign lands. In addition, ASTRO-DOME has developed a whole new concept of dome design, from the rotating mechanism to the labyrinth seal shutters, which are now standard on all ASTRO-DOMES. Our engineering department has recently completed exciting new designs which are now in production. Write and find out how ASTRO-DOME can meet your requirements in the new space age.



write for Brochure today

ASTRO-DOME
INCORPORATED

MANUFACTURERS OF ASTRONOMICAL OBSERVATORY DOMES
1801 BROWNLEE AVE. N. E. CANTON 5, OHIO Phone GL. 4-2755

PRINTED BY WELLESLEY PRESS, INC.

Johnny isn't watching the late show tonight . . .

The shadows that shiver and shake on the TV screen are shivering and shaking in somebody else's living room tonight. Johnny has discovered something new.

He's traded the fleeting, flickering "thrills" of the 24 inch screen for the timeless excitement and majesty of the night sky.

He's traded the nervous rattle of the private eye's gun for a ringside seat at the stupendous nightly fireworks in the heavens.

He has, in short, discovered astronomy.

Nothing better could happen than what happened to Johnny. And it happened simply because someone took the trouble to awaken, nourish and satisfy a lifetime of curiosity in Johnny by making him the gift of a fine telescope.

Someone, not so long ago, gave Johnny a Unitron.

Johnny abandoned his 24" screen for this 2.4" Unitron refractor, complete with its handy, handsome, easily portable carrying case.

This is a close-up of the Unihex Johnny is using. It's 6 eyepieces in one, an exclusive with Unitron. One of a complete line of accessories to multiply your viewing pleasure.

UNITRON INSTRUMENT DIVISION OF UNITED SCIENTIFIC CO. — 204-205 MILK STREET, BOSTON 9, MASS.

SEE PAGES 62 AND 63 FOR MORE ABOUT UNITRON.